

pharmaniaga®



145mm

Germacort

Cream

Fusidic acid with Betamethasone

DESCRIPTION

A white smooth cream.

COMPOSITION

Each gram contains Fusidic acid 2% w/w and
Betamethasone (as - 17 valerate) 0.1% w/w.

PHARMACODYNAMICS

Fusidic acid is potent topical antibacterial agent.

Fusidic acid is very active against staphylococci, notably *S.aureus* and *S.epidermis*. *Nocardia osteroides* and many clostridial strains are also highly susceptible. Most Gram-negative bacteria are intrinsically resistant but fusidic acid is active against *Neisseria* spp. and *Bacteroids fragilis*. It has some activity against strains of *Mycoba cterium tuberculosis* is highly active against *M. leprae*.

Betamethasone valerate is an anti-inflammatory and anti-pruritic agent.

PHARMACOKINETICS

Fusidic acid penetrates normal or inflamed skin and achieves bactericidal concentrations down to the lower

corium. Fusidic acid is widely distributed into tissues and body fluids, including bone, pus and synovial fluid; it penetrates cerebral abscesses but does not enter cerebrospinal fluid in appreciable amounts. It has been found in the foetal circulation and in breast milk. About 95% or more of fusidate in circulation is bound to plasma protein. Fusidic acid is excreted in the bile, almost entirely as metabolites some of which have weak antimicrobial activity. About 2% appears unchanged in the faeces. Little is excreted in the urine or removed by haemodialysis.

Betamethasone similar to other corticosteroids is absorbed systemically across the stratum corneum, increasing skin hydration, e.g. with occlusive dressings primarily increases penetration. Corticosteroids are metabolised primarily in the skin, or once absorbed systemically, in the liver. Repeated applications result in a cumulative depot effects in the skin, which may lead to a prolonged duration of action, increased side-effects and increased systemic absorption.

INDICATIONS

Inflammatory dermatoses where bacterial infection is present or likely to occur, e.g. atopic eczema, discoid eczema, stasis eczema, seborrheic dermatitis, lichen simplex chronicus, psoriasis, discoid lupus erythematosus.

CONTRAINDICATIONS

Topical corticosteroids are contraindicated in viral, tuberculous and fungal skin infections, perioral dermatitis, acne rosacea and ulcerative conditions. Fusidic acid is contraindicated in infections caused by non-susceptible organisms, in particular, *Pseudomonas aeruginosa*. This product is also contraindicated in known hypersensitivity to its components and in infants under 1 year old.

ADVERSE REACTIONS

Hypersensitivity reactions in the form of rashes, mild stinging and irritation may occur. As with other topical corticosteroids, prolonged and intensive treatment may cause local atrophic changes of the skin, such as striae, thinning and dilatation of the superficial blood vessels, particularly when occlusive dressing is used or when skin folds are involved. Infants and children are particularly at risk as systemic absorption, sufficient to produce hypercorticism/adrenal suppression, can occur with prolonged or extensive use, especially when used under occlusion/napkin.

Caution should be exercised if a topical steroid is used near the eye. Glaucoma might result if the preparation enters the eye.

Hypersensitivity reactions to fusidic acid are rare. However, if signs of hypersensitivity occur, treatment should be withdrawn.

WARNINGS

This product is not to be used in the eye as there is a potential risk of glaucoma. Dermal steroids should not be used in large amounts or for prolonged periods of time in pregnancy; foetal abnormalities have been seen in animals due to systemic absorption.

PRECAUTIONS

When steroids, particularly fluorinated steroids are applied for long period of time (more than 4 weeks), atrophic striae is likely to occur. Systemic absorption resulting in adrenal

100mm

145mm

suppression may occur especially under occlusion with weekly doses of over 30 g.

Prolonged use on flexures and intertriginous areas should be avoided.

Use in children

This product is contraindicated in children less than 1 year old. In infants, the napkin may acts as an occlusive dressing and increase absorption. Adrenal suppression is more likely to occur in infants and children.

PREGNANCY

Fusidic acid:

No effects during pregnancy are anticipated, since systemic exposure to fusidic acid is negligible. Studies in animals have not shown teratogenic effects with fusidic acid. Limited studies in animals have shown negligible systemic absorption of topical fusidic acid.

Betamethasone valerate:

There are no or limited amount of data from the use of topical betamethasone valerate in pregnant women. Studies in animals have shown reproductive toxicity / foetal abnormalities.

Germacort should not be used during pregnancy unless clearly necessary.

LACTATION

Germacort can be used during breast-feeding but should not be applied on the breasts to avoid accidental ingestion by the infant.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Germacort cream has no or negligible influence on the ability to drive or to use machines.

DRUG INTERACTIONS

No relevant information available.

DOSAGE AND ADMINISTRATION

Uncovered lesions : Apply 2-3 times a day.

Covered lesions : Less frequent applications may be adequate.

OVERDOSAGE

Topically applied corticosteroids can be absorbed in sufficient amounts to produce systemic effects (see adverse reactions). Since there is no specific antidote available, treatment is symptomatic, supportive and consists of discontinuance of topical corticosteroid therapy. Gradual withdrawal of the preparation may be necessary.

ROUTE OF ADMINISTRATION

Topical.

STORAGE CONDITIONS

Store below 30°C.

Protect from light.

Keep container tightly closed.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

Tube of 5g.

Date of revision: 20th March 2024

PRODUCT REGISTRATION HOLDER:

Pharmaniaga Manufacturing Berhad (198001006232)
No. 11A, Jalan P/1, Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi,
Selangor Darul Ehsan, Malaysia.

MANUFACTURER:

Idaman Pharma Manufacturing Sdn Bhd (200401023395)
Lot 24 & 25, Jalan Perusahaan Lapan,
Bakar Arang Industrial Estate,
08000 Sungai Petani, Kedah, Malaysia.

Lt-408.00

100mm