

pharmaniagaTM

Bromhexine



Tablet

DESCRIPTION

Green, round, flat, bevelled edge tablet with Pharmaniaga icon on one side and scoreline on the other.

COMPOSITION

Each tablet contains Bromhexine HCl 8 mg.

PHARMACODYNAMICS

Bromhexine HCL has been reported to change the structure of bronchial secretion and to increase volume and reduce the viscosity of sputum. It is absorbed from the gastrointestinal tract and peak plasma concentration has occurred after one hour. Most of the dose is excreted in the urine mainly as metabolites, only a small amount is excreted in the faeces.

PHARMACOKINETICS

Bromhexine Hydrochloride is absorbed from the gastrointestinal tract readily and peak plasma level occurs after an hour. Most of the dose is excreted in the urine, only a small amount is excreted in the faeces.

INDICATIONS

All forms of tracheobronchitis, emphysema with bronchitis, pneumoconiosis, chronic inflammatory pulmonary conditions, bronchiectasis, bronchitis and other conditions where tenacious mucous is a problem.

CONTRAINDICATION

Hypersensitivity to bromhexine.

ADVERSE REACTION/SIDE EFFECTS

Gastrointestinal side effect e.g. nausea and vomiting may occur occasionally and a transient rise in serum aminotransferase values has been reported.

Immune System Disorders

Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorder

Frequency not known: Severe skin reactions (including Steven Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

PRECAUTIONS

Bromhexine should be given with caution to patient with gastric ulceration.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as precaution.

DRUG INTERACTIONS

Bromhexine will increase the concentration of oxytetracycline in sputum and nasal mucous.

DOSAGE AND ADMINISTRATION

Adults and children over 10 years: 1-2 tablets three times daily.
Children 5-10 years: 1/2 tablets three times daily.

ROUTE OF ADMINISTRATION

Oral.

OVERDOSAGE

The toxicity associated with expectorant drugs like bromhexine is not reported.

STORAGE AND CONDITION

Store below 30°C.

Protect from light.

Keep container tightly closed.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

In bottles of 180 tablets.

Date of revision: 3rd September 2018

PRODUCT REGISTRATION HOLDER/

MANUFACTURER:

PHARMANIAGA MANUFACTURING BERHAD

(60016-D)

11A, Jalan P/1, Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi, Selangor Darul Ehsan,
Malaysia.

XXXXXXXXXX