

pharmaniaga™

Dextromethorphan

 Syrup

DESCRIPTION

Clear mixture, green in colour and with pineapple flavour.

COMPOSITION

Each 5 mL contains Dextromethorphan hydrobromide 15 mg.

ACTIONS AND PHARMACOLOGY

Dextromethorphan HBr has a depressant action on the cough reflex by a direct action on the cough center in the medulla of the brain. It has no analgesic or expectorant effect, no other depressant action on the central nervous system, nor any other established pharmacological action. Its use does not lead to addiction.

It is fully absorbed from the gastro-intestinal tract. It is metabolised in the liver to demethylated metabolites and dextorphan. The effectiveness in controlling pathological cough is as effective as codeine.

INDICATIONS

Treatment directed towards suppressed of cough is symptomatic, and has no direct effect on the

underlying disease. The aim is to decrease both the frequency and intensity of the cough.

CONTRAINDICATIONS

Productive cough should not be suppressed completely but only to reduce irritation and permit sleep. Cough persisting longer than 7-10 days requires medical diagnosis before symptomatic, treatment as the underlying disease may not be self-limiting (e.g. common cold) but of a more serious nature (e.g. lung cancer or congestive cardiac failure).

Cough suppressants may cause sputum retention and this may be harmful in patients with chronic bronchitis and bronchiectasis.

ADVERSE REACTIONS

Dextromethorphan HBr occasionally causes drowsiness, dizziness, excitation, mental confusion, and gastrointestinal disturbances, very high doses may produce respiratory depression.

PRECAUTIONS

Dextromethorphan should be administered with caution to patients with liver disease and to asthmatic patients.

Patients with mental depression or phobic-anxiety states who are being treated with monoamine oxidase inhibitors should also avoid dextromethorphan.

WARNING:

- Not suitable to be used for children below 2 years of age.
- Use carefully and follow the advice from doctor or pharmacist for children aged 2 to 6 years.

Pregnancy

Problems in humans have not been documented.

Lactation

It is not known whether dextromethorphan is distributed into breast milk. However, problems in humans have not been documented.

DRUG INTERACTIONS

Dextromethorphan is incompatible with penicillins, tetracyclines, salicylates, phenobarbitone sodium and high concentrations of sodium or potassium iodides.

Concurrent use with CNS depression-producing medications may potentiate the CNS depressant

effects of these medications or dextromethorphan. Concurrent use with monoamine oxidase inhibitors, including furazolidone, pargyline and procabazine may cause excitation, hypotension and hyperpyrexia.

DOSAGE AND ADMINISTRATION

Adults and children 12 years of age or older, 1 tsp. every 4 hours, not to exceed 120 mg daily.

OVERDOSAGE

Seek medical attention immediately if the following symptoms occurs. Confusion, drowsiness or dizziness; severe nausea or vomiting; severe unusual excitement; nervousness; restlessness or irritability.

ROUTE OF ADMINISTRATION

Oral.

STORAGE CONDITIONS

Store below 30°C.
Protect from light.
Keep container tightly closed.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

PRESENTATION

In bottles of 120 mL.

Date of revision: 03 April 2026

PRODUCT REGISTRATION HOLDER:

Pharmaniaga Manufacturing Berhad
(198001006232)
No. 11A, Jalan P/1, Kawasan Perusahaan Bangi,
43650 Bandar Baru Bangi,
Selangor Darul Ehsan, Malaysia.

MANUFACTURER:

InspiraPharma Sdn. Bhd. (200401023395)
Lot 24 & 25, Jalan 8,
Kawasan Perusahaan Bakar Arang,
08000 Sungai Petani,
Kedah Darul Aman, Malaysia.

Lt-317.01