



ARTWORK

LEO Pharma A/S
Internal Market Access

Scale	Get-up	Material No	Sent by e-mail
100%	LK+MY	066232-XX	▼
Subject	INS 175 x 280 mm		Date
Colour	Black		Sign.
		OMADK	Sign.

Preparation Strength Packsize	Fucidin® H cream	Place of production	Ireland
Comments: Page 2 of 2			

IIE007-01 - 175 x 280 mm 175 mm

Fucidin® H can be used during pregnancy if clinically needed. However, based on a general knowledge about systemic corticosteroids, caution should be exercised when using Fucidin® H during pregnancy.

Topical corticosteroids should not be used extensively, in large amounts or for prolonged periods in pregnant patients, as fetal abnormalities have been seen in animals due to systemic absorption.

Breastfeeding

No effects on the breastfed new-born/infant are anticipated since the systemic exposure of topically applied fusidic acid/hydrocortisone acetate to a limited area of skin of the breastfeeding woman is negligible.
Fucidin® H can be used during breastfeeding but it is recommended to avoid applying Fucidin® H on the breast.

Fertility

There are no clinical studies with Fucidin® H regarding fertility.

Effects on Ability to Drive and Use Machines

Fucidin® H has no or negligible influence on the ability to drive or to use machines.

Undesirable Effects

The estimation of the frequency of adverse reactions is based on a pooled analysis of data from clinical studies and spontaneous reporting.

The most frequently reported adverse reactions during treatment are application site reactions including pruritus, burning and irritation.

Adverse reactions are listed by MedDRA system organ class (SOC) and the individual adverse reactions are listed starting with the most frequently reported. Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

Very common ≥1/10
Common ≥1/100 and < 1/10
Uncommon ≥1/1,000 and <1/100
Rare ≥1/10,000 and <1/1,000
Very rare <1/10,000

Immune system disorders	
Uncommon (≥1/1,000 and <1/100)	Hypersensitivity
Skin and subcutaneous tissue disorders	
Uncommon: (≥1/1,000 and <1/100)	Dermatitis contact Eczema (condition aggravated) Rash
General disorders and administration site conditions	
Common: (≥1/1,00 and <1/10)	Application site reaction (incl. pruritus, burning and irritation)

Systemic undesirable class effects of mild corticosteroids, like hydrocortisone, include adrenal suppression especially during prolonged topical administration (see Special warnings and precautions for use).
Raised intra-ocular pressure and glaucoma may also occur after topical use of corticosteroids near the eyes, particularly with prolonged use and in patients predisposed to developing glaucoma (see Special warnings and precautions for use).

Dermatological undesirable class effects of mild corticosteroids like hydrocortisone include:
Atrophy, dermatitis (incl. dermatitis contact, dermatitis acneiform and perioral dermatitis), skin striae, telangiectasia, rosacea, erythema, depigmentation, hypertrichosis and hyperhidrosis. Ecchymosis may also occur with prolonged use of topical corticosteroids.

Class effects for corticosteroids have been uncommonly reported for Fucidin® H as described in the frequency table above.

Paediatric population

The observed safety profile is similar in children and adults (see Special warnings and precautions for use).

Overdose

For topically applied fusidic acid, no information concerning potential symptoms and signs due to overdose administration is available. Cushing’s syndrome and adrenocortical insufficiency may develop following topical application of corticosteroids in large amounts and for more than three weeks.

Systemic consequences of an overdose of the active substances after accidental oral intake are unlikely to occur. The amount of fusidic acid in one tube of Fucidin® H does not exceed the oral daily dose of systemic treatment. A single oral overdose of corticosteroids is rarely a clinical problem.

Shelf Life and Storage Conditions

Do not store above 30°C.
See the expiry date printed on the carton.

This leaflet was last revised in Jan 2023

Manufactured by:
LEO Laboratories Ltd
285, Cashel Road, Crumlin, Dublin 12, Ireland
LEO Pharma A/S
2750 Ballerup – Denmark

LEO



410

0 66232 - X X

