

BETAVAL-N CREAM

DESCRIPTION:

White smooth homogenous mass

COMPOSITION:

Each gm of the cream contains

Betamethasone 17-Valerate (Micronised) 1 mg Neomycin (as sulphate) 5mg

PHARMACODYNAMICS:

Betamethasone valerate is an active topical corticosteroids which produces a rapid response in those inflammatory dermatoses that are normally responsive to topical corticosteroids therapy and is often effective in the less responsive conditions, e.g.: psoriasis. Neomycin Sulphate, an aminoglycoside, is a broad spectrum bactericidal antibiotic. It is actively transported across the bacterial cell membrane, binds to a specific receptor protein on the 30S subunit of bacterial ribosomes, and interferes with an initiation complex between mRNA and the 30S subunit, inhibiting protein synthesis. DNA may be misread, thus producing nonfunctional proteins. Polyribosomes are split apart and are unable to synthesize protein.

PHARMACOKINETICS:

Absorbed systemically across the stratum corneum. Stratum corneum penetration is primarily enhanced by increasing skin hydration and or temperature particularly when occlusive dressing are used. Increased percutaneous absorption also occurs when the skin or mucosa is abraded or inflamed, with prolonged use or with excessive/extensive use. Readily absorbed from large denuded, burned or granulating areas.

INDICATIONS:

Treatment of the following conditions where secondary bacterial infection is present, suspected or likely to occur: Eczema, including atopic, infantile and discoid eczemas: prurigo; psoriasis; neurodermatoses, including lichen simplex; lichen planus; seborrhoeic dermatitis, intertrigo, contact sensitivity reactions and may be used as an adjunct to systemic therapy in generalized erythroderma.

Betaval-N can also be used in the management of secondarily infected insect bites, sunburn, prickly heat, anal and vulval pruritus (see Contraindications).

CONTRAINDICATIONS:

Contraindicated in those who are allergic to corticosteroids and neomycin sulphate and other aminoglycosides. Contraindicated in rosacea, acne and perioral dermatitis. Contraindicated in skin infections caused by viruses and fungi. Should not be used for the treatment of otitis externa when the ear drum is perforated, because of the risk of ototoxicity. Contraindicated in infants under 1 year.

SIDE EFFECTS:

The incidence and severity of the side effects increase with factors that increase percutaneous absorption. Prolonged and intensive treatment may cause local atrophic changes in the skin, eg. striae, thinning and dilatation of the superficial blood vessels, particularly when occlusive dressing are used or when skin folds are involved. Prolonged use of large amounts or treatment of extensive areas can result in sufficient systemic absorption to produce the features of hypercorticism. This effect is more likely to occur in infants and children and if occlusive dressings are used. Percutaneous absorption of topical corticosteroids has resulted in systemic effects such as hyperglycemia, glycosuria, and hypothalamic-pituitary-adrenal (HPA) axis suppression. There are reports of pigmentation changes and hypertrichosis with topical steroids. In rare instances, treatment

of psoriasis with corticosteroid preparations or its withdrawal is thought to have provoked the pustular form of the disease. Contact dermatitis and rarely auditory ototoxicity may occur.

PRECAUTIONS / WARNINGS:

Long-term continuous topical therapy should be avoided where possible, particularly in premature infants, infants and children, as adrenal suppression, Cushing's syndrome, intracranial hypertension and growth retardation can occur. If bacterial infection persists or any spread of infection occurs, withdraw the topical therapy. Bacterial infection is encouraged by the warm, moist conditions induced by occlusive dressing and the skin should be cleaned before a fresh dressing is applied. Avoid contact with the eyes. Gradual withdrawal of therapy by decreasing frequency of application may be necessary to avoid rebound flare-up of certain conditions such as psoriasis. Use of topical antibacterials may lead to skin sensitization, resulting in hypersensitivity reactions with subsequent topical or systemic use of the medication. The use of Neomycin may be associated with a theoretical risk of fetal toxicity.

Warning:

This product contains sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphite.

INTERACTIONS WITH OTHER MEDICAMENTS:

Co-administered drugs that can inhibit CYP3A4 (e.g. ritonavir, itraconazole) have been shown to inhibit the metabolism of corticosteroids leading to increased systemic exposure. The extent to which this interaction is clinically relevant depends on the dose and route of administration of the corticosteroids and the potency of the CYP3A4 inhibitor. Following significant systemic absorption, neomycin sulphate can intensify and prolong the respiratory depressant effects of neuromuscular blocking agents. Possibility of cumulative toxicity should be considered when neomycin sulphate is applied topically in combination with systemic aminoglycoside therapy

PREGNANCY AND LACTATION:

There is inadequate evidence of safety in human pregnancy.

It should not be applied to the breasts prior to nursing

ROUTE OF ADMINISTRATION:

Topical administration

DOSAGE & ADMINISTRATION:

A small quantity should be applied to the affected area 2 or 3 times daily until improvement occurs.

Maintenance: Apply once a day. If no improvement is seen within 2-4 weeks, reassessment of the diagnosis may be necessary.

OVERDOSAGE & TREATMENT:

Symptoms

: No specific symptoms are reported.

Treatment

: Since there is no specific antidote available, treatment is symptomatic and supportive and consists of discontinuance of therapy.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES:

Not applicable

PRESENTATION:

Aluminium tubes of 5g, 10g and 15g.

STORAGE CONDITIONS:

Store in a cool dry place below 30°C.

Keep containers tightly closed.

Protect from light and freezing.

Keep medicine out of reach of children.

MANUFACTURED BY:

Noripharma Sdn. Bhd. 200701034604 (792633-A)

Lot 5030, Jalan Teratai,

5 ½ Mile Off Jalan Meru,

41050 Klang,

Selangor

DATE OF REVISION:

August 2021