



DUZEN CREAM

Description: White colour cream

Content:

Fusidic Acid2.0%w/w

Preservative: Chlorocresol 0.1%w/w

Indication:

For the treatment of skin infections caused by staphylococci, streptococci (particularly due to penicillin sensitive strains), *Corynebacterium minustissimum* and other fusidic acid sensitive organisms e.g. in impetigo, boils, carbuncles, infected wounds, folliculitis, paronychia, sycosis barbae, hidradenitis and erythrasma.

Pharmacology:

Fusidic acid is a steroidal antibiotic with a bacteriostatic or bactericidal (in high concentrations) activity mainly against Gram+ve bacteria. Nearly all strains of *Staphylococcus aureus* are inhibited by very low concentrations of fusidic acid. It also has antiprotozoal properties and immunosuppressant actions. Fusidic acid inhibits bacterial protein synthesis, but in contrast to agents such as the macrolides or tetracyclines, it does not bind to the bacterial ribosome but inhibits a factor necessary for translocation of peptide subunits and elongation of the peptide chain. It is capable of inhibiting protein synthesis in mammalian cells but exerts a selective action against susceptible infecting organisms because of poor penetration of the host cell. Fusidic acid is very active against staphylococci (*Staph aureus* and *Staph epidermidis*), *Nocardia asteroides*, *Neisseria spp*, *Bacteroides fragilis*, *Mycobacterium tuberculosis* and *Mycobacterium leprae*, and many other clostridial strains.

Topically administered doses penetrate the skin and the extent is high when applied with dimethyl sulphoxide. Widely distributed throughout the body but does not enter the cerebrospinal fluid. It crosses the placenta and is secreted in the milk. Protein binding is about 95%. About 25% of a dose is excreted unchanged in the feces.

Contraindications:

It is contraindicated in infections caused by non-susceptible organisms such as *Pseudomonas aeruginosa* and patients hypersensitive to fusidic acid and propylene glycol.

Side effects / Adverse effects:

Low frequency of hypersensitivity reactions in the form of rashes and irritation. Topical use may lead to problems of resistance.

Precautions:

Safety for use in pregnancy and lactation have not been established. Risk benefits should be considered. Caution when using topical treatment near the eyes as it may cause irritation.

Drug Interactions

Nil for topical application.

Dosage and Administration

Apply to affected area 2 - 3 times daily for 7 days.

Treatment of overdose

Rashes and irritation may occur. Treatment is symptomatic and supportive.

Storage Conditions : Keep container tightly closed.
Store in a cool, dry place at 30°C. Protect from light.

Presentation : 5gm/tube

Shelflife : 3 years

Manufactured & : PRIME PHARMACEUTICAL SDN. BHD.

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia