



TRIDINE SYRUP

COMPOSITION:

Each 5ml contains
 Triprolidine HCl 1.25mg
 Pseudoephedrine HCl 30mg

Methylparaben 0.1% w/v as preservative.

PRESENTATION:

A cherry red coloured and fruity flavoured syrup.

INDICATIONS:

Axcel Tridine Syrup is indicated for symptomatic relief of nasal and respiratory congestion, common cold, acute sinusitis, allergic rhinitis, vasomotor rhinitis.

PHARMACOLOGY:

Mechanism of Action:

Triprolidine Hydrochloride is a histamine H₁-receptor antagonist. It acts by competing with histamine for H₁ receptor sites on the effector cells, to the exclusion of agonist molecules, without itself initiating a response. This antagonism is competitive and reversible. It does not inactivate or prevent the production of histamines.

Pharmacokinetics:

Pseudoephedrine is a direct and indirect acting sympathomimetic agent and is an effective upper respiratory tract decongestant. It is commonly combined with other ingredient in preparations intended for the relief of cough and cold symptoms.

DOSAGE & ADMINISTRATION:

For oral administration only.

To be taken 3 times daily.

Children:

6 - 12 years: 5ml

Adult and children above 12 years: 10ml

Do not use more than 5 days.

CONTRAINDICATION:

Contraindicated in patients with known hypersensitivity to antihistamines, propylene glycol and pseudoephedrine; severe hypertension or severe coronary artery disease and MAOI intake.

PRECAUTION:

Use with caution in patients with narrow angle glaucoma, diabetes mellitus, urinary retention, history of bronchial asthma, hyperthyroidism,

epilepsy, cardiovascular disease, prostatic enlargement, caution in severe renal or hepatic impairment. May cause drowsiness, avoid driving or operating machinery. Do not use in children under 6 years of age. Do not use for more than 5 days. If symptoms persist, please consult your doctor.

Use in Pregnancy and Lactation:

Use is not recommended in nursing mothers, pregnancy.

Use in Newborn / Premature:

Not recommended for newborn and premature infants.

SIDE EFFECTS:

The most common side effects is sedation including inability to concentrate, lassitude, dizziness, hypotension, muscular weakness, and incoordination. Sedative effects, when they occur, may diminish after a few days. Other side effects include gastrointestinal disturbances, such as nausea, vomiting, diarrhoea or constipation, colic and epigastric pain. It may also produce headache, blurred vision, tinnitus, elation or depression, irritability, anxiety, restlessness, tachycardia, palpitations, tremor, insomnia, confusion, anorexia, difficulty in micturition, urinary retention, dyspnoea, dryness of the mouth, tightness of chest.

DRUG INTERACTIONS:

Axcel Tridine Syrup may enhance the sedative effects of CNS depressants including alcohol, narcotic analgesics, sedatives and tranquilisers. Interact with MAO inhibitors, antihypertensives, cardiac glycosides, furazolidone, sympatholytics, quinidine, or tricyclic antidepressants. It is also important to note that it may mask the warning signs of damage caused by ototoxic drugs such as aminoglycoside antibiotics.

OVERDOSAGE AND TREATMENT:

Overdosage is associated with antimuscarinic, gastrointestinal, and CNS effects. In children CNS stimulation predominates over CNS depression, causing ataxia, excitement. Tremors, psychoses, hallucinations, and convulsions; hyperpyrexia may also occur. Deepening coma and cardiorespiratory collapse may follow.

Treatment for overdosage:

If the drug has been taken recently by mouth, the stomach should be emptied by aspiration and lavage. Emetics are generally of little value. The patients should be kept quiet to minimise the excitation which occurs particularly in children. Convulsion and marked CNS stimulation may be controlled with diazepam. Severe hypotension may require fluid replacement, and assisted respiration may be necessary.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN JAUHI DARI KANAK-KANAK

PACK QUANTITIES:

Available in 60ml & 100ml PET bottle and glass bottle of 120ml.

Further information can be obtained from pharmacist, physician or the manufacturer.

Product Registration Holder & Manufactured By:



Kotra Pharma (M) Sdn. Bhd.
 No.1, 2 & 3, Jalan TTC 12, (90082-V)
 Cheng Industrial Estate,
 75250 Melaka, Malaysia.

MAL20012739AZ

Regn - 280723