



DEXTROZINE SYRUP

COMPOSITION:

Each 5ml contains
Dextromethorphan Hydrobromide 15mg
Promethazine Hydrochloride 3.6mg

Methylparaben 0.1% w/v as preservative.

PRESENTATION:

Dark brown coloured syrup with sour plum flavour.

INDICATIONS:

Relief of congestion and dry irritating cough eg. those associated with common cold, upper respiratory tract infection & allergic rhinitis.

PHARMACOLOGY:

Mechanism of Action

Dextromethorphan is rapidly absorbed from gastrointestinal tract. It is metabolised in the liver and excreted in the urine as unchanged dextromethorphan and demethylated metabolites including dextrophan, which has some cough suppressant activity. Promethazine is well absorbed after oral or intramuscular administration. Peak plasma concentration have been observed 2 to 3 hours after administration by these routes, although there is low systemic bioavailability after oral administration, due to high first pass metabolism in the liver. Promethazine is widely distributed; it enters the brain and crosses the placenta. Values ranging from 76 to 93% have been reported for plasma-protein binding. Promethazine undergoes extensive metabolism, predominantly to promethazine sulfoxide, and also to N-desmethyl promethazine. It is excreted slowly via the urine and bile, chiefly as metabolites. Elimination half-life of 5 to 14 hours have been reported.

DOSAGE AND ADMINISTRATION:

To be taken 6 - 8 hourly.

Adults & Children > 12 years: 10ml (max. 40ml in 24 hours);

6 - 12 years: 5ml (max. 20ml in 24 hours);

2 - 6 years: 2.5ml (max. 10ml in 24 hours).

CONTRAINDICATION:

Contraindicated in infants less than one or two years of age, or patients with liver disease or with known hypersensitivity to dextromethorphan and other ingredients of the product.

PRECAUTION:

May cause drowsiness and hence impair ability to drive vehicles or operate machinery. Should not be given to patients with or at risks of developing respiratory failure, liver disease. Use with caution in patients with narrow angle glaucoma, hyperthyroidism, cardiovascular disease. Administration to patients receiving monoamine oxidase inhibitors may produce severe reactions such as hyperpyrexia and fatalities.

WARNING :

This preparation contains Sodium Metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites. This product contains Promethazine hydrochloride. It should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression. To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

Use in Pregnancy and Lactation:

Use is not recommended in nursing mothers, pregnant women.

Use in Children:

Use is not recommended in newborn, premature infants and children under 2 years of age.

SIDE EFFECTS:

Most common side effects include dizziness, drowsiness and gastrointestinal disturbances which may involve nausea, diarrhea and epigastric pain. It may also produce headache, blurred vision, dry mouth, urinary retention, constipation, increased gastric reflux, thickened respiratory tract secretions and tightness of the chest. In high doses, transient bradycardia may occur, followed by tachycardia with palpitation, arrhythmias and CNS stimulation.

DRUG INTERACTIONS:

Quinidine, CNS depressants including alcohol, barbiturates, hypnotic, narcotic analgesics. Sedatives and tranquillisers may enhance the CNS depression effects of dextromethorphan HBr. Severe reactions may occur if it is taken concomitantly with MAO inhibitors.

OVERDOSAGE AND TREATMENT:

Symptoms of overdosage may include excitation, confusion, respiratory depression.

Treatment for overdosage:

If the drug has been taken recently by mouth, the stomach should be emptied by emesis. A report by Shaul WL et. al. toxicity can be reversed with intravenous naloxone.

STORAGE:

Keep container well close. Store below 30°C. Protect from light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK**

PACK QUANTITIES:

Available in PET bottle of 60ml & 100ml and glass bottle of 120ml.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL20012740AZ

Product Registration Holder & Manufactured by :



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision : 31/10/23