

DYNA LAXATIVE PILL

MAL20014531XZ

DESCRIPTION:

TABLET

Colour : Brown
Shape : Round and Biconvex
Coating : Enteric-coated

CONTENT:

Each tablet contains:

Bisacodyl.....5 mg

PHARMACODYNAMICS:

Bisacodyl is a contact laxative acting directly on the colonic mucosa to produce normal peristalsis throughout the intestine where it stimulates sensory nerve to produce parasympathetic contraction of the colon.

PHARMACOKINETICS:

Following administration by mouth a small amount is absorbed and excreted in the urine as the glucuronide, but up to 38% of the dose has been stated to be excreted in the urine in some patients. About 3% of the glucuronide appear in the bile after about 10 hours. Bisacodyl is mainly excreted in the faeces.

INDICATION:

For the treatment of constipation and bowel evacuation before investigation procedures or surgery.

RECOMMENDED DOSE:

Children 6 years and above: 1 tablet.

Adults: 2 tablets.

To be taken before bedtime.

Do not chew the tablet. Swallow whole with a glass of water. Do not exceed the stated dose unless advised by a physician. Milk and antacid should not be taken within 1 hour.

ROUTE OF ADMINISTRATION:

FOR ORAL USE ONLY

CONTRAINDICATIONS:

Bisacodyl is contraindicated to patients hypersensitive to Bisacodyl. It should also not be given to patients with intestinal obstruction, abdominal pain, nausea or vomiting.

WARNINGS AND PRECAUTIONS:

Avoid prolonged use.

DRUG INTERACTIONS:

The concomitant use of antacids and milk products may reduce the resistance of the coating of the tablets and result in dyspepsia and gastric irritation.

The concomitant use of diuretics or adreno-corticosteroids may increase the risk of electrolyte imbalance. However, this situation only arises if excessive doses of Bisacodyl are taken.

PREGNANCY AND LACTATION:

There are no reports of undesirable or damaging effects during pregnancy or to the foetus attributable to the use of Bisacodyl. Nevertheless, medicines should not be used in pregnancy, especially the first trimester, unless the expected benefit is thought to outweigh any possible risk to the foetus.

Although the active ingredient of Bisacodyl is not known to be excreted in breast milk, its use during breast feeding is not recommended.

SIDE EFFECTS:

Bisacodyl can cause abdominal discomfort such as colic or cramps. Prolonged use can cause diarrhoea (with loss of water and electrolyte example calcium, potassium), muscular weakness and weight loss.

SYMPTOMS AND TREATMENT OF OVERDOSE:

Symptoms:

If high doses are taken diarrhoea, abdominal cramps and a clinically significant loss of potassium and other electrolytes can occur. This may also lead to increased sensitivity to cardiac glycosides.

Chronic overdose with laxatives may cause chronic diarrhoea, abdominal pain, hypokalaemia, secondary hyperaldosteronism and renal calculi.

Renal tubular damage, metabolic alkalosis and muscle weakness secondary to hypokalaemia have also been described in association with chronic laxative abuse.

Therapy:

Within a short time after ingestion of oral forms of Bisacodyl, absorption can be minimised or prevented by inducing vomiting. Otherwise, gastric lavage should be performed. Replacement of fluids and correction of electrolyte imbalance (particularly hypokalaemia) may be required. This is especially important in the elderly and the young. Administration of antispasmodics may be of some value.

PACKING/PACK SIZE(S):

Blister pack of 10x10's, 30x10's and 100x10's

**JAUHI UBAT DARIPADA KANAK-KANAK
KEEP OUT OF REACH OF CHILDREN**

MANUFACTURER/PRODUCT REGISTRATION HOLDER:

DYNAPHARM (M) SDN. BHD. 198001011897 (65683-V)

2497, Mk. 1, Lorong Perusahaan Baru 5,
Kawasan Perusahaan Perai 3,
13600 Perai, Pulau Pinang, MALAYSIA

PI2NT L001-03
Latest Revision: Jan 2022