

| Artwork Document Information                                                                                                                            |  | Artwork Type [ Please Tick / ]                                                                                                                                                                                                                                                                                                                                      |  | Sign / Stamp                         |  |
|---------------------------------------------------------------------------------------------------------------------------------------------------------|--|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--------------------------------------|--|
| Product : T ALLAVIN EXPECTORANT<br>SAP No. :<br>Date : 24 March 2025                                                                                    |  | <input type="checkbox"/> Unit Box<br><input type="checkbox"/> Outer Box<br><input type="checkbox"/> Label<br><input checked="" type="checkbox"/> Leaflet<br><input type="checkbox"/> Fix-A-Form<br><input type="checkbox"/> Shipper Carton<br><input type="checkbox"/> Aluminium Foil<br><input type="checkbox"/> Sachet<br><input type="checkbox"/> Others : _____ |  | <input type="checkbox"/> Generic Box |  |
| Document Category [ Please Tick / ]                                                                                                                     |  | Pharmacode / QR Code / Others Info                                                                                                                                                                                                                                                                                                                                  |  |                                      |  |
| <input checked="" type="checkbox"/> RA/NPRA Artwork<br><input type="checkbox"/> Internal Artwork Circulation<br><input type="checkbox"/> Others : _____ |  | <input type="checkbox"/> Recirculation Artwork<br><input type="checkbox"/> Previous Approved Artwork<br><input type="checkbox"/> Approval Copy Artwork<br><input type="checkbox"/> Draft Artwork<br><input type="checkbox"/> Clean Copy                                                                                                                             |  |                                      |  |
| Remarks                                                                                                                                                 |  | Colour                                                                                                                                                                                                                                                                                                                                                              |  |                                      |  |
| 1. Font : Arial Regular ( Spacing -8.3pt / Size - 7.5pt )<br>2. Dimension : 148.5mm(w) x 210mm(h)                                                       |  | Black                                                                                                                                                                                                                                                                                                                                                               |  |                                      |  |
| Colours shown on this proof are to indicate correct colour separations.                                                                                 |  |                                                                                                                                                                                                                                                                                                                                                                     |  |                                      |  |

210 mm

ALLAVIN  
EXPECTORANT

RELIEVES CHESTY COUGH

DESCRIPTION

Each 5 ml of Allavin Expectorant contains Diphenhydramine HCl 14mg and Ammonium Chloride 135mg. It is a green, prune flavoured syrup. Preservatives: Methylparaben: 0.1% w/v.

PHARMACOKINETICS

Diphenhydramine is well absorbed from the GI tract and is widely distributed throughout the body including the CNS. It crosses the placenta and is found in breast milk. It is mainly excreted in urine as metabolites. Ammonium Cl is also absorbed from the GI tract. The ammonium ion is converted into urea in the liver; the anion hence liberated into the blood stream and extracellular fluid causes a metabolic acidosis and decreases the pH of the urine; this is followed by transient diuresis.

INDICATIONS

For cough and complication associated with hypersensitivity and allergic manifestation of upper respiratory tract illness eg. common cold, flu and bronchitis.

PHARMACODYNAMICS

Diphenhydramine:

Frequently employed as antitussives in compound cough preparations, although it is not certain how they act in suppressing the cough reflex. Suggested mechanisms have included reduction in cholinergic nerve transmission, or cough suppression as a result of their sedative effects.

Ammonium :

Increases the volume of secretions in the respiratory tract and therefore facilitate their removal by ciliary action and coughing. It is also believed to achieve this by a reflex irritant effect on the gastric mucosa.

SIDE EFFECTS

The most common side effect of antihistamine is sedation, varying from slight drowsiness to deep sleep, and including lassitude, dizziness, and incoordination. Antihistamines may also produce antimuscarinic effects including dry mouth; thickened respiratory-tract secretions and tightness of the chest; blurred vision; urinary difficulty and retention; a reduction in tone and motility of the gastro-intestinal tract resulting in constipation and increased gastric reflux; in high doses, transient bradycardia followed by tachycardia with palpitations and arrhythmias; also in high doses, CNS stimulation. Ammonium chloride is irritant to the gastric mucosa and may produce nausea and vomiting particularly in large doses. Large doses may cause a profound acidosis and hypokalaemia which should be treated symptomatically.

DRUG INTERACTIONS

Allavin Expectorant may enhance the sedative effects of CNS depressants including alcohol, barbiturates, hypnotics, sedatives and tranquillisers. It may cause drowsiness and dulling of mental alertness. Patients undergoing treatment should not take charge of vehicles, other means of transport, machinery where loss of attention may lead to accident. Patients should abstain from alcohol.

PRECAUTIONS

Allavin Expectorant may enhance the sedative effects of CNS depressants including alcohol, barbiturates, hypnotics, sedatives and tranquillisers. It may cause drowsiness and dulling of mental alertness. Patients undergoing treatment should not take charge of vehicles, other means of transport, machinery where loss of attention may lead to accident. Patients should abstain from alcohol.

Pregnancy:

Adequate and well-controlled studies in humans have not been done.

WARNING

When used for treatment of cough and cold:

- Not to be used in children less than 2 years of age
- To be used with caution and doctor's/pharmacist's advice in children 2 to 6 years of age.

CONTRAINDICATIONS

Porphyria patients, hypersensitivity to any active ingredients and patients taking concomitant MAOI's. should be used with care in conditions such as epilepsy, prostatic hypertrophy, urinary retention, glaucoma and hepatic disease.

DOSAGE AND ADMINISTRATION

Adult and children above 12 years old - 10ml three or four times daily.  
Children 4-12 years old - 2.5 ml to 5 ml three or four times daily.  
Infants and children below 4 years old - As directed by physician.

SYMPTOMS AND TREATMENT OF OVERDOSE

Overdosage is associated with antimuscarinic, extrapyramidal, gastro-intestinal and CNS effects. In infants and children CNS stimulation predominates over CNS depression, causing ataxia, excitement, tremors, psychoses, hallucinations, and convulsions; hyperpyrexia may also occur. Deepening coma and cardiorespiratory collapse may follow. In adults, CNS depression is more common with drowsiness, coma, and convulsions, progressing to respiratory failure or possibly cardiovascular collapse. In severe overdosage the stomach should be emptied. Emetics may be tried if the patient is alert and there are no symptoms of toxicity. Other treatment is supportive and symptomatic and may include artificial respiration, external cooling for hyperpyrexia, and intravenous fluids.

PACK AND PRESENTATION

60ml and 90ml.

STORAGE CONDITIONS

Keep container tightly closed.  
Store in a dry place below 30°C.  
Protect from light.  
Keep out of reach of children.  
Jauhi daripada kanak-kanak.

Shelf life: Please refer to outer package.

Route of administration:

Oral

Product Registration Holder:

**Duopharma Manufacturing (Bangi) Sdn. Bhd.**  
Lot No. 2, 4, 6, 8 & 10, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

Manufacturer:

**Duopharma Manufacturing (Bangi) Sdn. Bhd.**  
Lot No. 2 & 4, Jalan P/7, Section 13, Bangi Industrial Estate, 43650 Bandar Baru Bangi, Selangor, Malaysia.

Revision Date :

March 2025

148.5 mm

