



DUZEN - B CREAM

Description: White colour cream

Content:

Fusidic acid 2.00%/w/w
Betamethasone 17 valerate (equiv to betamethasone 0.1%) 0.12%/w/w

Indications:

Treatment of inflammatory conditions due to bacterial infection e.g. eczema, seborrheic dermatitis and psoriasis.

Pharmacology:

Fusidic acid is a steroidal antibiotic with a bacteriostatic or bactericidal (in high concentrations) activity mainly against Gram +ve bacteria. Nearly all strains of Staphylococcus aureus are inhibited by very low concentrations of fusidic acid. It also has antiprotozoal properties and immunosuppressant actions. Fusidic acid inhibits bacterial proteins synthesis, but in contrast to agents such as the macrolides or tetracyclines, it does bind to the bacterial ribosome but inhibiting protein synthesis in mammalian cells but exerts a selective action against susceptible infecting organism because poor penetration of the host cell fusidic acid is very active against staphylococci (Staph. aureus and Staph. epidermidis), Nocardia asteroides, Neisseria spp., Bacteroides fragilis, Mycobacterium tuberculosis and Mycobacterium leprae, and many other clostridial strains. Betamethasone 17 valerate is a glucocorticoid applied topically in the management of various disorders in which corticosteroids are indicated. It has marked antipruritic, vasoconstrictor and anti-inflammatory activity. Topically administered fusidic acid doses penetrate the skin and the extent is increased when applied with dimethyl sulphoxide. From urinary excretion data, the half-life is 4 - 6 hours. It is widely distributed throughout the body but does not enter the cerebrospinal fluid; crosses the placenta and is secreted in the milk. Protein binding is about 95% bound to plasma proteins. About 2% of a dose is excreted unchanged in the feces. Betamethasone 17 valerate is well absorbed from the sites of local application, particularly under occlusive dressing or when the skin is broken. It may be absorbed to give systemic effects. It is more slowly metabolised in the skin and this tend to be systemically absorbed in a greater extent.

Contraindications:

Contraindicated in infections caused by non-susceptible organisms e.g. Pseudomonas aeruginosa, viral tuberculosis, fungal skin infections, perioral dermatitis, acne rosacea and ulcerative conditions. Also contraindicated in patients hypersensitive to fusidic acid, betamethasone 17 valerate, propylene glycol or any related components.

Sideeffects/Adverse reactions:

Low frequency of hypersensitivity reactions in the form of rashes and irritation. Topical use may lead to problems of resistance. When applied topically, particularly to large areas when the skin is broken or under occlusive dressings, betamethasone 17 valerate may be absorbed in sufficient amounts to cause systemic effects. With prolonged use, maceration of the skin, skin atrophy, striae and secondary infection may be present.

Precaution:

Safety for use in pregnancy and lactation has not been established. Risk benefits should be considered. Caution when using topical treatment near the eyes as it may cause irritation. Steroids applied topically over large areas, particularly when the skin is broken, with or without occlusive dressings may result in systemic effects. Therefore, suitable precautions should be taken. Special precautions have to be observed when prolonged use on infants and children. It should be used with caution on the face.

Dosage and administration:

Apply to the affected area 2 - 3 times daily.

Treatment of overdosage:

Rashes and irritation may occur. Excessive or prolonged use of topical corticosteroids can suppress pituitary-adrenal function, resulting in secondary adrenal insufficiency, and produce manifestations of hypercorticism, including Cushing's disease. In cases of chronic toxicity, slow withdrawal of steroids is advised. Appropriate antifungal or antibacterial therapy is indicated if overgrowth occurs. Treatment is symptomatic and supportive.

Storage conditions: Keep container tightly closed. Store in a dry place at 30°C. Protect from light. Keep away from children.

Presentation : 5gm/tube
Shelflife : 3 years

Product registration holder & Manufacturer: **PRIME PHARMACEUTICAL SDN. BHD.**

1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

30/07/2024