

LADOCORT®

Description : It is homogeneous white cream

Composition : LADOCORT® consists of 2 important drugs; 2% Miconazole Nitrate and 1% Hydrocortisone

Pharmacology/Pharmacokinetic :

Pharmacodynamics :

LADOCORT® has miconazole nitrate which is a synthetic antifungal agent that inhibits the growth of the common dermatophytes, the yeast-like fungus, and the organism responsible for tinea versicolor. LADOCORT® has also hydrocortisone as one of its active ingredients so it is anti-inflammatory and antipruritic too.

Pharmacokinetic :

Small amounts of Miconazole are absorbed systemically when the drug is applied intravaginally. Average peak serum concentrations of 4.2 mg/mL have been reported after 6 daily applications of a 14-day regimen for treatment of vulvovaginal candidacies; serum concentrations did not increase further during the remainder of the treatment period. Following intravaginal administration of a single dose of Miconazole (cream or suppository) to healthy women, about 1% of the dose is recovered in urine and feces. There has been no report to date indicating that Miconazole nitrate is absorbed following application to intact skin. Following application of hydrocortisone, which is in LADOCORT cream, to most areas of normal skin, only minimal amounts of the drug reach the dermis and subsequently. The systemic circulation; however, absorption is markedly increased when the skin has lost its keratin layer and can be increased by inflammation and/or diseases of the epidermal barrier (e.g., psoriasis, eczema). The drugs are absorbed to a greater degree from the scrotum, axial, eyelid, face, and scalp than from the forearm, knee, elbow, palm, and sole. Even after washing the area being treated, prolonged absorption of the corticosteroid occurs, possibly because the drug is retained in the stratum corneum.

Indications : LADOCORT® is indicated for the initial treatment of fungal infections which symptoms of pruritus or inflammation are prominent. Once the inflammatory symptoms have disappeared, treatment should be continued with Miconazole nitrate alone

Dosage & Usage : Apply LADOCORT® cream thinly to the infected lesion once to twice a day, than it should be continued with Miconazole nitrate alone should be instated after the symptoms of inflammation have disappeared.

Used in Elderly : Natural thinning of the skin occurs in the elderly; hence corticosteroids should be used sparingly and for short periods of time.

If after about seven days of treatment no improvement has occurred, cultural isolation of the infecting organism should be followed by appropriate local or systemic antimicrobial therapy.

Contraindication : Miconazole Nitrate preparation and topical Hydrocortisone therapy are contraindicated in patients with known hypersensitivity to the drug or any ingredient in the formulation. It is contraindicated in Tuberculous skin infections, herpes simplex, vaccine and all forms of varicella.

Warning and Precautions :

General Precautions : If a reaction suggesting sensitivity or irritation should occur, the treatment should be discontinued. Avoid the introduction of the cream into the eyes. Miconazole nitrate preparations should not come in contact with the eyes.

As with any topical corticosteroids, care is advised with infants and children when it is to be applied to extensive surface areas or under occlusive dressing including baby napkin. Similarly application to the face should be avoided. Patients receiving topical corticosteroids should be instructed to use the preparations only as directed by the physician, only for the disorder for which it was prescribed, and for no longer than the time period prescribed.

In infants, long term continuous topical corticosteroid therapy should be avoided. Adrenal suppression can occur even without occlusion.

Pediatric Used : Topical Miconazole nitrate preparations should be used in children younger than 2 years of age only under the direction and supervision of a physician. Use of the topical preparations in children 2-11 years of age should be supervised by an adult. Topical corticosteroid therapy in children should be limited to the minimum amount necessary for therapeutic efficacy.

Warning : This product contains Sodium Metabisulfite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulfites. For external use only. Avoid contact with the eyes.

In patients on warfarin, caution should be exercised and the anticoagulant effect should be monitored (see Interactions).

Drug Interaction : Miconazole enhances the anticoagulant effect to coumarin drugs. When it is used with these drugs, the anticoagulant effect should be monitored and dosage of the anticoagulant be adjusted accordingly. Severe hypoglycemia has occurred when Miconazole was used concomitantly with and oral sulfonylurea antidiabetic agent. It has been suggested that Miconazole may inhibit hepatic metabolism of these agents. Miconazole administered systemically is known to inhibit CYP2C9 enzyme system. Due to the limited systemic availability after topical application, clinically relevant interactions occur very rarely. In patients on warfarin which is subjected to metabolism by CYP2C9, caution should be exercised and the anticoagulant effect should be monitored (see Warnings and Precautions).

Pregnancy and Lactation : Caution is recommended during pregnancy and lactation. Treatment of large surfaces and the application under occlusive dressing should be avoided during that time.

Side Effect : The side effects are irritation; burning; maceration of the skin; allergic contact dermatitis; erythema; dryness; folliculitis; hypertrichosis; pruritus; acneiform eruptions; hypopigmentation; stinging and cracking/tightening of skin. As with all substances applied to the skin, an allergic reaction to any of the ingredients of the cream may occur.

Overdose : Hydrocortisone in LADOCORT® cream can be absorbed in sufficient amounts to produce systemic effects such as reversible HPA axis suppression, Cushing's syndrome, hyperglycemia and glycosuria. Recovery of HPA axis function is generally prompt and complete upon discontinuation of the drug.

Overdosage of Miconazole Nitrate in humans has not been reported to date. If accidental ingestion of large quantities of the product occurs, an appropriate method of gastric emptying may be used is considered to be necessary.

Storage conditions : Store below 30°C

Shelf-life : 4 years

Packing size : In tubes of 5 g and 15 g

Reg No. MAL20020820AZ

Date of revision : 19 April 2023

Manufactured by :
 **L.B.S. LABORATORY LTD., PART.**
602 Soi Panichanant, Sukhumvit 71 Road,
Khlong Tan Nuea, Wattana, Bangkok 10110, Thailand
Tel : (662) 392-4082 Fax : (662) 391-7122

Product Registration Holder :
HEALOL PHARMACEUTICALS SDN. BHD.
74-3, Jalan Wangsa Delima 6, KLSC Wangsa Maju,
53300 Kuala Lumpur, Malaysia
Tel : (60) 3-4142-7106, 4142-7107 Fax : (60) 3-4149-7108

Front

Back

Code: R 039/23
Project: LADOCORT (CREAM)
LF F&B
Date: 19-04-2023
Printed Size: 11x15 cm (WxH)
Special Colours: **K = 100**

R039-23_LADOCORT (CREAM)_LF F&B_ML_SM_2ndDraft_SM_19-04-2023