

pharmaniaga®

ALLERYL SYRUP 2MG/5ML

DESCRIPTION

Colourless to yellowish clear syrup.

**This product is presented as mobile liquid with pleasant pineapple odour and sweet, pleasant and slightly sourish after taste.*

COMPOSITION

Each 5mL of the syrup contains:

Chlorpheniramine Maleate 2 mg

Preservative: Sodium Benzoate 5 mg

PHARMACODYNAMICS

Chlorpheniramine Maleate is a potent antihistamine (H₁-antagonist).

Antihistamine diminish or abolish the actions of histamine in the body by competitive reversible blockade of histamine H₁-receptor sites on tissues. Chlorpheniramine Maleate also has anticholinergic activity.

Antihistamines prevent the release of histamine, prostaglandins and leukotrienes and shown to prevent the migration of inflammatory mediators. The actions of Chlorpheniramine Maleate include inhibition of histamine on smooth muscle, capillary permeability and reduction of oedema as well as wheal hypersensitivity reactions due to allergy and anaphylaxis.

PHARMACOKINETICS

Chlorpheniramine Maleate is well absorbed from the gastrointestinal tract, following oral administration. The effects develop within 30 minutes, are maximal within 1 to 2 hours and last 4 to 6 hours. The plasma half-life is estimated to be 12 to 15 hours. Chlorpheniramine Maleate is metabolised to monodesmethyl and didesmethyl derivatives. About 22% of an oral dose is excreted unchanged in the urine.

INDICATIONS

- 1) Symptomatic treatment of perennial and seasonal allergic rhinitis, vasomotor rhinitis and allergic conjunctivitis due to inhalant allergens and foods.
- 2) Symptomatic treatment of pruritus associated with allergic reactions and of mild, uncomplicated allergic skin manifestations of urticaria and angioedema.
- 3) May be used in combination with a decongestant (Pseudoephedrine) for the temporary relief of nasal and sinus congestion associated with the common cold and allergic rhinitis.

RECOMMENDED DOSAGE

Usual Paediatric Dose:

- 1) Children 2 to 5 years – 1 mg (or 2.5 mL) every four to six hours [maximum 6 mg (or 15 mL) daily].
- 2) Children 6 years to 12 years – 2 mg (or 5 mL) four to six hours, not to exceed 12 mg (or 30 mL) per day.

ROUTE OF ADMINISTRATION: Oral

CONTRAINDICATIONS

Alleryl Syrup 2mg/5ml is contraindicated in patients with known hypersensitivity to Chlorpheniramine Maleate or to any of the syrup ingredients.

Patients intolerant of other antihistamines or sympathomimetics (for e.g. amphetamines, epinephrine, ephedrine, isoproterenol, etc.) may be intolerant of this medication also.

The anticholinergic properties of Chlorpheniramine Maleate are intensified by monoamine oxidase inhibitors (MAOIs). Alleryl Syrup 2mg/5ml is therefore contraindicated in patients who have been treated with MAOIs within the last fourteen days.

WARNINGS AND PRECAUTIONS

Chlorpheniramine Maleate, in common with other drugs having anticholinergic effects, should be used with caution in epilepsy; raised intra-ocular pressure including glaucoma; prostatic hypertrophy; severe hypertension or cardiovascular disease; bronchitis, bronchiectasis or asthma; hepatic impairment; renal impairment. Children and the elderly are more likely to experience the neurological anticholinergic effects and paradoxical excitation (e.g. increased energy, restlessness, nervousness).

The anticholinergic properties of Chlorpheniramine Maleate may cause drowsiness, dizziness, blurred vision and psychomotor impairment in some patients which may seriously affect ability to drive and use machinery.

The effects of alcohol may be increased and therefore concurrent use should be avoided. Should not be used with other antihistamine containing products, including antihistamine containing cough and cold medicines.

Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

Alleryl Syrup 2mg/5ml contains 3 g of granular sugar per 5 ml. This should be taken into account in patients with diabetes mellitus.

Long term use increases the risk of dental caries and it is essential that adequate dental hygiene is maintained.

Use in New Born or Premature Infants: Alleryl Syrup 2mg/5ml is not recommended in new born or premature infants as this age group is at a higher risk than others, because of an increased susceptibility to antimuscarinic side effects such as central nervous system (CNS) excitation and increased tendency towards convulsions.

Use in Children and The Elderly: Caution is to be exercised when prescribing Alleryl Syrup 2mg/5ml to children and the elderly patients as they are more likely to experience the neurological anticholinergic effects.

When used in the treatment of cough and cold:

- a. Do not use for children below 2 years of age.
- b. Use with caution on the advice of doctor / pharmacist for children aged 2 - 6 years.

INTERACTIONS WITH OTHER MEDICAMENTS

The following drug interactions are significant:

- **CNS depressants** - Antihistamines may enhance the sedative effects of CNS depressants including alcohol, barbiturates, hypnotics, opioid analgesics, anxiolytics, sedatives and neuroleptics.
- **Monoamine Oxidase Inhibitors (MAOIs) Including Furazolidone, Pargyline And Procarbazine** - MAOIs may enhance the antimuscarinic effects of antihistamines and antihistamines have an additive antimuscarinic action with other antimuscarinic drugs, such as atropine and tricyclic antidepressants.
- **Phenytoin** - Concomitant administration of phenytoin sodium and Chlorpheniramine Maleate has been shown to cause drowsiness, ataxia, diplopia, tinnitus and episodes of occipital headache associated with vomiting. Chlorpheniramine Maleate might have delayed the hepatic metabolism of phenytoin thereby increasing the plasma concentrations.

PREGNANCY AND LACTATION

There is inadequate evidence of safety in human pregnancy. Alleryl Syrup 2mg/5ml should only be used during pregnancy when clearly needed and when the potential benefits outweigh the potential unknown risk to the fetus. Use during the third trimester may result in reaction in newborn or premature neonates.

Alleryl Syrup 2mg/5ml is not to be used during lactation unless considered essential by a physician because of the risk of adverse effects, such as unusual excitement or irritability as small amounts of Chlorpheniramine Maleate can be excreted in the breast milk.

ADVERSE EFFECTS AND UNDESIRABLE EFFECTS

- **Blood and lymphatic system disorders**
Unknown: haemolytic anaemia, blood dyscrasias.
- **Immune system disorders**
Unknown: allergic reaction, angioedema, anaphylactic reactions.
- **Metabolism and nutritional disorders**
Unknown: anorexia.
- **Psychiatric disorders**
Unknown: confusion*, excitation*, irritability*, nightmares*, depression.
- **Nervous system disorders***
Very common: sedation, somnolence.
Common: disturbance in attention, abnormal coordination, dizziness, headache.
- **Eye disorders**
Common: blurred vision.
- **Ear and labyrinth disorders**
Unknown: tinnitus.
- **Cardiac disorders**
Unknown: palpitations, tachycardia, arrhythmias.

- **Vascular disorders**
Unknown: hypotension.
- **Respiratory, thoracic and mediastinal disorders**
Unknown: thickening of bronchial secretions.
- **Gastrointestinal disorders**
Common: nausea, dry mouth.
Unknown: vomiting, abdominal pain, diarrhoea, dyspepsia.
- **Hepatobiliary disorders**
Unknown: hepatitis including jaundice.
- **Skin and subcutaneous disorders**
Unknown: exfoliative dermatitis, rash, urticaria, photosensitivity.
- **Musculoskeletal and connective tissue disorders**
Unknown: muscular twitching, muscle weakness.
- **Renal and urinary disorders**
Unknown: urinary retention.
- **General disorders and administration site conditions**
Common: fatigue.
Unknown: chest tightness.

**Children and the elderly are more susceptible to neurological anticholinergic effects and paradoxical excitation (e.g. increased energy, restlessness, nervousness)*

OVERDOSAGE

Symptom

The estimated lethal dose of Chlorpheniramine Maleate is 25 to 50 mg/kg body weight. Symptoms and signs include sedation, paradoxical excitation of the CNS, toxic psychosis, convulsions, apnoea, anticholinergic effects, dystonic reactions and cardiovascular collapse including arrhythmias.

Treatment

Symptomatic and supportive measures should be provided with special attention to cardiac, respiratory, renal and hepatic functions and fluid and electrolyte balance. If overdosage is by the oral route, treatment with activated charcoal should be considered provided there are no contraindications for use and the overdose has been taken recently (treatment is most effective if given within an hour of ingestion). Treat hypotension and arrhythmias vigorously. CNS convulsions may be treated with IV diazepam. Haemoperfusion may be used in severe cases.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

The anticholinergic properties of Chlorpheniramine Maleate may cause drowsiness, dizziness, blurred vision and psychomotor impairment, which can seriously hamper the patients' ability to drive and use machinery.

PACKING / PACK SIZE

Plastic container of 60ml.

STORAGE CONDITIONS

Keep container tightly closed in a dry place, below 30°C; Protect from light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

REGISTRATION NUMBER

MAL20021005AZ

CONTROLLED MEDICINE**UBAT TERKAWAL****KEEP MEDICINES OUT OF REACH OF CHILDREN****JAUHI UBAT-UBATAN DARI KANAK-KANAK**

For further information, please consult your doctor or your pharmacist.

Revision number : 02

Revision date : 17 Dec 2025

LI-002.02 PRODUCT REGISTRATION HOLDER AND MANUFACTURER

NSPIRAPHARMA SDN. BHD. (200401023395)

LOT 24 & 25, JALAN 8,
KAWASAN PERUSAHAAN BAKAR ARANG,
08000 SUNGAI PETANI,
KEDAH DARUL AMAN, MALAYSIA.