

105mm

Body as a whole: Headache, photosensitivity, asthenia.

Cardiovascular system: Hypotension, palpitations, tachycardia, arrhythmias

Digestive system: Constipation, diarrhoea, dry mouth and dry throat, dyspepsia, nausea, vomiting. Nausea and vomiting have been reported with high doses of ammonium chloride.

Nervous system: Agitation/ excitation, anxiety, confusion, convulsions, disturbed coordination, dizziness, hallucinations, insomnia, irritability, nervousness, paraesthesia, somnolence/ sedation, tremor, impaired performance (impaired driving performance, poor work performance, incoordination, reduced motor skills and impaired information processing), appetite stimulation, muscle dyskinesias and activation of epleptogenic foci. Somnolence was the most frequently reported adverse effects.

Paradoxical CNS stimulation may occur especially in children, with insomnia, nervousness, euphoria, irritability, tremors and rarely nightmares, hallucinations and convulsions.

Respiratory system: Dryness of nose, thickening of bronchial secretions, tightness of chest or throat, wheezing.

Skin: Pruritis, rash, urticaria.

Special senses: Dryness of the eyes, blurred vision, tinnitus.

Urogenital system: Urinary hesitancy and retention.

Adverse Drug Reactions Identified during Post Marketing Experience with Ammonium Chloride and Diphenhydramine Hydrochloride

Frequency	Adverse Event
Immune system disorders	
Very rare	Angioedema, hypersensitivity
Psychiatric disorders	
Very rare	Confusional state, hallucination, irritability, nervousness
Nervous system disorders	
Very rare	Agitation, coordination abnormal, convulsion, dizziness, headache, insomnia, paraesthesia, sedation, somnolence, tremor
Eye disorders	
Very rare	Vision blurred
Ear and labyrinth disorders	
Very rare	Tinnitus
Cardiac disorders	
Very rare	Palpitations, tachycardia
Vascular disorders	
Very rare	Hypotension
Respiratory, thoracic and mediastinal disorders	
Very rare	Chest discomfort, dry throat, nasal dryness
Gastrointestinal disorders	
Very rare	Abdominal pain, application site reaction, constipation, diarrhoea, dry mouth, dyspepsia, nausea, vomiting
Skin and subcutaneous tissue disorders	
Very rare	Pruritis, rash, urticaria

Renal and urinary disorders	
Very rare	Urinary retention
General disorders and Administrative site conditions	
Very rare	Asthenia

SYMPTOMS AND TREATMENT OF OVERDOSAGE

Overdosage of antihistamine may be fatal especially in infants and children.

Symptoms

For antihistamines:

In infants and children, CNS stimulation predominates over CNS depression, causing ataxia, excitement, tremors, psychoses, hallucinations and convulsion; hyperpyrexia may also occur. Deepening coma and cardiorespiratory collapse may follow.

In adults, CNS depression is more common with drowsiness, coma and convulsions, progressing to respiratory failure or possibly cardiovascular collapse. Large doses of ammonium chloride may cause a profound acidosis and hypokalaemia.

Treatment

There is no specific antidote for overdosage with antihistamines and Ammonium chloride, treatment is symptomatic and supportive until possible utilisation of the following:

- induction of emesis (syrup of ipecac is recommended, however, precaution against respiration is necessary especially in infants and children).
- gastric lavage if patient is unable to vomit within 3 hours of ingestion.
- saline cathartics are sometimes used.
- vasopressors to treat hypotension, however, epinephrine should not be used since it may further lower blood pressure.
- oxygen and intravenous fluids.
- precaution against use of stimulant because they may cause seizures.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

Diphenhydramine may cause drowsiness and may increase the effects of alcohol. Drowsiness may continue the following day. Those affected should not drive or operate machinery. Alcohol should be avoided.

PACKING / PACK SIZE

Plastic bottle of 90 mL

STORAGE CONDITIONS

Keep container tightly closed in a dry place, below 30°C; Protect from light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

REGISTRATION NUMBER

MAL20021006AZ

For further information, please consult your doctor or your pharmacist.

Revision number : 09

Revision Date : 11 Dec 2025

CONTROLLED MEDICINE

UBAT TERKAWAL

KEEP MEDICINES OUT OF REACH OF CHILDREN

JAUHI UBAT-UBATAN DARI KANAK-KANAK

LI-087.09 PRODUCT REGISTRATION HOLDER AND MANUFACTURER

INSPIRAPHARMA SDN. BHD. (200401023395)

LOT 24 & 25, JALAN 8

KAWASAN PERUSAHAAN BAKAR ARANG,

08000 SUNGAI PETANI,

KEDAH DARUL AMAN, MALAYSIA.

148mm