



## BEPROSALIC Lotion

**NAME AND STRENGTH OF ACTIVE SUBSTANCES:** Betamethasone dipropionate 0.064% w/v (equiv. to Betamethasone 0.05% w/v) and Salicylic acid 2% w/v.

**PRODUCT DESCRIPTION:** A colourless and slightly viscous solution.  
Other ingredients: Isopropyl alcohol, Methylparaben, Sodium hydroxide and Purified water.

**PHARMACODYNAMICS :** Betamethasone dipropionate is a topically active fluorinated corticosteroid which has anti-inflammatory, anti-pruritic and vasoconstrictive actions. Salicylic Acid has keratolytic properties, as well as bacteriostatic and fungicidal actions.

**PHARMACOKINETICS :** Topical corticosteroids have anti-inflammatory, anti-pruritic and vasoconstrictive actions. When administered topically, particularly under occlusive dressing or when the skin is broken sufficient corticosteroid may be absorbed to give systemic effects. Corticosteroids are extensively bound to plasma proteins. Only unbound corticosteroids has pharmacological effects or is metabolised. They are metabolised mainly in the liver, also in the kidney and are excreted in the urine. Salicylic Acid has keratolytic properties. It produces slow and painless destruction of the epithelium. When administered topically to large areas of the body, it may cause symptoms of acute systemic salicylate poisoning.

**INDICATIONS :** For the relief of the inflammatory manifestations of hyperkeratotic and dry corticosteroid-responsive dermatoses which include psoriasis, chronic atopic dermatitis, neurodermatitis and seborrhoeic dermatitis.

**CONTRAINDICATIONS :** Hypersensitivity to Betamethasone dipropionate, other corticosteroids, Salicylic Acid or other Salicylates or any component in the base. Topical corticosteroids are contraindicated in most viral infections of the skin and also tuberculosis and acne rosacea.

**DRUG INTERACTIONS :** Unknown.

**ADVERSE EFFECT :** Reported local adverse reactions associated with corticosteroids are burning, itching, irritation, dryness, folliculitis, hypertrichosis, acneiform eruptions, hypopigmentation, maceration of the skin, secondary infection, skin atrophy, striae and miliaria. Systemic adverse reactions, such as vision blurred, have also been reported with the use of topical corticosteroids. Salicylic Acids preparations may cause dermatitis.

**WARNINGS AND PRECAUTIONS :** Discontinue use if irritation, sensitivity, excessive dryness or other reactions develop. In the presence of infection, appropriate treatment should be given. Corticosteroids applied topically can be absorbed in sufficient amount to produce systemic effects like adrenal suppression, manifestation of Cushing's syndrome, hyperglycaemia and glucosuria in some patients, particularly in infants and children. Salicylic Acids in preparations applied to the skin can also be absorbed in sufficient amounts to produce salicylism. Therefore suitable precautions must be taken with Beprosalic Lotion during prolonged treatment, when treating extensive body surface areas, when using the occlusive technique and when treating children (because of a larger skin surface area to bodyweight ratio). Suitable precautions must be taken when treating stasis dermatitis and other skin diseases with impaired circulation. Discontinue use if irritation, sensitivity or other reactions develop and institute appropriate treatment. Safety of its use during pregnancy and lactation has not been established. Thus it

should be used only if the potential benefit justifies the potential risk to the foetus or nursing infant. Not suitable for ophthalmic use. Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma, or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids.

**ROUTE OF ADMINISTRATION :** Topical.

**RECOMMENDED DOSAGE :** Apply a few drops to the affected area two times daily.  
Maintenance : Apply less often.

**SYMPTOMS AND TREATMENT FOR OVERDOSE AND ANTIDOTE(S) :**

Symptoms: Excessive prolonged use of topical corticosteroids can suppress pituitary-adrenal functions resulting in secondary adrenal insufficiency. Excessive prolonged use of topical preparations containing Salicylic Acid may cause symptoms of salicylism.  
Treatment: Appropriate symptomatic treatment is indicated. Acute hypercortoid symptoms are virtually reversible. Treat electrolyte imbalance if necessary. In cases of chronic corticosteroid toxicity, slow withdrawal of steroids is advised. Treatment of salicylism is symptomatic. Measures should be taken to rid the body of salicylate. Administer oral sodium bicarbonate to alkalinize and force diuresis.

**PACKING :** 30ml Plastic bottle.

**STORAGE :** Keep containers well closed. Protect from light. Store below 30°C.  
For external use only. Keep out of reach of children.

**RECOMMENDED SHELF-LIFE:** 18 months.

**MANUFACTURED BY :**

HOE Pharmaceuticals Sdn. Bhd.  
Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman,  
42000 Port Klang, Selangor, MALAYSIA.

Revision date : January 2022