

Important information. Please read carefully.



Oral Solution

COMPOSITION

Each ml contains Cetirizine Hydrochloride 1 mg, Methyl Hydroxybenzoate 0.135% w/v and Propyl Hydroxybenzoate 0.015% w/v as preservative.

PHARMACODYNAMICS/ PHARMACOKINETICS

The principle effects of Cetirizine are mediated via selective inhibition of peripheral H₁ receptors. It is reported to have negligible anticholinergic and antiseroenergetic activity. Cetirizine is an antihistamine reported to be long acting and with some mast cell stabilizing activity. It inhibits the histamine-induced early phase of allergic reaction. It also inhibits the late phase recruitment of eosinophils, neutrophils and basophils associated with the allergic inflammatory response.

Cetirizine is rapidly absorbed from the gastro-intestinal tract after oral administration, with peak plasma concentration being attained in approximately 1 hour in adults. Food had no effect on the extent of cetirizine absorption but time to maximum concentration (T_{max}) was delayed by 1.7 hours and maximum concentration (C_{max}) was decreased by 23% in the presence of food. Cetirizine is highly bound to plasma proteins and has an elimination half-life of about 11 hours. The first-pass metabolism is reported to be low, and excretion is primarily through renal as unchanged drug.

INDICATION

Adezio is indicated for the relief of symptoms associated with seasonal allergic rhinitis and perennial allergic rhinitis in adults and children over 2 years of age such as sneezing, rhinorrhea, postnasal discharge, nasal pruritus, ocular pruritus, tearing and redness of the eyes.

Adezio is also indicated for the treatment of the uncomplicated skin manifestations of chronic urticaria in adults and children over 2 years of age. It significantly reduces the occurrence, severity and duration of hives and significantly reduces pruritus.

DOSEAGE AND ADMINISTRATION

Adezio is given as a single daily dose, with or without food. The time of administration may be varied to suit individual patient needs. To be administered orally.

Adults and Children over 12 Years: 5 or 10 mg depending on symptom severity.	1 teaspoon (5ml) or 2 teaspoons (10ml) once daily.
Children 6 to 11 Years: 5 or 10 mg depending on symptom severity.	1 teaspoon (5ml) or 2 teaspoons (10ml) once daily
Children 2 to 5 Years: 2.5mg; can be increased to a maximum dose of 5 mg per day depending on symptom severity and patient response.	½ teaspoon (2.5ml) once daily; can be increased to 1 teaspoon (5ml) once daily or ½ teaspoon (2.5ml) twice daily.

Dose Adjustment for Renal and Hepatic Impairment:

In patients 12 years old and older with decreased renal function (creatinine clearance 11-31 ml/min), patients on hemodialysis (creatinine clearance less than 7 ml/min), and in hepatically impaired patients, a dose of 5mg once daily is recommended. Similarly, pediatric patients aged 6 to 11 years with impaired renal or hepatic function should use the lower recommended dose. Because of the absence of pharmacokinetic and safety information for cetirizine in children below the age of 6 years with impaired renal or hepatic function, its use in this impaired patient population is not recommended.

CONTRAINDICATIONS

Adezio is contraindicated in those patients with a known hypersensitivity to it or any of its ingredients or hydroxyzine. It is also contraindicated during lactation.

PRECAUTIONS

Activities Requiring Mental Alertness: In clinical trials, the occurrence of somnolence has been reported in some patients taking Cetirizine, due caution should therefore be exercised when driving a car or operating potentially dangerous machinery. Central sedation or antimuscarinic effects do not generally occur with usual doses of Cetirizine.

Use in Pregnancy and Lactation

Teratogenic studies in animals have not demonstrated any special malformations. However as a precaution, Adezio should not be administered during the first trimester of pregnancy. Cetirizine has been reported to be excreted in human breast milk. Because many drugs are excreted in human milk, use of cetirizine in nursing mothers is not recommended.

Use in Paediatric

When used for treatment of cough and cold,

(a) Not to be used in children less than 2 years of age

(b) To be used with caution and doctor's/ pharmacist's advice in children 2 to 6 years of age.

DRUG INTERACTIONS

No clinically significant drug interactions have been reported with other drugs. However, caution should be exercised when other CNS depressants or alcohol is taken concurrently.

SIDE EFFECTS / ADVERSE REACTIONS

Most adverse reactions reported during therapy with cetirizine were mild or moderate. There have been occasional reports on headache, dizziness, drowsiness, agitation, dry mouth, gastrointestinal discomfort and hypersensitivity.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Overdosage has been reported with cetirizine. Symptoms reported include somnolence, restlessness, irritability and drowsiness. Should overdose occur, treatment should be symptomatic or supportive, taking into account any concomitantly ingested medications. There is no known specific antidote to cetirizine. Cetirizine is not effectively removed by dialysis, and dialysis will be ineffective unless a dialyzable agent has been concomitantly ingested.

SHELF-LIFE

The expiry date is indicated on the packaging.

STORAGE

Store in a cool place below 30°C.

KEEP ALL MEDICINES OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

DESCRIPTION

Clear, colourless oral solution with fruity flavour.

PACKING / PACK SIZES

Packs of 60 ml and 90ml in PET (polyethylene terephthalate) bottles.

Revision Date : 15-Aug-2023



Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd Company No. 8504-M 1-5, Cheng Industrial Estate, 75250 Melaka, Malaysia.

AD0200-004