



FLEMIN JUNIOR

EXPECTORANT

COMPOSITION:

Each 5ml contains
Diphenhydramine HCl 7mg
Ammonium Chloride 67.5mg

Sodium Benzoate 0.1% w/v as preservative.
Sodium Citrate 30mg / 5ml

PRESENTATION:

Brown color liquid with caramel flavor.

INDICATION:

Relief of cough and alleviation of nasal & bronchial congestion.

PHARMACOLOGY:

Mechanism of Action:

Diphenhydramine Hydrochloride is a histamine H₁-receptor antagonist. The antihistamine acts by competing with histamine for H₁ receptor sites on the effectors cells, to the exclusion of agonist molecules, without itself initiating a response. This antagonism is competitive and reversible. It does not prevent the production of histamines. Diphenhydramine Hydrochloride is readily absorbed from gastrointestinal tract and parental sites of administration. Systemic availability is however affected by high first pass metabolism. Following oral administration, its peak plasma concentration is within 1-4 hours. It is highly bound to plasma protein, thus widely distributed throughout the body including the CNS. As it crosses the plasma, amount may be detected in breast milk. Metabolism is extensive. Most of diphenhydramine is excreted as metabolites through urine with complete excretion achieved within 24 hours of administration. Ammonium Chloride is used as an expectorant. It absorbed from the gastrointestinal tract. The ammonium ion is converted into urea in the liver; the anion thus liberated into the blood stream and extracellular fluid causes a metabolic acidosis and decreases the pH of the urine; this is followed by transient diuresis.

DOSAGE & ADMINISTRATION:

For oral administration only.

To be taken 4 - 6 hourly.

Children 2 - 5 years : 5ml, not to exceed 40ml in 24 hours.

Children 6 - 12 years : 5 - 10ml, not to exceed 60ml in 24 hours.

Children above 12 years : 10 - 20ml, not to exceed 120ml in 24 hours.

CONTRAINDICATION:

Product is also contraindicated in patients who are hypersensitive to antihistamines or in patients who are receiving MAOI therapy and also the presence of impaired hepatic or renal function.

PRECAUTION:

May cause drowsiness and hence impair ability to drive vehicles or operate machinery. Use with caution in patients with narrow angle glaucoma, history of bronchial asthma, hyperthyroidism, cardiovascular disease and prostatic hypertrophy, stenosing peptic ulcer, pyloroduodenal obstruction, and bladder neck obstruction. It should be used cautiously, in patients with glucose-6-phosphate deficiency since they may induce hemolysis in presence of infection or stress. When used for treatment of cough and cold:

- a) Not to be used in children less than 2 years of age.
- b) To be used with caution and doctor's / pharmacist's advice in children 2 to 6 years of age.

Use in Newborn / Premature:

It should not be given to newborn or premature infants as this group of patients has an increased susceptibility to antimuscarinic effects.

Use in Children:

A paradoxical reaction characterised by hyperexcitability may occur in older children taking antihistamines.

SIDE EFFECTS:

The most common effect is sedation, varying from drowsiness to deep sleep, and including inability to concentrate, lassitude, dizziness, hypotension, muscular weakness, and incoordination. Sedative effects, when they occur, may diminish after a few days. Other side effects include gastrointestinal disturbances, such as nausea, vomiting, diarrhea or constipation, colic and epigastric pain. It may also produce headache, blurred vision, tinnitus, elation or depression, irritability, nightmares, anorexia, difficulty in micturition, dryness of the mouth, tightness of chest, tingling, heaviness and weakness of the hands. It may potentiate other CNS depressant compounds. Large doses may precipitate fits in epileptics. The ammonium salts content are irritant to the gastric mucosa and may produce nausea and vomiting particularly in large doses.

DRUG INTERACTIONS:

Additive effects if used with hypnotics, sedatives and tranquilizers. This preparation should not be given to patients receiving MAOIs, aminoglycosides, antibiotics and antimuscarinics. Avoid drinking alcoholic beverages when taking this syrup.

OVERDOSAGE & TREATMENT:

Symptoms of overdosage may vary from CNS depression including sedation, cardiovascular collapse, and stimulation

including insomnia, hallucination, tremors or convulsion. Other symptoms include dizziness, tinnitus, ataxia, blurred vision and hypotension.

Treatment For Overdosage:

If the drug has been taken recently by mouth, the stomach should be emptied by aspiration and lavage. Emetics are generally of little value. The patient should be kept quiet to minimise the excitation which occurs particularly in children. Convulsion and marked CNS stimulation should preferably be treated with diazepam or phenobarbitone intramuscularly; paraldehyde, thiopentone sodium, and chlorpromazine have also been recommended. Severe hypotension may require fluid replacement and assisted respiration may be necessary. Forced diuresis is of little value since antihistamine are rapidly metabolised and only traces are recovered in the urine. Large doses of Ammonium Chloride may cause a profound acidosis and hypokalemia which should be treated symptomatically.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN

JAUHI DARI KANAK-KANAK

PACK QUANTITIES:

Available in PET bottle of 60ml & 100ml and 120ml in amber glass bottle. Not all pack sizes may be marketed.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL20021157AZ

Product Registration Holder & Manufactured By :



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

Date of revision : 19/05/26