

For the use of a Registered Medical Practitioner or a Hospital or a Laboratory only OR for Specialist Use only

Generic Name

Telmisartan tablets 80 mg

Brand Name

Cresar 80

Composition

Each tablet contains:

Telmisartan Ph Eur.....80mg

Dosage Form

Uncoated Tablet

Product Description

White to off white, oval shaped tablets debossed with “D” on one side and plain on other side

Pharmacology

Pharmacodynamics

ATC Classification

C09CA07 - telmisartan; Belongs to the class of angiotensin II antagonists. Used in the treatment of cardiovascular disease.

Mechanism of Action

Pharmacology: Telmisartan is an orally effective and specific angiotensin-II receptor (type AT1) antagonist. Telmisartan displaces angiotensin II with very high affinity from its binding site at the AT1 receptor subtype, which is responsible for the known actions of angiotensin II. Telmisartan does not exhibit any partial agonist activity at the AT1 receptor. Telmisartan selectively binds the AT1 receptor. The binding is long lasting.

Telmisartan does not show affinity for other receptors, including AT2 and other less characterised AT receptors. The functional role of these receptors is not known, nor is the effect of their possible overstimulation by angiotensin II, whose levels are increased by telmisartan. Plasma aldosterone levels are decreased by telmisartan. Telmisartan does not inhibit human plasma renin or block ion channels. Telmisartan does not inhibit angiotensin-converting enzyme (kininase II), the enzyme which also degrades bradykinin. Therefore, it is not expected to potentiate bradykinin-mediated adverse effects.

In man, an 80 mg dose of telmisartan almost completely inhibits the angiotensin II evoked blood pressure increase. The inhibitory effect is maintained over 24 hours and still measurable up to 48 hours.

Treatment of Essential Hypertension: After the first dose of telmisartan, the antihypertensive activity gradually becomes evident within 3 hours. The maximum reduction in blood pressure is generally attained 4 weeks after the start of treatment and is sustained during long-term therapy.

The antihypertensive effect persists constantly over 24 hours after dosing and includes the last 4 hours before the next dose as shown by ambulatory blood pressure measurements. This is confirmed by trough to peak ratios consistently above 80% seen after doses of telmisartan 40 and 80 mg. There is an apparent trend to a dose relationship to a time to recovery of baseline systolic blood pressure (SBP). In this respect, data concerning diastolic blood pressure (DBP) are inconsistent. In patients with hypertension, telmisartan reduces both systolic and diastolic blood pressure without affecting pulse rate. The antihypertensive efficacy of telmisartan has been compared to antihypertensive drugs such as amlodipine, atenolol, enalapril, hydrochlorothiazide, losartan, lisinopril, ramipril and valsartan.

Upon abrupt cessation of treatment with telmisartan, blood pressure gradually returns to pre-treatment values over a period of several days without evidence of rebound hypertension. Telmisartan treatment has been shown statistically significant reductions in Left Ventricular Mass and Left Ventricular Mass Index in patients with hypertension and Left Ventricular Hypertrophy.

Telmisartan treatment has been shown (including comparators like losartan, Ramipril and valsartan) to be associated with statistically significant reductions in proteinuria (including microalbuminuria and macroalbuminuria) in patients with hypertension and diabetic nephropathy. The incidence of dry cough was significantly lower in patients treated with telmisartan than in those given with angiotensin converting enzyme inhibitors.

Cardiovascular Risk Reduction: The primary endpoint was a composite of cardiovascular death, non-fatal myocardial infarction, nonfatal stroke, or hospitalization for congestive heart failure.

Adherence to treatment was better for telmisartan than for ramipril or the combination of telmisartan and ramipril, although the study population had been pre-screened for tolerance to treatment with an ACE-inhibitor. The analysis of adverse events leading to permanent treatment discontinuation and of serious adverse events showed that cough and angioedema were less frequently reported in patients treated with telmisartan than in patients treated with ramipril, whereas hypotension was more frequently reported with telmisartan.

Telmisartan had similar efficacy to ramipril in reducing the primary endpoint. The incidence of the primary endpoint was similar in the telmisartan (16.7%), ramipril (16.5%) and telmisartan plus Ramipril combination (16.3%) arms. The hazard ratio for telmisartan vs. ramipril was 1.01 [97.5% CI 0.93-1.10, p (non-inferiority) =0.0019]. The treatment effect was found to persist following corrections for differences in systolic blood pressure at baseline and over time. There was no difference in the primary endpoint based on age, gender, race, baseline therapies or underlying disease.

Combining telmisartan with ramipril did not add further benefit over ramipril or telmisartan alone. In addition, there was a significantly higher incidence of hyperkalaemia, renal failure, hypotension and syncope in the combination arm. Therefore, the use of a combination of telmisartan and ramipril is not recommended in this population.

Pharmacokinetics

Absorption of telmisartan is rapid although the amount absorbed varies. The mean absolute bioavailability for telmisartan is about 50%.

When telmisartan is taken with food, the reduction in the area under the plasma concentration-time curve (AUC) of telmisartan varies from approximately 6% (40 mg dose) to approximately 19% (160 mg dose). By 3 hours after administration, plasma concentrations are similar whether telmisartan is taken fasting or with food.

The small reduction in AUC is not expected to cause a reduction in the therapeutic efficacy.

Gender differences in plasma concentrations were observed, C_{max} and AUC being approximately 3- and 2-fold higher, respectively, in females compared to males without relevant influence on efficacy. Telmisartan is largely bound to plasma protein (>99.5%), mainly albumin and alpha-1 acid glycoprotein. The mean steady-state apparent volume of distribution (V_{ss}) is approximately 500 L. Telmisartan is metabolised by conjugation to the glucuronide of the parent compound. No pharmacological activity has been shown for the conjugate.

Telmisartan is characterised by biexponential decay pharmacokinetics with a terminal elimination half-life of >20 hours. The maximum plasma concentration (C_{max}) and, to a smaller extent, area under the plasma concentration-time curve (AUC) increase disproportionately with dose. There is no evidence of clinically relevant accumulation of telmisartan.

After oral (and intravenous) administration, telmisartan is nearly exclusively excreted with the faeces, exclusively as unchanged compound. Cumulative urinary excretion is <2% of dose. Total plasma clearance (Cl_{Tot}) is high (approximately 900 mL/min) compared with hepatic blood flow (about 1,500 mL/min).

Elderly Patients: The pharmacokinetics of telmisartan do not differ between younger and elderly patients.

Patients with Renal Impairment: Lower plasma concentrations were observed in patients with renal insufficiency undergoing dialysis. Telmisartan is highly bound to plasma protein in renal-insufficient subjects and cannot be removed by dialysis. The elimination half-life is not changed in patients with renal impairment.

Patients with Hepatic Impairment: Pharmacokinetic studies in patients with hepatic impairment showed an increase in absolute bioavailability up to nearly 100%. The elimination half-life is not changed in patients with hepatic impairment.

Indications

Hypertension: Treatment of essential hypertension.

Cardiovascular Risk Reduction: Telmisartan is indicated for reduction of the risk of myocardial infarction, stroke, or death from cardiovascular causes in patients 55 years of age or older at high risk of developing major cardiovascular events who are unable to take ACE inhibitors.

High risk of cardiovascular events can be evidenced by history of coronary artery disease, peripheral arterial disease, stroke, transient ischemic attack, or high-risk diabetes (insulin-dependent or non-insulin dependent) with evidence of end-organ damage. Telmisartan can be used in addition to other needed treatment (such as antihypertensive, antiplatelet or lipid-lowering therapy).

Studies of telmisartan in this setting do not exclude that it may not preserve a meaningful fraction of the effect of the ACE inhibitor to which it was compared. Consider using the ACE inhibitor first, and, if it is stopped for cough only, consider re-trying the ACE inhibitor after the cough resolves.

Use of telmisartan with an ACE inhibitor is not recommended.

Dosage and Method of Administration

Adults:

Essential Hypertension: The recommended dose is 40 mg once daily. In cases where the target blood pressure is not achieved, telmisartan dose can be increased to a maximum of 80 mg once daily. Alternatively, telmisartan may be used in combination with thiazide-type diuretics such as hydrochlorothiazide, which has been shown to have an additive blood pressure lowering effect with telmisartan. When considering raising the dose, it must be borne in mind that the maximum antihypertensive effect is generally attained four-eight weeks after the start of treatment. In patients with severe hypertension treatment with telmisartan at doses up to 160 mg alone and in combination with hydrochlorothiazide 12.5-25 mg daily was well tolerated and effective.

Cardiovascular Risk Reduction: The recommended dose of telmisartan tablets is 80 mg once a day and can be administered with or without food. It is not known whether doses lower than 80 mg of telmisartan are effective in reducing the risk of cardiovascular morbidity and mortality.

When initiating telmisartan therapy for cardiovascular risk reduction, monitoring of blood pressure is recommended, and if appropriate adjustment of medications that lower blood pressure may be necessary.

Renal Impairment: No posology adjustment is required for patients with renal impairment including those on haemodialysis. Telmisartan is not removed from the blood by haemofiltration.

Hepatic Impairment: In patients with mild to moderate hepatic impairment, the posology should not exceed 40 mg once daily.

Elderly: No dosing adjustment is necessary.

Children and Adolescents: The safety and efficacy of telmisartan for use in children below 18 years have not been established

Administration:

Telmisartan may be taken with or without food.

Contraindications

Hypersensitivity to telmisartan or any of the excipients.

Biliary obstructive disorders.

Severe hepatic impairment.

The concomitant use of telmisartan with aliskiren is contraindicated in patients with diabetes mellitus or renal impairment (GFR <60 ml/min/1.73 m²).

In cases of rare hereditary conditions that may be incompatible with an excipient of the product (see *Warnings and Precautions*), the use of the product is contraindicated.

Second and third trimester of pregnancy (see *Fertility, pregnancy and lactation*)

Lactation. (See *Fertility, pregnancy and lactation*)

Warnings and Precautions

Pregnancy: Angiotensin II receptor antagonists should not be initiated during pregnancy.

Unless continued angiotensin II receptor antagonist therapy is considered essential, patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy.

When pregnancy is diagnosed, treatment with angiotensin II receptor antagonists should be stopped immediately, and if appropriate, alternative therapy should be started.

Renovascular Hypertension: There is an increased risk of severe hypotension and renal insufficiency when patients with bilateral renal artery stenosis or stenosis of the artery to a single functioning kidney are treated with medicinal products that affect the renin-angiotensin-aldosterone system.

Renal Impairment and Kidney Transplant: When telmisartan is used in patients with impaired renal function, a periodic monitoring of potassium and creatinine serum levels is recommended. There is no experience regarding the administration of telmisartan in patients with a recent kidney transplant.

Intravascular Volume Depletion: Symptomatic hypotension, especially after the first dose, may occur in patients who are volume and/or sodium depleted by vigorous diuretic therapy, dietary salt restriction, diarrhoea or vomiting. Such conditions, especially volume and/or sodium depletion, should be corrected before the administration of telmisartan.

Dual Blockade of the Renin-Angiotensin-Aldosterone System: As a consequence of inhibiting the renin-angiotensin-aldosterone system changes in renal function (including acute renal failure) have been reported in susceptible individuals, especially if combining medicinal products that affect this system. Dual blockade of the renin-angiotensin-aldosterone system (e.g. by adding an ACE-inhibitor or the direct renin-inhibitor aliskiren to an angiotensin II receptor antagonist) should therefore be limited to individually defined cases with close monitoring of renal function (see *Contraindication*).

Other Conditions with Stimulation of the Renin-Angiotensin-Aldosterone System: In patients whose vascular tone and renal function depend predominantly on the activity of the renin-angiotensin-aldosterone system (e.g. patients with severe congestive heart failure or underlying renal disease, including renal artery stenosis), treatment with medicinal products that affect this system has been associated with acute hypotension, hyperazotaemia, oliguria, or rarely acute renal failure.

Primary Aldosteronism: Patients with primary aldosteronism generally will not respond to antihypertensive medicinal products acting through inhibition of the renin-angiotensin system. Therefore, the use of telmisartan is not recommended.

Aortic and Mitral Valve Stenosis, Obstructive Hypertrophic Cardiomyopathy: As with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy.

Hyperkalaemia: During treatment with medicinal products that affect the renin-angiotensin-aldosterone system hyperkalaemia may occur, especially in the presence of renal impairment and/or heart failure. Monitoring of serum potassium in patients at risk is recommended. Based on experience with the use of medicinal products that affect the renin-angiotensin system, concomitant use with potassium-sparing diuretics, potassium supplements, salt substitutes containing potassium or other medicinal products that may increase the potassium level (heparin, etc.) may lead to an increase in serum potassium and should therefore be co-administered cautiously with telmisartan.

Hepatic Impairment: Telmisartan is mostly eliminated in the bile. Patients with biliary obstructive disorders or hepatic insufficiency can be expected to have reduced clearance. telmisartan should be used with caution in these patients.

Diabetes Mellitus: In diabetic patients with an additional cardiovascular risk, i.e. patients with diabetes mellitus and coexistent coronary artery disease (CAD), the risk of fatal myocardial infarction and unexpected cardiovascular death may be increased when treated with blood pressure lowering agents such as ARBs or ACE-inhibitors. In patients with

diabetes mellitus CAD may be asymptomatic and therefore undiagnosed. Patients with diabetes mellitus should undergo appropriate diagnostic evaluation, e.g. exercise stress testing, to detect and to treat CAD accordingly before initiating treatment with telmisartan.

Other: As observed for angiotensin converting enzyme inhibitors, angiotensin receptor blockers including telmisartan are apparently less effective in lowering blood pressure in black people than in non-blacks, possibly because of higher prevalence of low-renin states in the black hypertensive population.

As with any antihypertensive agent, excessive reduction of blood pressure in patients with ischaemic cardiopathy or ischaemic cardiovascular disease could result in a myocardial infarction or stroke.

Effects on Ability to Drive and Use Machines: No studies on the effect on the ability to drive and use machines have been performed. However, when driving vehicles or operating machinery it should be taken into account that dizziness or drowsiness may occasionally occur when taking antihypertensive therapy.

Drug Interactions

Telmisartan may increase the hypotensive effect of other antihypertensive agents. Other interactions of clinical significance have not been identified.

Co-administration of telmisartan did not result in a clinically significant interaction with digoxin, warfarin, hydrochlorothiazide, glibenclamide, ibuprofen, paracetamol, simvastatin and amlodipine. For digoxin, a 20% increase in median plasma digoxin trough concentration has been observed (39% in a single case), monitoring of plasma digoxin levels should be considered.

The co-administration of telmisartan and ramipril led to an increase of up to 2.5 fold in the AUC_{0-24} and C_{max} of ramipril and ramiprilat. The clinical relevance of this observation is not known.

Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin converting enzyme inhibitors. Cases have also been reported with angiotensin-II receptor antagonists including telmisartan. Therefore, serum lithium level monitoring is advisable during concomitant use.

Treatment with NSAIDs (i.e. ASA at anti-inflammatory dosage regimens, COX-2 inhibitors and non-selective NSAIDs) is associated with the potential for acute renal insufficiency in patients who are dehydrated. Compounds acting on the Renin-Angiotensin-System like telmisartan may have synergistic effects. Patients receiving NSAIDs and telmisartan should be adequately hydrated and be monitored for renal function at the beginning of combined treatment.

A reduced effect of antihypertensive drugs like telmisartan by inhibition of vasodilating prostaglandins has been reported during combined treatment with NSAIDs.

Fertility, pregnancy and lactation

The use of angiotensin II receptor antagonists is not recommended during the first trimester of pregnancy and should not be initiated during pregnancy. When pregnancy is diagnosed, treatment with angiotensin II receptor antagonists should be stopped immediately, and, if appropriate, alternative therapy should be started.

The use of angiotensin II receptor antagonists is contraindicated during the second and third trimester of pregnancy.

Non-clinical studies with telmisartan do not indicate teratogenic effect, but have shown fetotoxicity. Angiotensin II receptor antagonists exposure during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalaemia).

Unless continued angiotensin II receptor antagonist therapy is considered essential, patients planning pregnancy should be changed to alternative antihypertensive treatments which have an established safety profile for use in pregnancy.

Should exposure to angiotensin II receptor antagonists have occurred from the second trimester of pregnancy, ultrasound check of renal function and skull is recommended.

Infants whose mothers have taken angiotensin II receptor antagonists should be closely observed for hypotension.

Telmisartan is contraindicated during lactation since it is not known whether it is excreted in human milk. Animal studies have shown excretion of telmisartan in breast milk.

Impairment of Fertility: No studies on fertility in humans have been performed. In non-clinical studies, an effect of telmisartan on male and female fertility was not observed.

Undesirable Effects

The overall incidence of adverse events reported with telmisartan (41.4%) was usually comparable to placebo (43.9%) in patients treated for hypertension. The incidence of adverse events was not dose related and showed no correlation with gender, age or race of the patients.

The safety profile of telmisartan in patients treated for cardiovascular risk reduction was consistent with that obtained in hypertensive patients.

Infections and Infestations: Urinary tract infections (including cystitis), upper respiratory tract infections, sepsis including fatal outcome.

Blood and Lymphatic System Disorders: Anaemia, eosinophilia, thrombocytopenia.

Immune System Disorders: Anaphylactic reaction, hypersensitivity.

Metabolism and Nutrition Disorders: Hyperkalaemia, hypoglycaemia (in diabetic patients).

Psychiatric Disorders: Insomnia, depression, anxiety.

Nervous System Disorders: Syncope (faint).

Eye Disorders: Visual disturbance.

Ear and Labyrinth Disorders: Vertigo.

Cardiac Disorders: Bradycardia, tachycardia.

Vascular Disorders: Hypotension, orthostatic hypotension.

Respiratory, Thoracic and Mediastinal Disorders: Dyspnoea.

Gastrointestinal Disorders: Abdominal pain, diarrhoea, dyspepsia, flatulence, vomiting, dry mouth, stomach discomfort.

Hepatobiliary Disorders: Abnormal hepatic function or liver disorder*.

*Most cases of hepatic function abnormal/liver disorder from post-marketing experience with telmisartan occurred in patients in Japan, who are more likely to experience these adverse reactions.

Skin and Subcutaneous Tissue Disorders: Pruritus, hyperhidrosis, rash, angioedema (with fatal outcome), eczema, erythema, urticaria, drug eruption, toxic skin eruption.

Musculoskeletal, Connective Tissue and Bone Disorders: Back pain, muscle spasms (cramps in legs), myalgia, arthralgia, pain in extremity (leg pain), tendon pain (tendinitis-like symptoms).

Renal and Urinary Tract Disorders: Renal impairment including acute renal failure (see *Warning and Precautions*).

General Disorders and Administration Site Conditions: Chest pain, asthenia (weakness), influenza-like illness.

Investigations: Increased blood creatinine, decreased haemoglobin, increased blood uric acid, increased hepatic enzymes, increased blood creatinine phosphokinase (CPK).

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the local reporting system.

Overdose

Limited information is available with regard to overdose in humans. The most prominent manifestations of telmisartan overdose were hypotension and tachycardia, bradycardia also occurred. If symptomatic hypotension should occur, supportive treatment should be instituted. Telmisartan is not removed by haemodialysis.

Incompatibility

Not applicable.

Shelf-Life

24 months

Storage and Handling Instruction

Store below 30°C.

Telmisartan should be kept in the sealed blister due to the hygroscopic property of the tablets. Tablets should be taken out of the blister shortly before administration.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Packaging Information

A carton containing blister pack of 3 X 10 tablets

Manufactured by

Jubilant Generics Limited,
Village Sikandarpur Bhainswal, Roorkee- Dehradun Highway,
Bhagwanpur, Roorkee,
Distt-Haridwar,
Uttarakhand- 247 661 India.

Product Registration Holder

CIPLA MALAYSIA SDN BHD
Suite 1101, Amcorp Tower, Amcorp Trade Centre,
18 Persiaran Barat, 46050
Petaling Jaya, Selangor, Malaysia

Last Updated: *October 2019*