

SAME SIZE ARTWORK
LEAFLET SIZE: 130 mm x 100 mm

33 mm

25 mm

hairy sites, the hair should be parted before application to allow direct contact with the lesions. The frequency of application is generally 2 to 3 times daily. After controlling the eczematous symptoms, a switch over to CANDID Cream or Lotion may be advised till the primary fungal infection is completely eradicated.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE AND ANTIDOTE (S):
Not applicable since this formulation is for topical use and are normally well tolerated. However, if irritation develops, the preparation should be discontinued.

PACKING / PACK SIZES : Bottle of 15ml.

SHELF LIFE : 2 Years

STORAGE CONDITIONS, USER INSTRUCTIONS AND PHARMACEUTICAL PRECAUTIONS
Store below 30°C. Protect from light.
MAL 20033330AZ
Date of Revision: 31 March 2021

Product Registration Holder:
Glenmark Pharmaceuticals (Malaysia)
Sdn Bhd (660397-W) D-31-02,
Menara Suezcap 1, No. 2, Jalan Kerinchi,
59200 Kuala Lumpur, Malaysia

Manufactured by:
glenmark
PHARMACEUTICALS LTD.
Plot No. E-37, 39, D-Road,
MIDC, Satpur, Nashik - 422 007,
Maharashtra State, India.
© Trade Mark

MY
PE00000

For the use only of Registered Medical Practitioner or
a Hospital or a Laboratory

Clotrimazole+
Anhydrous Beclometasone
candid-B®
Lotion

SPACE FOR
PHARMACODE

COMPOSITION :
Clotrimazole USP 1%
Anhydrous Beclometasone Dipropionate BP 0.025%
Base q.s.

THERAPEUTIC CLASS : T0130

DESCRIPTION: A printed carton containing a leaflet and a labelled LDPE bottle fitted with plug and green coloured cap containing a clear, colourless to pale yellow, viscous liquid.

ACTIONS AND MODE OR MECHANISM OF ACTION :
CANDID-B Lotion is a broad spectrum Antimycotic, Anti-inflammatory and Antipruritic. It offers effective therapy for dermatomycoses complicated by inflammation, pruritus, erythema or other eczematous features.
The active ingredients of CANDID-B Lotion are Clotrimazole and Beclometasone Dipropionate.
Clotrimazole has broad spectrum fungicidal activity against Dermatophytes (Trichophyton, Epidermophyton and Microsporum spp.), Candida and Aspergillus spp., Malassezia furfur and other fungi. It also inhibits some strains of Gram-positive and Gram-negative bacteria.
Beclometasone Dipropionate, the potent and fast acting topical corticosteroids exerts marked anti-inflammatory, antipruritic, anti-allergic and vaso-constrictive effects. On account of these properties it gives prompt relief in inflammation, pruritis, erythema etc.

130 mm

100 mm

SAME SIZE ARTWORK
LEAFLET SIZE: 130 mm x 100 mm

PHARMACOLOGY (SUMMARY OF PHARMACODYNAMICS AND PHARMACOKINETICS):
PHARMACODYNAMICS:
CANDID-B Lotion is a combination of Clotrimazole and Beclometasone Dipropionate. The pharmacological aspects of both the ingredients are given below:
Clotrimazole
Clotrimazole is an imidazole antifungal agent and is effective against a wide variety of fungi including Epidermophyton, Microsporum and Trichophyton spp., Coccidioides immitis, Histoplasma capsulatum, Aspergillus and Candida spp. Other fungi responding to Clotrimazole include Cryptococcus neoformans, Malassezia furfur (Pityrosporum orbiculare), Paracoccidioides brasiliensis, Sporothrix schenckii, Pseudallescheria, Medurella and Phialophora spp.
Clotrimazole also has antibacterial action and some strains of Staphylococcus aureus and Streptococcus pyogenes have been reported to be sensitive.
Clotrimazole is believed to exert its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisation and the incorporation of the acylated amino acids essential for fungi.
Beclometasone Dipropionate
Beclometasone Dipropionate is a synthetic glucocorticoid and is used by topical application for the treatment of various dermatoses.
The topical application of Beclometasone Dipropionate produces dramatic suppression of dermatoses by potent and rapid anti-inflammatory, anti-pruritic, anti-allergic, anti-proliferative and vaso-constrictive actions.
Unlike fluorinated topical steroids, Beclometasone Dipropionate is less likely to produce adrenal suppression or changes in the skin as Beclometasone Dipropionate is metabolised in the skin.
PHARMACOKINETICS :
Clotrimazole appears to be well absorbed in humans following oral administration and is eliminated mainly as inactive metabolites. Following topical and vaginal administration, however, clotrimazole appears to be minimally absorbed. Six hours after the application of radioactive clotrimazole 1% cream and 1% solution into intact and acutely inflamed skin, the concentration of clotrimazole varied from 100 mcg/cm² in the stratum corneum to 0.5 to 1 mcg/cm² in the stratum reticulare, and 0.1 mcg/cm² in the subcutis. No measurable amount of radioactivity (± 0.001 mcg/ml) was found in the serum within 48 hours after application under occlusive dressing of 0.5 ml of the solution or 0.8 g of the cream. Only 0.5% or less of the applied radioactivity was excreted in the urine.
The extent of percutaneous absorption of topical corticosteroids is determined by many factors including the vehicle, the integrity of the epidermal barrier, and the use of occlusive dressings. Topical corticosteroids can be absorbed from normal intact skin.
Inflammation and/or other disease processes in the skin increases

percutaneous absorption. Occlusive dressings substantially increase the percutaneous absorption of topical corticosteroids.
Once absorbed through the skin, topical corticosteroids are handled through pharmacokinetic pathways similar to systemically administered corticosteroids. Corticosteroids are bound to plasma proteins in varying degrees. Corticosteroids are metabolised primarily in the liver and are then excreted by the kidneys. Some of the topical corticosteroids and their metabolites are also excreted into the bile.

INDICATIONS :
CANDID-B Lotion can be of benefit in the fungal infections of the skin, particularly when accompanied by eczematous features :
● Tinea Capitis ● Tinea Pedis
● Tinea Corporis ● Tinea Versicolor
● Tinea Barbae ● Erythrasma
● Tinea Cruris ● Candidal Vulvitis
● Tinea Manuum ● Candidal Nappy Rash
● Tinea Unguium ● Candidal Paronychia
CANDID-B Lotion will be particularly useful in large lesions and for hairy and intertriginous skin areas.

CONTRAINDICATIONS :
These formulations should not be used in tuberculosis of the skin, herpes simplex, chicken pox, measles, vaccinia, syphilitic skin lesions and in cases of hypersensitivity to any of the components.

SIDE EFFECTS / ADVERSE REACTIONS :
The use of steroids topically over large skin areas, with or without occlusive dressing, may result in systemic absorption of the steroid and suitable precautions should be taken by physician. When occlusive dressings are put, miliaria, folliculitis or pyoderma may sometimes develop beneath the occlusive material. Localised atrophy and striae have been reported with topical steroids.

DRUG INTERACTIONS :
None reported.

PRECAUTIONS / WARNINGS:
CANDID-B Lotion is normally well tolerated. If irritation develops, the preparation should be discontinued.
This formulation should not be used in the eyes or in the ear if the eardrum is perforated.
Though Topical steroids have not been reported to cause adverse effect on pregnancy, their safety in pregnancy has not been firmly established. They should, therefore, not be used in large amounts or for long periods of time during pregnancy.

RECOMMENDED DOSAGE, DOSAGE SCHEDULE AND ROUTE OF ADMINISTRATION :
The affected area is washed and dried. CANDID-B Lotion is then rubbed in. In

PE00000 MY

130 mm

100 mm

ICONGRAPHICS CODE:

PANTONE SHADE

PANTONE
BLACK
PROCESS C

Supersedes
Artwork Code

PHARMACODE: