

Insertion can be carried out more easily when lying on the back with the knees bent and spread out.

6. The Applicator after each use should be taken apart for cleaning by holding the cylinder with one hand and pulling out the plunger with slight force. Wash with soap and lukewarm water (not boiling water). Rinse thoroughly. To reassemble the Applicator, insert the plunger into the cylinder.

SYMPTOMS AND TREATMENT FOR OVERDOSAGE AND ANTIDOTE (S):

Acute overdose with vaginal application of Clotrimazole is unlikely and it is not expected to lead to a life threatening situation. In the event of accidental oral ingestion, routine measures such as gastric lavage should be performed as soon as possible after ingestion.

PACKING / PACK SIZES : Tube of 30g with applicator.

STORAGE CONDITIONS, USER INSTRUCTIONS AND PHARMACEUTICAL PRECAUTIONS:

Store below 30°C. Protect from freezing & light.

MAL2003331AZ

19 October 2023

Product Registration Holder:

Glenmark Pharmaceuticals
(Malaysia) Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1,
No. 2, Jalan Kerinchi,
59200 Kuala Lumpur,
Malaysia

Manufactured by:

glenmark
PHARMACEUTICALS LTD.
Plot No. E-37, 39, D-Road, MIDC,
Satpur, Nashik - 422 007,
Maharashtra State, India.
™ Trade Mark

PE000000 MY

*For the use only of Registered Medical Practitioner
or a Hospital or a Laboratory*

candid[®]-v GEL

SPACE FOR
PHARMACODE

COMPOSITION:

Clotrimazole USP 2%

Preservatives:

Benzyl Alcohol USP-NF 2%

Chlorocresol USP-NF 0.1%

THERAPEUTIC CLASS: T0130

DESCRIPTION: A printed carton with an insert, applicator and a printed aluminium collapsible tube containing white, smooth, semi-solid gel.

ACTIONS AND MODE OR MECHANISM OF ACTION :

The active substance of CANDID is Clotrimazole which is bis-phenyl-(2-chlorophenyl)-1-dazolylmethane.

CANDID has a wide spectrum of activity against Dermatophytes (Trichophyton, Microsporum, Epidermophyton), Yeasts (Candida sp., Torulopsis sp.), Mould fungi and other fungi (Aspergillus sp., Malassezia furfur, Nocardia sp., etc.). It also has action against Trichomonas and some strains of Gram - positive bacteria (Streptococci, Staphylococci and Corynebacterium minitissimum) and Gram - negative bacteria (Bacteroides and Haemophilus vaginalis). CANDID acts on the cell membrane of the fungus. It appears that it inhibits protein synthesis and influences the utilisation and incorporation of the acylated amino acids essential to fungi.

PHARMACOLOGY (SUMMARY OF PHARMACODYNAMICS AND PHARMACOKINETICS):

PHARMACODYNAMICS:

Clotrimazole is a broad-spectrum antifungal agent that is used for treatment of dermal infections caused by various species of pathogenic dermatophytes, yeasts and Malassezia furfur. The primary action of Clotrimazole is against dividing any growing organisms.

In vitro Clotrimazole exhibits fungistatic and fungicidal activity against isolates Trichophyton rubrum, Trichophyton mentagrophytes, Epidermophyton floccosum, Microsporum canis and Candida species including Candida albicans. In general, the in vitro activity of Clotrimazole corresponds to that of tolnaftate and griseofulvin against the mycelia of dermatophytes

ICONGRAPHICS CODE:

PANTONE SHADE



PANTONE
BLACK
PROCESS C

**Supersedes
Artwork Code:
PE55269**

PHARMACODE:

(Trichophyton, Microsporum and Epidermophyton), and to that of polyenes (amphotericin B and nystatin), against budding fungi (Candida). Using as in vitro Clotrimazole was effective in preventing the growth of pseudomycelia and mycelia of Candida albicans. Strains of fungi having a natural resistance to Clotrimazole are rare. Only single isolate of Candida guilliermondii has been reported to have primary resistance to Clotrimazole. No single step or multiple-step resistance to Clotrimazole had developed during successive passages of Candida albicans and Trichophyton metagrophytes.

No appreciable change in sensitivity was detected after successive passages of isolates of Candida albicans, Candida Krusei or Candida Pseudotropicalis in liquid or solid media containing Clotrimazole.

Also, resistance could not be developed in chemically induced mutant strains of polyene-resistant isolates of Candida albicans.

In studies of the mechanism of action, the minimum fungicidal concentration of Clotrimazole caused leakage of intracellular phosphorus compounds into the ambient medium with concomitant breakdown of cellular nucleic acids and accelerated potassium efflux. Both these events began rapidly and extensively for addition of the drug.

PHARMACOKINETICS :

Clotrimazole appears to be well absorbed in humans following oral administration and is eliminated mainly as inactive metabolites. Following topical and vaginal administration, however, clotrimazole appears to be minimally absorbed. Six hours after the application of radioactive clotrimazole 1% cream and 1% solution into intact and acutely inflamed skin, the concentration of clotrimazole varied from 100 mcg/cm³ in the stratum corneum to 0.5 to 1 mcg/cm³ in the stratum reticulare, and 0.1 mcg/cm³ in the subcutis. No measurable amount of radioactivity (< 0.001 mcg/ml) was found in the serum within 48 hours after application under occlusive dressing of 0.5 ml of the solution or 0.8 g of the cream. Only 0.5% or less of the applied radioactivity was excreted in the urine.

INDICATIONS :

For the treatment of vulvovaginal candidiasis.

CONTRAINDICATIONS:

Except for rare cases of hypersensitivity no other cases of contraindications have been reported with the use of **Candid V gel**. **Candid V gel** is contraindicated in individuals who have shown hypersensitivity to any of the ingredients of the preparation.

SIDE EFFECTS/ADVERSE REACTIONS :

This is virtually no absorption of the active ingredients through the vaginal mucosa. No systemic effects are therefore to be expected. Candid V gel is well tolerated

locally. Local mild burning or irritation, immediately after application Hypersensitivity reactions may occur.

DRUG INTERACTIONS :

None reported.

PRECAUTIONS/WARNINGS:

If irritation or sensitivity develops with the use of Clotrimazole, treatment should be discontinued and appropriate therapy instituted. Animal studies have not shown any evidence of carcinogenicity or mutagenicity.

Pregnancy

There are no adequate and well-controlled studies in pregnant women during first trimester although in clinical trials, use of vaginally applied Clotrimazole in pregnant women in their second and third trimester has not been associated with ill effects. **Candid V gel** should be used during pregnancy only if clearly needed.

Lactation

It is not known whether this drug is excreted in human milk because many drugs are excreted in human milk, caution should be exercised when Candid V gel is used by nursing women.

Candid V gel contains propylene glycol as one of the inactive ingredients, which may cause hypersensitivity to any of the ingredients of the preparation.

Candid V gel may reduce the efficacy and safety of latex contraceptive such as condom & diaphragm.


RECOMMENDED DOSAGE, DOSAGE SCHEDULE AND ROUTE OF ADMINISTRATION :

To be applied intravaginally, once a day at bedtime for 3 days using the application provided.


DIRECTIONS FOR USE OF THE APPLICATOR :

1. Remove the cap from the tube and attach the applicator in its place.

2. Squeeze the tube from the bottom to fill the Gel into the cylinder and push out the plunger as far out till the cylinder is filled.



3. Always roll the tube from the bottom to allow easy filling of the Applicator and complete utilisation of the contents.



4. Remove the filled Applicator from the tube and close the tube with the cap.

5. Hold the filled Applicator by the cylinder and gently insert the Applicator into the vagina as deeply as possible. Press the plunger completely to empty the contents of the Applicator. With the plunger still depressed, remove the Applicator from the vagina, hold it by the cylinder.



PE00000 MY

ICONGRAPHICS CODE:

PANTONE SHADE



PANTONE
BLACK
PROCESS C

**Supersedes
Artwork Code:
PE55269**

PHARMACODE: