

PACKAGE INSERT

SUNADINE TABLET 10MG/ SUNADINE SYRUP 5MG/5ML

Tablet : Loratadine 10mg / tablet
Syrup : Loratadine 5mg / 5ml

Product Description:

Tablet : Oval, white, convex, film-coated tablet with single score on one side only.
Syrup : A clear, colourless syrup with peach flavour.

Pharmacodynamics:

Antihistamines (loratadine) diminish or abolish the main actions of histamine in the body by competitive, reversible blockade of histamine receptor sites on tissues. Histamines H₁-receptors are responsible for vasodilatation, increased capillary permeability, flare and itch reactions in the skin, and to some extent for contraction of smooth muscle in the bronchi and gastrointestinal tract.

Loratadine is a long-acting tricyclic antihistamine with selective peripheral histamine H₁-receptor antagonistic activity. Studies have been shown that neither loratadine nor its metabolites readily cross the blood-brain barrier. It has higher affinity for the peripheral rather than the central nervous system H₁-receptors. Hence, less problem of daytime sedation for those who have to drive or operate machinery.

Pharmacokinetics:

It is rapidly absorbed from the gastrointestinal tract after oral administration. The peak plasma concentration is being attained at about one hour. Loratadine undergoes extensive metabolism. Descarboethoxyloratadine, which is the major metabolite has potent histamine H₁-receptor blocking activity. Reported half-lives for loratadine and descarboethoxyloratadine are about 12 and 18 hours respectively.

Loratadine is about 98% bound to plasma proteins while its major metabolite, descarboethoxyloratadine is less extensively bound. Loratadine and its metabolite have been detected in breast milk, but do not appear to cross the blood-brain barrier to a significant extent. Loratadine and its metabolites are excreted in the urine and faeces.

Indication:

Loratadine is indicated for the relief of nasal as well as non-nasal symptoms of seasonal allergic rhinitis, chronic urticaria and other allergic dermatological disorders.

Recommended Dosage:

Category	Dose
Children 2-6 years of age: Body weight > 30kg Body weight ≤ 30kg	Two 5ml spoonful (10mg) syrup once daily. One 5ml spoonful (5mg) syrup once daily.
Children 6-12 years of age: Body weight > 30kg Body weight ≤ 30kg	One tablet (10mg)/Two 5ml spoonful (10mg) syrup once daily. Half a tablet (5mg)/One 5ml spoonful (5mg) syrup once daily.
Adults and children 12 years of age and over:	One tablet (10mg)/Two 5ml spoonful (10mg) syrup once daily.

The use of loratadine in children under the age of 2 have not yet been established.

Route of Administration:

Oral

Contraindications:

Contraindicated in patients with known hypersensitivity or idiosyncrasy to loratadine or other components in the product.

Warnings and Precautions:

Dose should be lowered (5mg daily or 10mg every other day) in patients with liver impairment or renal impairment (GFR < 30mL/min). It should also be used with caution in patients with prostatic hypertrophy, urinary retention, glaucoma and epilepsy. Children and elderly are more susceptible to side effects.

Safety and efficacy of loratadine in children younger than 2 years of age have not yet been established. Drug known to inhibit hepatic metabolism should be coadministered with caution until definitive interaction studies can be completed.

Interactions with Other Medicaments:

Loratadine has no potentiating effect when administered concurrently with alcohol. Increase in plasma concentrations of loratadine has been reported with concomitant use of ketoconazole, erythromycin or cimetidine in healthy individuals but without clinically significant changes. Caution should be exercised when coadministering drugs known to inhibit hepatic metabolism. Drug/lab test: Loratadine should be discontinued approximately 48 hours prior to skin testing procedures. Since antihistamines may prevent or diminish otherwise positive reactions to dermal reactivity indicators.

Pregnancy and Lactation:

There are no well-controlled studies in pregnant women. Loratadine and its metabolite, descarboethoxyloratadine have been detected in breast milk. Hence, loratadine should only be used when the potential benefit outweighs the potential risk.

Side Effects:

Gastritis, alopecia, anaphylaxis, abnormal hepatic function, tachycardia, palpitation, fatigue, headache, somnolence, dry mouth, nervousness, wheezing, abdominal pain, hyperkinesia, malaise, nausea and rash has been reported.

Symptoms and Treatment of Overdose:

In adults, somnolence, tachycardia and headache have been reported with doses greater than 10mg (40-180mg). In children, extrapyramidal side effects and palpitations have been reported in doses greater than 10mg.

General symptomatic and supportive measures should be instituted in the event of overdosage. The treatment should consist of emesis, preferably by administration of ipecac syrup except in patients with impaired consciousness. Following emesis, activated charcoal should be administered to absorb any remaining drug. However, if emesis is unsuccessful or contraindicated, gastric lavage should be performed using normal saline. Saline cathartics may be of value as it draw water into the bowel by osmosis, hence rapidly diluting the bowel content. Hemodialysis is of no value as loratadine is not eliminated by hemodialysis.

Effects on Ability to Drive and Use Machine:

In clinical trials that assessed driving ability, no impairment occurred in patients receiving loratadine. However, patients should be informed that very rarely some people experience drowsiness, which may affect their ability to drive or use machines.

Storage Conditions: Store at or below 30°C. Protect from light.

Pack Size: Tablet: Bottle pack : A bottle of 30 tablets.
Blister pack: A box of 3x10s, 10x10s, 50x10s and 100x10s per strip.
Syrup: Bottle pack : A bottle of 60ml, 100ml and 120ml.

Pack Size (export only): Tablet: A bottle of 100 tablets and 500 tablets.
Syrup: A bottle of 1 litre.

Shelf-life: Tablet - 3 years.
Syrup - 1.5 years.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer & Product Registration Holder:
Sunward Pharmaceutical Sdn. Bhd.
No. 9, 11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

CL673A-R10
Revision Date: 21/10/2022