

DHACTULOSE SYRUP

Product Description:

Clear, viscous liquid, colourless to pale brownish-yellow.

Composition:

Each 5ml contains Lactulose BP 3.35g.

Pharmacodynamics:

Lactulose is a synthetic disaccharide which is absorbed minimally from the GI tract, reaching the colon unchanged. It is then broken down by colonic bacteria into simple organic acids, mainly lactic acids, which lead to a lowering of pH in the colon and exert a local osmotic effect. Fluids retained in the colon by osmosis increase and soften the faecal bulk and hence, stimulate peristalsis. Constipation is relieved and the physiological rhythm of the colon is restored. It may take up to 48 hours before an effect is obtained.

Pharmacokinetics:

Following oral administration, a small amount (less than 3%) of the lactulose dose is absorbed from the small intestine. The absorbed lactulose is not metabolised and subsequently excreted unchanged in the urine. The unabsorbed lactulose reaches the colon unchanged where it is metabolised by colonic bacteria to form lactic acids and small amounts of acetic and formic acids.

Indications:

Relief of occasional or non-persistent constipation.

Dosage: Oral

Adults: Initially, 15 to 30ml daily as a single dose (preferably during breakfast) or in 2 divided doses. The daily dosage should be adjusted according to the patient's needs.

Children: 6-12 years : 10ml

1-5 years : 5ml

under 1 year : 2.5ml. To be given 2 times a day.

Side Effects:

Abdominal discomfort, flatulence and cramps. Nausea and vomiting have been reported with high doses. Diarrhoea with excessive loss of water and electrolytes (particularly potassium) may result from prolonged use or excessive dosage.

Precautions:

Patients with lactose intolerance or in diabetic patients. Consult your doctor if constipation is not relieved after a few days or reoccur after treatment.

Product Name	: Leaflet Dhactulose 200ml Malaysia
Product Code	: 41312-02
Size	: 110 x 175 mm
Paper	: HVS 70 gr
Finishing	: ---
Colour	: Black (1/1)
Date / Revise #	: 
Filename	: 41312-02 Lf Dhactulose

Use in Pregnancy:

No evidence of teratogenicity has been reported. Lactulose should be used during pregnancy only if clearly needed.

Contraindications:

Patients with intestinal obstruction, galactosemia, on low galactose diet or hypersensitive to any of the ingredients.

Drug Interactions:

Because of the mode of action of lactulose, in which the colonic pH is lowered, the action of substances with release dependent on pH (such as 5-ASA formulations) may be reduced. No other drug interactions have been reported.

Overdosage and Treatment:

No information is available on cases of overdosing. High doses of lactulose can cause abdominal cramps and diarrhoea. Treatment should be ceased and replacement of fluid and electrolyte is given if required.

Packing: 200ml in amber glass

Storage:

Store below 30 °C and away from direct sunlight. Freezing should be avoided.

Expiry Period: Do not use after the expiry date stated on the label.

Date of Revision: November 2021

For further information, please consult your doctor or pharmacist.

Manufactured by:

PT. Actavis Indonesia

Jl. Raya Bogor Km. 28, Kelurahan Pekayon, Kecamatan Pasar Rebo
Kota Jakarta Timur, Provinsi DKI Jakarta, Indonesia

W
Actavis

AAAN8680

41312-02

Product Name	: Leaflet Dhactulose 200ml Malaysia
Product Code	: 41312-02
Size	: 110 x 175 mm
Paper	: HVS 70 gr
Finishing	: ---
Colour	: Black (1/1)
Date / Revise #	: 
Filename	: 41312-02 Lf Dhactulose