



CLOMAZOL SOLUTION

Description: Colourless Solution

Contents:

Clotrimazole 1.00% w/v

Indications:

For topical treatment of tinea pedis, tinea cruris, tinea corporis, candidiasis and tinea versicolor.

Pharmacology:

Clotrimazole has fungistatic and fungicidal (depending on concentration) activity against Dermatophytes (Trichophyton, Epidermophyton and Microsporum sp.), Candida, Aspergillus sp., Malassezia furfur and other fungi. It inhibits biosynthesis of ergosterol or other sterols, damaging the fungal cell wall membrane and altering permeability. As a result, Loss of essential intracellular elements occur. It inhibits biosynthesis of triglycerides and phospholipids by fungi. In addition, it inhibits oxidative and peroxidative enzyme activity, resulting in intracellular buildup of toxic concentration of hydrogen peroxide, which may contribute to deterioration of subcellular organelles and cellular necrosis. In Candida albicans, it inhibits transformation of blastospores to invasive mycelial form. Others action include some antibacterial activity.

When applied topically, clotrimazole penetrates the dermis but there is little, if any systemic absorption.

Contraindications:

It is contraindicated in patients hypersensitive to clotrimazole.

Side effects / Adverse reactions:

When applied topically, local reactions including erythema, urticaria, itching, stinging and irritation may occur in patients.

Precautions:

When this medication is used for treatment of candidiasis, occlusive dressings should not be used since they provide conditions which favour growth of yeast and release of its irritating endotoxin. Discontinue treatment if irritation occurs.

Pregnancy & Lactation:

Risk benefits should be considered when use in pregnant and lactating mothers, although safety of use has not been established.

Drug Interactions:

Nil.

Dosage and Administration:

Apply two to three times daily on affected area. Continue for another two weeks after symptoms disappear.

Treatment of overdosage:

Unlikely to have overdosage.

Storage : Keep container tightly closed. Store in a cool, dry place, below 30°C.

Conditions : Protect from light. Keep away from children.

Presentation : 10ml/bottle.

Shelf-life : 3 years.

Manufactured & : **Prime Pharmaceutical Sdn. Bhd.**

Distributed by : 1505, Lorong Perusahaan Utama 1, Taman Perindustrian Bukit Tengah,
14000 Bukit Mertajam, Penang, Malaysia.

17/01/2022