



Axcil[®]

CETIRIZINE

TABLETS

COMPOSITION:

Each tablet contains
Cetirizine Dihydrochloride 10mg

PRESENTATION:

White, long oblong shaped tablet with score line on both sides.

INDICATIONS:

Adults and children of 3 years or above: symptomatic treatment of seasonal rhinitis, perennial allergic rhinitis and urticaria of allergic origin.

PHARMACOLOGY:

Mechanism of Action:

Cetirizine is an antihistamine as its principal effects are mediated via selective inhibition of peripheral H1 receptors.

Pharmacokinetics:

Cetirizine is rapidly absorbed from the gastro-intestinal tract after oral administration, peak plasma concentrations being attained in about 1 hour. It is highly bound to plasma proteins and has an elimination half-life of about 11 hours. Cetirizine has been detected in breast milk. It is excreted primarily in the urine mainly as unchanged drug.

DOSAGE AND ADMINISTRATION:

Adults and Children 12 Years and above
10mg once daily.

Children 6 to 11 Years
5mg twice daily or 10mg once daily.

Children 3 to 6 Years
2.5mg twice a day or 5mg once daily.

In patients with renal insufficiency, dosage should be reduced to half the daily dose.

Tablets: 1 tablet = 10mg, 1/2 tablet = 5mg

CONTRAINDICATION:

Contraindicated in those patients with known hypersensitivity to any of its ingredients. Cetirizine should not be administered to pregnant women during the first 3 month of pregnancy and women who are breastfeeding.

PRECAUTIONS:

Activities Requiring Mental Alertness:

In clinical trials, the occurrence of somnolence has been reported in some patients taking cetirizine; due caution should therefore be exercised when driving car or operating potentially dangerous machinery. Concurrent use of cetirizine with alcohol or other CNS depressants should be avoided because additional reductions in alertness and additional impairment of CNS performance may occur.

Carcinogenesis, Mutagenesis, and Impairment of Fertility:

In a 2 years carcinogenicity study in rats, cetirizine was not carcinogenic at dietary doses up to 20mg/kg. In a 2 years carcinogenicity study in mice, cetirizine caused an increased incidence of benign liver tumors in males at a dietary dose of 16mg/kg. No increase in the incidence of liver tumors was observed in mice at a dietary dose of 4mg/kg. The clinical significance of these findings during long-term use of cetirizine is not known. Cetirizine was not mutagenic in the Ames test, and not clastogenic in the human lymphocyte assay, the mouse lymphoma assay, and *in vivo* micronucleus test in rats. In a fertility and general reproductive performance study in mice, cetirizine did not impair fertility at an oral dose of 64mg/kg.

Pediatric Under 2 Years Old:

The safety and effectiveness of cetirizine in pediatric patients under the age of 2 years have not yet been established.

Pregnancy and Lactation:

Safe uses of cetirizine during pregnancy and lactation have not been established, therefore it is not recommended.

SIDE EFFECTS:

There have been occasional reports of mild and transient subjective side-effects such as, headache, dizziness, drowsiness, agitation, dry mouth and gastrointestinal discomfort. In objective tests of psychomotor function, the incidence of sedation with cetirizine was similar to that of placebo. Occasionally symptoms of hypersensitivity have been reported.

DRUG INTERACTIONS:

To date there are no known interactions with other drugs. Nevertheless, Axcil Cetirizine should be used with caution if sedatives are also being taken.

OVERDOSAGE AND TREATMENT :

In initial observation, overdosage of Cetirizine may cause restlessness and irritability, followed by drowsiness. Should overdose occur, treatment should be symptomatic or supportive, taking into account any concomitantly ingested medications. In the case massive overdosage, gastric lavage should be performed as soon as possible. Usual supportive measures should be provided and routine observation carried out regularly.

STORAGE:

Store below 30°C. Protect from light.

KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK - KANAK

PACKAGE QUANTITIES:

Available in blister pack of 10 x 10's.

Further information can be obtained from pharmacist, physician or the manufacturer.



Product Registration Holder & Manufacturer

Kotra Pharma (M) Sdn. Bhd.

No.1, 2 & 3, Jalan TTC 12, (90082-V)

Cheng Industrial Estate,
75250 Melaka, Malaysia.

MAL20040892AZ

REGN - 280720