



CETIRIZINE SYRUP

COMPOSITION:

Each 5ml contains
Cetirizine Dihydrochloride 5mg
Methylparaben 0.1% w/v as preservative.

PRESENTATION:

Clear, colourless syrup with banana flavour.

INDICATIONS:

For symptomatic treatment of seasonal rhinitis, perennial allergic rhinitis and urticaria of allergic origin.

PHARMACOLOGY:*Mechanism of Action:*

Cetirizine is an antihistamine as its principal effects are mediated via selective inhibition of peripheral H1 receptors.

Pharmacokinetics:

Cetirizine is rapidly absorbed from the gastrointestinal tract after oral administration, peak plasma concentrations being attained in about 1 hour. It is highly bound to plasma proteins and has an elimination half-life of about 11 hours. Cetirizine has been detected in breast milk. It is excreted primarily in the urine mainly as unchanged drug.

DOSAGE AND ADMINISTRATION:

Adult : 10ml (10mg) once daily.

Children above 6 years : 5ml (5mg) twice daily
or 10ml (10mg) once daily.

Children 2-6 years : 2.5ml (2.5mg) twice daily
or 5ml (5mg) once daily.

Children 1-2 years : 2.5ml (2.5mg) twice daily.

In patients with renal insufficiency, dosage should be reduced to half the daily dose.

CONTRAINDICATION:

Contraindicated in those patients with known hypersensitivity to any of its ingredients. Cetirizine should not be administered to pregnant women during the first 3 month of pregnancy and women who are breastfeeding.

PRECAUTION:

Activities Requiring Mental Alertness : In clinical trials, the occurrence of somnolence has been reported in some patients taking cetirizine; due caution should therefore be exercised when driving car or operating potentially dangerous machinery. Concurrent use of cetirizine with alcohol or other CNS depressants should be avoided because additional reductions in alertness and additional impairment of CNS performance may occur.

Carcinogenesis, Mutagenesis and Impairment of Fertility : In a 2-year carcinogenicity study in rats, cetirizine was not carcinogenic at dietary doses up to 20mg/kg. In a 2-year carcinogenicity study in mice, cetirizine caused an increased incidence of benign liver tumors in males at a dietary dose of 16mg/kg. No increase in the incidence of liver tumors was observed in mice at a dietary dose of 4 mg/kg. The clinical significance of these findings during long-term use of cetirizine is not known. Cetirizine was not mutagenic in the Ames test, and not clastogenic in the human lymphocyte assay, the mouse lymphoma assay, and *in vivo* micronucleus test in rats. In a fertility and general reproductive performance study in mice, cetirizine did not impair fertility at an oral dose of 64 mg/kg.

Pediatric Under 2 Years Old : The safety and effectiveness of cetirizine in pediatric patients under the age of 2 years have not yet been established unless when necessary it should be used under supervision of physician.

Pregnancy and Lactation : Safe uses of cetirizine during pregnancy and lactation have not been established, therefore it is not recommended.

SIDE EFFECTS:

There have been occasional reports of mild and transient subjective side-effects such as, headache, dizziness, drowsiness, agitation, dry mouth and gastrointestinal discomfort. In objective tests of psychomotor function, the incidence of sedation with cetirizine was similar to that of placebo. Occasionally symptoms of hypersensitivity have been reported.

DRUG INTERACTIONS:

To date there are no known interactions with other drugs. Nevertheless, Axcel Cetirizine should be used with caution if sedatives are also being taken.

OVERDOSAGE AND TREATMENT:

In initial observation, overdosage of cetirizine may cause restlessness and irritability, followed by drowsiness. Should overdose occur, treatment should be symptomatic or supportive, taking into account any concomitantly ingested medications. In the case of massive overdosage, gastric lavage should be performed as soon as possible. Usual supportive measures should be provided and routine observation carried out regularly.

STORAGE:

Keep container well closed. Store below 30°C. Protect from light.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK****PACK QUANTITIES:**

Available in 60ml & 100ml PET bottle.
Not all pack sizes may be marketed.

Further information can be obtained from pharmacist, physician or the manufacturer.

MAL20040893AZ

Product Registration Holder & Manufactured By:



Kotra Pharma (M) Sdn. Bhd.
No.1, 2 & 3, Jalan TTC 12, (90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

PMSFAAXC07900 - 130825(06)
Date of revision : 2108/25