

"PBF" FUSIDIC ACID CREAM 2% W/W

Active Ingredient:

Each gram of "PBF" Fusidic Acid Cream 2% w/w contains 20mg Fusidic Acid.

Preservative(s):

Potassium Sorbate	0.2% w/w
Butylated Hydroxyanisole	0.02% w/w
Tocopherol Acetate	1% w/w

Excipients:

Cetyl Alcohol	Propylene Glycol
Polysorbate 60/ Tween 60	White Petrolatum
Hydroxyethylcellulose	Butylated Hydroxyanisole
Purified Water	Tocopherol Acetate (Vitamin E)
Potassium Sorbate	Liquid Paraffin

Product Description:

"PBF" Fusidic Acid Cream 2% w/w is a white to off-white cream.

Pharmacodynamics:

Fusidic Acid is an antibiotic with powerful antibacterial activity against a number of gram-positive organisms. Staphylococci, including the strains resistant to penicillin or to other antibiotics, are particularly susceptible to Fusidic Acid. Besides pronounced antibacterial activity against the majority of organisms responsible for skin infections, "PBF" Fusidic Acid Cream 2% w/w also possesses unique ability for penetration to the focus of infection, even when applied to intact skin.

Pharmacokinetics:

The absorption of topical applied Fucidin is about 2%. Fucidin is bound reversibly to plasma proteins, the binding being about 95%. Fucidin is metabolized in the liver and metabolites are excreted with the bile.

Indications:

"PBF" Fusidic Acid Cream 2% w/w is indicated for treatment of skin infections caused by staphylococci, streptococci, Propionibacterium acnes, Corynebacterium minutissimum, and other organisms sensitive to fusidic acid. The most important indications being:

<u>Impetigo</u>	<u>Boils</u>	<u>Carbuncles</u>	<u>Paronychia</u>
Infected wounds	Sycosis barbae	Hidradenitis	Erythrasma
Folliculitis			

"PBF" Fusidic Acid Cream 2% w/w is an invisible, non-staining preparation, which is cosmetically acceptable for treatment of face and scalp infections.

Recommended Dose:

"PBF" Fusidic Acid Cream 2% w/w is applied to the affected area 2-3 times daily, generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of this disease.

Route of Administration:

For external (Topical) use only.

Contraindications:

Infections caused by non-susceptible organisms. Patients with hypersensitivity to any of the ingredients.

Warnings and Precautions:

"PBF" Fusidic Acid Cream 2% w/w is a ready-for-use preparation and should, therefore, not be mixed with other ingredients or preparations.

Interactions with Other Medicaments:

There is no information about "PBF" Fusidic Acid Cream 2% w/w interacts with other drugs.

Pregnancy and Lactation:

There is inadequate evidence of safety in human pregnancy. Animal studies and many years of clinical experience have suggested that fusidic acid is devoid of teratogenic effects. Safety in nursing mothers has not been established.

Side Effects:

"PBF" Fusidic Acid Cream 2% w/w is well tolerated, and there is an extremely low frequency of hypersensitivity reactions.

Symptoms and Treatment of Overdosage:

If patient accidentally apply too much cream or a small quantity of the cream is accidentally swallowed, it is unlikely to cause any harm. If a larger quantity of the cream is accidentally swallowed, the patient should contact their doctor or pharmacist for advice. If patient forget to use this medicine, they need to use it as soon as they remember. The next use this medicine at the usual time. Further questions regarding the use of this medicine can be asked to doctor or pharmacist.

Effects on Ability to Drive and Use Machine:

Driving and using machines

Usually your medicine will have very little effect on your ability to drive or use machines. Check with your doctor if you feel any side effect that may stop you from driving or using machines.

Instructions for Use:

This medicine is only for using on your skin or the skin of a child. Do not swallow it. Do not put it inside your body.

Remove the cap. Check the seal is not broken before you first use the cream. Then push the spike in cap through the seal on the tube.

Always wash your hand before using "PBF" Fusidic Acid Cream 2% w/w. Rub the medicine gently on the skin. If you use it on the face, be careful to avoid the eyes area.

Unless you are using the cream to treat your hands, always wash your hands after using "PBF" Fusidic Acid Cream 2% w/w.

Storage Condition: Store below 30°C. Protect from light.

Shelf Life: 3 years.

Package: 5g/ tube.

MANUFACTURED BY:
 **PANION & BF BIOTECH INC.**
No. 266 Singlong Rd.,
Pingzhen Dist., Taoyuan City 32460, Taiwan, R.O.C.

Product Registration Holder:

MALAYSIA:
Sunward Pharmaceutical Sdn. Bhd.
No. 9, 11 & 17, Jalan Kempas 4,
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

SINGAPORE:
Sunward Pharmaceutical Pte. Ltd.
11, Wan Lee Road,
Singapore 627943

Revised Date: December 3, 2019