



MUPRIN OINTMENT

DESCRIPTION: A white ointment containing 2.0% w/w Mupirocin.

PHARMACODYNAMICS: Mupirocin inhibits bacterial protein synthesis by binding to isoleucyl-transfer RNA synthetase. Due to this particular mode of action and its unique molecular structure, Mupirocin does not show any cross-resistance with other antibiotics. It is mainly active against Gram-positive aerobes, most strains of staphylococci including methicillin-resistant and multiple resistant staphylococcus aureus and streptococci. It is also active against some Gram-negative bacterium.

PHARMACOKINETICS: Mupirocin is rapidly inactivated after absorption and its systemic level is undetectable. Virtually no systemic absorption (< 1.1 nanograms per mL of whole blood) following application to lower arm of normal males with occlusion for 24 hours.

INDICATIONS: Bacterial skin infections.

CONTRAINDICATIONS: Hypersensitivity to Mupirocin, polyethylene glycol or any other constituent in the ointment base.

DRUG INTERACTIONS: Unknown.

ADVERSE EFFECTS: Mupirocin is usually well tolerated but local reactions such as burning, stinging and itching may occur after the application of Mupirocin to the skin. Muprin Ointment has not been reported to cause contact sensitization or photosensitivity reactions. The polyethylene glycol vehicle in Muprin Ointment may irritate broken skin or mucous membranes. Muprin Ointment may cause some unwanted effects. The following side effects may go away during treatment as your body adjusts to the medicine. However, check with your doctor if any of these persist or are bothersome.

Less frequently reported unwanted effects include dry skin; skin burning, itching, pain, rash, redness, stinging or swelling; headache; nausea.

Rarely reported are abdominal pain; dizziness; secondary wound infection; ulcerative stomatitis (sores in mouth or on lips).

PRECAUTIONS/WARNINGS: Not for ophthalmic or intranasal use. Avoid contact with eyes: use with caution if there is evidence of moderate or severe renal impairment. Check with physician or pharmacist if there is no improvement within 3 to 5 days.

Pregnancy: Topical Mupirocin has not been studied in pregnant women. However, this medication has not been shown to cause birth defects or other problems in animal studies.

Breast feeding: It is not known whether topical mupirocin passes into the breast milk. However, this medicine is unlikely to pass into the breast milk in large amounts since very little Mupirocin is absorbed into the mother's body when applied to the skin.

Children: Safety and effectiveness of mupirocin ointment have not been established in children up to 2 months of age.

Older adults: No overall difference in safety and efficacy was observed in patients over 65 years of age.

RECOMMENDED DOSAGE: Adult and children: Apply one to three times daily for up to 10 days.

SYMPTOMS AND TREATMENT FOR OVERDOSE AND ANTIDOTE(S): Less frequently reported unwanted effects include: dry skin, skin burning, itching, pain, rash, redness, stinging or swelling, headache and nausea. Rarely reported are: abdominal pain, dizziness, secondary wound infection, ulcerative stomatitis (sores in mouth or on lips).

PACKING: 5g and 15g collapsible aluminium tubes.

STORAGE: Keep container well closed. Protect from strong light. Store below 30oC. Keep out of reach of children.

RECOMMENDED SHELF-LIFE : 2 years

Manufactured by :

HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohd. 6, Bandar Sultan Suleiman,
42000 Port Klang, MALAYSIA.

Revision date : October 2013



MUPRIN 軟膏

產品內容: 本品為白色軟膏，含有2.0% W/W 莫匹羅星。

產品作用: 莫匹羅星能夠藉由與異白胺醯-轉移核醣核酸合成酶的結合來抑制細菌蛋白質的合成，由於這種藥品具有特殊的作用機轉與獨特的分子結構，因此並無呈現與其他抗生素具有交叉抗藥性。其主要對抗革蘭氏陽性好氧菌、大部分的葡萄球菌（包括對於methicillin有抗藥性與多重抗藥性的金黃色葡萄球菌）、以及鏈球菌等，也可以有效對抗某些革蘭氏陰性菌。

藥理學機轉: 莫匹羅星可以在吸收後快速地被轉化成無活性的代謝物，同時其全身濃度也低至不可測得，而在塗抹於正常男性下臂並包覆24小時後，幾乎沒有發現全身性的吸收（全血中<每毫升1.1毫微克）。

適應症: 皮膚細菌性感染

禁忌症: 已知對於莫匹羅星、丙烯乙二醇、或軟膏基劑配方中任何一種成分過敏者。

藥品交互作用: 不詳。

藥品不良反應: 莫匹羅星通常耐受性良好，然而在局部使用莫匹羅星於皮膚後，仍然有可能會產生灼熱、刺痛、搔癢等不良反應。Muprin軟膏尚未有引發接觸性過敏或光過敏反應的病例，而Muprin軟膏中的丙烯乙二醇基劑則有可能會刺激皮膚傷口或黏膜。Muprin軟膏可能會造成一些不良反應。而隨著治療的進行，以下這些不良反應可能會在身體適應藥品之後消失。然而，如果這些反應持續發生或令人困擾，請與醫師聯絡。

較少見的不良反應包括：皮膚乾燥、皮膚灼熱、搔癢、疼痛、出疹、發紅、刺痛、或腫脹等，此外還包括頭痛與噁心。

極為少見的不良反應則包括：腹痛、頭暈、傷口二度感染、潰瘍性口炎（口腔或唇部的疼痛）等。

注意事項/警語: 不可使用於眼部或鼻腔內部。避免接觸眼部。使用於中度或嚴重功能不全病患身上時，應格外謹慎。如果在使用3至5天之後症狀尚未有所改善，請與醫師或藥師聯絡。

懷孕期間: 尚未有懷孕期間局部使用莫匹羅星的研究報告。然而，在動物試驗當中，尚未有本藥品造成胎兒缺陷或其他病症的病例。

哺乳期間: 莫匹羅星是否會進入母乳尚且未知，然而，由於莫匹羅星局部使用於母體皮膚時，僅會產生極少的吸收，因此本藥品較不可能會大量進入母乳當中。

孩童: 莫匹羅星軟膏使用於2歲以下孩童的安全性與有效性資料尚未建立。

年長者: 使用在65歲以上長者的安全性與有效性在整體來說並無不同。

建議劑量: 成人與孩童：每日使用一至三次，最多使用十日。

服用過量之症狀、治療方式、以及解毒劑: 較少見的不良反應包括：皮膚乾燥、皮膚灼熱、搔癢、疼痛、出疹、發紅、刺痛、或腫脹等，此外還包括頭痛與噁心。

極為少見的不良反應則包括：腹痛、頭暈、傷口二度感染、潰瘍性口炎（口腔或唇部的疼痛）等。

包裝: 5公克與15公克鋁製軟管裝。

儲存方式: 請緊密關好容器。避免強光直射。儲存於攝氏30度以下。請放置於孩童無法拿取之處。

建議有效期限: 2年

製造廠商: HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohd. 6, Bandar Sultan Suleiman,
42000 Port Klang, MALAYSIA.