

———— IDAMAN PHARMA MANUFACTURING ————

CITREX COUGH SYRUP 100MG/5ML

PRODUCT DESCRIPTION

A clear, orange coloured solution with a sweet, pleasant smell.

COMPOSITION

Each 5ml contains:

- Active Ingredient: Guaifenesin 100mg
- Preservative: Sodium Benzoate 5mg

ROUTE OF ADMINISTRATION

Oral

PHARMACODYNAMICS

Guaifenesin is an expectorant which is thought to act by irritating the gastric mucosa and subsequently stimulating respiratory tract secretions. This increase in fluid increases the volume and decreases the viscosity of bronchial secretions.

PHARMACOKINETICS

Guaifenesin is well absorbed from the gastrointestinal tract. It is metabolised and then excreted in the urine. The half-life in plasma is approximately 1 hour.

INDICATIONS

Citrex Cough Syrup 100mg/5ml helps to:

- Relieve of chesty coughs.
- Loosen phlegm and thin bronchial secretion.

RECOMMENDED DOSAGE

The elderly, adults and children over 12 years: Take 10ml four times daily.

Children

6 – 12 years: Take 5ml four times daily

2 – 6 years: Take 2.5ml four times daily

Under 2 years: Not recommended

CONTRAINDICATIONS

Hypersensitivity to Guaifenesin or any of the ingredients/components of the product.

WARNINGS AND PRECAUTIONS

Do not exceed the stated dose. Please consult with a doctor or pharmacist before taking this medicine if the following occurs:

- Constant cough for more than 7 days or cough recurs or is accompanied by a fever, rash or persistent headache.
- Persistent cough that may be caused by asthma.
- Cough which brings up a lot of mucus (phlegm).
- Suffer from liver or kidney problems.
- Concomitant use of cough suppressants and other cough and cold medicine.

INTERACTIONS WITH OTHER MEDICAMENTS

Diagnostic Interference:

Guaifenesin has been shown to produce a colour interference with certain clinical laboratory determinations of 5-hydroxyindoleacetic acid (5-HIAA) and vanillylmandelic acid (VMA).

Guaifenesin could interfere with the diagnosis of carcinoid syndrome. Patients being evaluated for carcinoid syndrome should therefore discontinue any preparation containing Guaifenesin for 24 hours before collection of urine specimens for the determination of 5-hydroxyindoleacetic acid (5-HIAA).

It is also not recommended to take Guaifenesin with other cough suppressants as they may interfere with the action of this medicine and expose patients to unnecessary adverse effects.

PREGNANCY AND LACTATION

Available evidence and/or expert consensus is inconclusive or is inadequate for determining infant risk when used during pregnancy and breastfeeding. Guaifenesin is excreted in breast milk in small amounts. Weigh the potential benefits of drug treatment against potential risks before taking this drug during pregnancy and breastfeeding. It is recommended to consult with doctor or pharmacist before taking this medicine.

SIDE/ADVERSE EFFECTS

Gastrointestinal discomfort, nausea, and vomiting have occasionally been reported with Guaifenesin, particularly in very large doses. Others include dizziness, diarrhoea, headache and rash (including urticaria, pruritus and hypersensitivity).

SYMPTOMS & TREATMENT OVERDOSE

The effects of acute toxicity from Guaifenesin may include gastrointestinal discomfort, nausea, drowsiness and vomiting. When taken in excess, Guaifenesin may cause renal calculi. In the event of overdosage, discontinue medication and seek medical help immediately. Treatment should be symptomatic and supportive.

EFFECTS ON ABILITY TO DRIVE & USE MACHINE

Guaifenesin has no or negligible influence on the ability to drive or operate machinery. However, if patients do feel unwell, do not drive or use machinery.

PACK SIZES

Available in a bottle of 120ml syrup.

STORAGE CONDITION

Store below 30°C. Keep container tightly closed. Do not use if temper evident seal is broken or missing.

SHELF LIFE

Product should not be consumed beyond 2 years of the manufacturing date imprinted on the product packaging.

REGISTRATION NUMBER

MAL_____

KEEP MEDICINES OUT OF REACH OF CHILDREN JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 01

Revision Date : 3rd April 2020

PRODUCT REGISTRATION HOLDER AND MANUFACTURER

IDAMAN PHARMA MANUFACTURING SDN BHD (200401023395)

LOT 24 & 25, JALAN PERUSAHAAN LAPAN,
BAKAR ARANG INDUSTRIAL ESTATE, 08000 SUNGAI PETANI,
KEDAH DARUL AMAN, MALAYSIA.