



VIFUSxx-0\_6

## CP PHARMA FUSIDIC ACID CREAM 2% W/W

### DESCRIPTION

White, opaque cream.

### COMPOSITION

Fusidic Acid (micronised) 2.03% w/w  
Equivalent to Fusidic Acid 2% w/w

### PHARMACODYNAMICS

Pharmacotherapeutic group: Other antibiotics for topical use, ATC code: D06AX01.

Fusidic acid is a potent antibacterial agent. Fusidic acid and its salts show fat and water solubility and strong surface activity and exhibit unusual ability to penetrate intact skin. Concentrations of 0.03 - 0.12 mcg fusidic acid per ml inhibit nearly all strains of Staphylococcus aureus. Topical application of fusidic acid is also effective against streptococci, corynebacteria, neisseria and certain clostridia.

### PHARMACOKINETICS

In vitro studies show that fusidic acid can penetrate intact human skin. The degree of penetration depends on factors such as the duration of exposure to fusidic acid and the condition of the skin. Fusidic acid is excreted mainly in the bile with little excreted in the urine.

### INDICATIONS

CP Pharma Fusidic Acid Cream 2% is indicated for the treatment of skin infections caused by Staphylococci, Streptococci, Propionibacterium acnes, Corynebacterium minutissimum and other organisms sensitive to fusidic acid. The most important indication being: Impetigo, Boils, Carbuncles, Paronychia, Infected wounds, Sycosis barbae, Hidradenitis, Erythrasma, Folliculitis.

### CONTRAINDICATION

Hypersensitivity to the active substance or to any of the excipients in this product.

### WARNINGS AND PRECAUTIONS

Bacterial resistance among staphylococcus aureus has been reported to occur with the use of topical CP Pharma Fusidic Acid Cream 2%. As with all antibiotics, extended or recurrent use may increase the risk of developing antibiotic resistance.

Extended or recurrent use may increase the risk of developing contact sensitisation.

CP Pharma Fusidic Acid Cream 2% should therefore be used with care when applied in the proximity of the eyes.

### DRUG INTERACTION

No interaction studies have been performed. Interactions with systemically administered medicinal products are considered minimal as the systemic absorption of topical CP Pharma Fusidic Acid Cream 2% is negligible.

### PREGNANCY AND LACTATION

#### Pregnancy

No effects during pregnancy are anticipated, since systemic exposure to topically-applied fusidic acid/sodium fusidate is negligible. Topical CP Pharma Fusidic Acid Cream 2% can be used during pregnancy.

#### Breast-feeding

No effects on the breast-fed new-born/infant are anticipated since the systemic exposure of topically-applied fusidic acid/sodium fusidate to the breast-feeding woman is negligible. Topical CP

Pharma Fusidic Acid Cream 2% can be used during breast-feeding but it is recommended to avoid applying topical CP Pharma Fusidic Acid Cream 2% on the breast.

### MAIN SIDE/ADVERSE EFFECTS

The most frequently reported adverse reactions during treatment are various skin reactions such as pruritus and rash, followed by application site conditions such as pain and irritation.

**Immune system disorders (rare):** Hypersensitivity

**Eye disorders (rare):** Conjunctivitis

**Skin and subcutaneous tissue disorders (uncommon):** Dermatitis (including dermatitis contact, eczema), Rash, Pruritus, Erythema.

**Skin and subcutaneous tissue disorders (rare):** Angioedema, Urticaria, Blister

**General disorders and administration site conditions (uncommon):** Application site pain (including skin burning sensation), Application site irritation.

#### Paediatric population

Frequency, type and severity of adverse reactions in children are expected to be the same as in adults.

#### Effects on ability to drive and use machines

CP Pharma Fusidic Acid Cream 2% administered topically has no or negligible influence on the ability to drive and use machines.

### SYMPTOMS AND TREATMENT OF OVERDOSE

Overdose is unlikely to occur.

Unless hypersensitivity to Fusidic Acid or any of the excipients exists, accidental ingestion of CP Pharma Fusidic Acid Cream is unlikely to cause any harm. The total quantity of fusidic acid will usually not exceed the approved total daily oral dose of fusidic acid containing products except in children aged less than 1 year and weighing  $\leq 10$  kg. Although in this instance a child of this particular age group is unlikely to ingest a whole tube of CP Pharma Fusidic Acid Cream. The concentration of the excipients is too low to constitute a safety risk.

### DOSAGE AND ADMINISTRATION

FOR EXTERNAL USE ONLY.

CP Pharma Fusidic Acid Cream 2% w/w is applied to the affected area 2-3 times daily generally for a period of 7 days, except in treatment of acne where therapy should be extended for a longer period according to the nature of this disease.

Note: The information given here is limited. For further information, consult your doctor or pharmacist.

Storage: Store below 30°C.

Presentation/Packing: Cream of 15 g in an aluminium tube.

Product registration holder: HOVID Pharmacy Sdn. Bhd., 121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak, Malaysia.

Manufactured by: HOVID Bhd., 121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak, Malaysia.

For: CARING Pharmacy Retail Management Sdn Bhd (757411-U) No. 1, Jalan 51/203A, Kawasan Perindustrian Tiong Nam, Seksyen 51, 46050 Petaling Jaya.

Information date: April 2020

XXX