Synergy Meditech
Malaysia

For the use of a Registered Medical Practitioner or a Hospital

PIROBACT

MUPIROCIN OINTMENT USP 2%w/w

Category: Topical Antibiotic
Composition:
Each gram of ointment contains:
Mupirocin USP .....20 mg
In a water soluble ointment base.....q.s

Description:
White to off white translucent ointment.

Indications:
Bacterial skin infections, e.g. impetigo, folliculitis and furunculosis.

Clinical Pharmacology
Pharmacodynamics
Mechanism of action
Mupirocin is a novel antibiotic produced through fermentation by *Pseudomonas fluorescens.* Mupirocin inhibits isoleucyl transfer-RNA synthetase, thereby arresting bacterial protein synthesis.

Mupirocin has bacteriostatic properties at minimum inhibitory concentrations and bactericidal properties at the higher concentrations reached when applied locally.

Low-level resistance in staphylococci is thought to result from point mutations within the usual staphylococcal chromosomal gene (ileS) for the target isoleucyl tRNA synthetase enzyme. High-level resistance in staphylococci has been shown to be due to a distinct, plasmid encoded isoleucyl tRNA synthetase enzyme.

Intrinsic resistance in Gram negative organisms such as the Enterobacteriaceae could be due to poor penetration of the outer membrane of the Gram-negative bacterial cell wall.
Due to its particular mode of action, and its unique chemical structure, mupirocin does not show any cross-resistance with other clinically available antibiotics.

Pharmacokinetics:
After topical application of Mupirocin Ointment, mupirocin is only very minimally absorbed systemically and that which is absorbed is rapidly metabolised to the antimicrobially inactive metabolite, monic acid. Penetration of mupirocin into the deeper epidermal and dermal layers of the skin is enhanced in traumatised skin and under occlusive dressings.

Elderly:
No restrictions unless there is evidence of moderate or severe renal impairment

DOSAGE
Adults and children:
PIROBACT (Mupirocin Ointment USP 2% w/w) should be applied to the affected area 2 to 3 times a day, for up to 10 days, depending on the response.
The area may be covered with a dressing or occluded if desired.
Do not mix with other preparations as there is a risk of dilution, resulting in a reduction of the antibacterial activity and potential loss of stability of the mupirocin in the ointment.

Side effects:
Common Side effects
Burning localised to the area of application

Uncommon Side effects
Itching, erythema, stinging and dryness localised to the area of application. Cutaneous sensitisation reactions to mupirocin or the ointment base.

CONTRAINDICATIONS
Hypersensitivity to the active substance(s) or to any of the excipients.

Mupirocin Ointment formulation is not suitable for ophthalmic or intranasal use

Precaution:
Should a possible sensitisation reaction or severe local irritation occur with the use of Mupirocin Ointment, treatment should be discontinued, the product should be washed off and appropriate therapy instituted.

As with other antibacterial products, prolonged use may result in overgrowth of non-susceptible organisms. Pseudomembranous colitis has been reported with the use of antibiotics and may range in severity from mild to life-threatening. Therefore, it is important to consider its diagnosis in patients who develop diarrhoea during or after antibiotic use. Although this is less likely to occur with topically applied mupirocin, if prolonged or significant diarrhoea occurs or the patient experiences abdominal cramps, treatment should be discontinued immediately and the patient investigated further.

Renal Impairment
Polyethylene glycol can be absorbed from open wounds and damaged skin and is excreted by the kidneys. In common with other polyethylene glycol-based ointments, Mupirocin Ointment should not be used in conditions where absorption of large quantities of polyethylene glycol is possible, especially if there is evidence of moderate or severe renal impairment.

Mupirocin Ointment is not suitable for:
- ophthalmic use;
- intranasal use (in neonates or infants);
- use in conjunction with cannulae;
- at the site of central venous cannulation.

Avoid contact with the eyes. If contaminated, the eyes should be thoroughly irrigated with water until the ointment residues have been removed.

DRUG INTERACTION
No interaction studies have been performed.

Pregnancy and Lactation
Pregnancy:
No evidence of harm to the foetus in animals.
Mupirocin Ointment should only be used in pregnancy when the potential benefits outweigh the possible risks of treatment.
Breast-feeding:
It is unknown whether mupirocin is excreted in human milk. If a cracked nipple is to be treated, it should be thoroughly washed prior to breast-feeding.

Effects on Ability to Drive and Use Machine
Mupirocin 2% w/w Ointment has no or negligible influence on the ability to drive and use machines.

Overdosage:
Symptoms
There is currently limited experience with overdosage of mupirocin.

Management
The toxicity of mupirocin is very low. In the event of accidental ingestion of the ointment, symptomatic treatment should be given.
In case of erroneous oral intake of large quantities of the ointment, renal function should be closely monitored in patients with renal insufficiency because of the possible side effects of polyethylene glycol.
There is no specific treatment for an overdose of mupirocin. In the event of overdose, the patient should be treated supportively with appropriate monitoring as necessary. Further management should be as clinically indicated or as recommended by the national poisons centre, where available.

Storage:
Store in a cool dry place below 30°C, protect from light. Do not freeze.
Keep out of reach of children

PRESENTATION
Box containing 5g or 15g ointment in printed polyethylene lamitube sealed with aluminium foil and closed with a white polypropylene cap.

Manufactured by

Swiss Garnier LIFE SCIENCES

No. 21-23, Industrial Area, Mehatpur, Dist: UNA,
Himachal Pradesh - 174 315. INDIA

Product Registration Holder :
Synergy Meditech (M) Sdn Bhd,
A-17-7, BLOCK A, JAYA ONE, NO. 72A,
JALAN PROFESOR DIRAJA UNGKU AZIZ,
SEKSYEN 13, 46200 PETALING JAYA, MALAYSIA.

Revision date : 21 January 2025

PMLF0656R1

Brand Name	**PIROBACT**			Artwork Date	**08.07.2024**
Generic Name	**Mupirocin Ointment USP 2% w/w**				
Client/Country	**Synergy Meditech / Malaysia**			Language	**English**
ERP Code	**-------**	Artwork Code	**PMLF0656R1**	Item Type	**Leaflet**
Specification	**60 GSM Maplitho Paper with Folding**				
Size	**120 x 180 mm**	Packing	**NA**	Software	Coral Draw X8
Colours	■ **Black**				
Changes Made:					

Swiss Garnier LIFE SCIENCES