

Front - 80mm

Back - 80mm

Unimed's ChlorSept 2% with IPA Antiseptic Solution

Composition and Strength of Active Ingredients:

10% v/v of Chlorhexidine gluconate 20% solution equivalent to Chlorhexidine gluconate	2.0% v/v
Isopropyl alcohol	70% v/v

Product Description

A clear, pink color solution.

Pharmacodynamics

Chlorhexidine gluconate is a type of chemical agent used to destroy or inhibit the growth of pathogenic micro-organisms in the non-sporing or vegetative state. Disinfectants not necessarily kill all micro-organisms but may reduce to the level of limiting or preventing infection. The combination with alcohols may have a synergistic effect and the antimicrobial efficacy of chlorhexidine is much lower and slower with possibility of occurrence in bacterial resistance. Disinfection of catheter insertion sites with aqueous chlorhexidine 2% has been reported to be associated with fewer local and systemic infections. Chlorhexidine is a bisguanide antiseptic and disinfectant that is bactericidal or bacteriostatic with a wide range of Gram-positive and Gram-negative bacteria. It is more effective against Gram-positive than Gram-negative bacteria, and some species of *Pseudomonas* and *Proteus* have low susceptibility. It is relatively ineffective against mycobacteria. Chlorhexidine inhibits some viruses and is active against some fungi. It is inactive against bacterial spore at room temperature. Chlorhexidine is most active at a neutral or slightly acid pH.

Isopropyl alcohol is a bactericidal agent that acts by the denaturation of proteins. It is a rapidly bactericidal and a fast-acting broad-spectrum antiseptic, but is not considered persistent. At concentrations of 70% v/v, it is a much effective antibacterial preservative than ethanol (95%). The bactericidal effect of aqueous solutions increases steadily as the concentration approaches 100% v/v. However, isopropyl alcohol is ineffective against bacterial spores.

Pharmacokinetics

Chlorhexidine is cationic nature causing it to bind strongly to skin mucosa and other tissues and is thus very poorly absorbed. As a consequence, no general pharmacological studies on chlorhexidine available with minimal effects on internal organs. No detectable blood levels or insignificant amount was found in man subject to oral use and percutaneous absorption. Chlorhexidine is poorly absorbed from the gastrointestinal tract.

As for the pharmacokinetic properties, there is little absorption of isopropyl alcohol through intact skin. It is readily absorbed from the gastrointestinal tract and may be slowly absorbed through intact skin. Isopropyl alcohol is metabolized more slowly than ethanol, primarily to acetone. Metabolites and unchanged isopropyl alcohol are mainly excreted in the urine.

Indications

As an antiseptic for the skin prior to invasive procedures.

Recommended Dose

Skin Preparation: Use minimum amount of Unimed's ChlorSept 2% with IPA Antiseptic Solution and allow to dry. The solution is used undiluted.

Route of Administration

Topical

Contraindications

Known hypersensitivity to Unimed's ChlorSept 2% with IPA Antiseptic Solution or any of its components, especially those with a known history of Chlorhexidine-related allergic reactions.

Warning and Precautions

The solution is flammable. Do not use electrocautery procedures or other ignition sources until the skin is completely dry.

Unimed's ChlorSept 2% with IPA Antiseptic Solution contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of

chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. Unimed's ChlorSept 2% with IPA Antiseptic Solution should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Interactions with Other Medicaments

Chlorhexidine salts are incompatible with soaps and other anionic materials. Activity may be reduced in the presence of suspending agents such as alginates and tragacanth, insoluble powders such as kaolin, and insoluble compounds of calcium, magnesium, and zinc.

As for the isopropyl alcohol, it should not be brought into contact with some vaccines and skin test injections (patch tests) and you should consult the vaccine manufacturer's literature when in doubt. Also, isopropyl alcohol is known to be incompatible with oxidizing agents such as hydrogen peroxide and nitric acid, which may cause decomposition. Isopropyl alcohol may be salted out from aqueous mixtures by the addition of sodium chloride, sodium sulfate, and other salts, or by the addition of sodium hydroxide.

Pregnancy & Lactation

No special precautions need to be taken when used in pregnancy and lactation. Thus, Unimed's ChlorSept 2% with IPA Antiseptic Solution can be used during pregnancy or during breast-feeding.

Side Effects

Skin sensitivity to Unimed's ChlorSept 2% with IPA Antiseptic Solution has been reported rarely with occasional occurrence. Immune system disorders. Frequency not known: Hypersensitivity including anaphylactic shock.

Symptoms and Treatment of Overdose

There are no reports of overdose with this product. However, Chlorhexidine Gluconate taken orally is poorly absorbed into the blood. Accidental ingestion may require appropriate gastric lavage using milk, egg white, gelatine or mild soap. Haemolysis has been reported following ingestion or accidental intravenous administration which necessitate blood transfusion.

Effects on Ability to Drive and Use Machine

Unimed's ChlorSept 2% with IPA Antiseptic Solution has no influence on ability to drive and use machines.

Storage Condition

Keep container tightly closed. Protect from heat and light. Store below 30°C.

Dosage Forms and Packaging Available

60 ml, 250 ml and 500 ml bottle.

Registration Number

MAL20096023XZ

Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.
Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Manufactured by

KCK Pharmaceutical Industries Sdn. Bhd.
No. 5 & 7, Hala Beringin,
Diamond Valley Industrial Park, 11960 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision

11 December 2023

PI0064-1

125mm