

VENTAMOL TABLET 2 mg

VIVEN01-M

Description

Round, light purplish uncoated tablet, bevel-edged, flat-faces, "HD" embossed and scored on the same face.

Composition

Salbutamol (as sulphate) 2 mg/tablet.

Actions and Pharmacology

Salbutamol is a direct-acting sympathomimetic amine. It acts selectively on beta-2 adrenergic receptors in the lungs to relax bronchial smooth muscle, thereby relieving bronchospasm and reducing airway resistance. It is rapidly absorbed orally, undergoes first pass hepatic metabolism and renal excretion.

Indications

For treatment of asthma, chronic bronchitis and other bronchopulmonary disorders involving bronchospasm.

Contraindications

Avoid in patients hypersensitive to sympathomimetics.

Warning and Precautions

- Caution in patients with cardiovascular disorders, including coronary insufficiency and hypertension; diabetes mellitus and hyperthyroidism.
- Check with physician immediately if difficulty in breathing persists after use of this medication or if conditions become worse.
- Tocolysis: Serious adverse reactions including death have been reported after administration of salbutamol to women in labor. In the mother, these include increased heart rate, transient hyperglycaemia, hypokalaemia, cardiac arrhythmias, pulmonary oedema and myocardial ischaemia. Increased fetal heart rate and neonatal hypoglycaemia may occur as a result of maternal administration.

Use in Pregnancy and Lactation

- Safety for use in pregnancy and lactation has not been established.

Main Side/Adverse Effects

Tremor, palpitations, slight tachycardia, nervousness, nausea.

Drug Interactions

- Co-administration of non-selective beta-adrenergic blocking agents is not recommended.
- Caution in patients receiving the following drug therapy :
Oral sympathomimetics, xanthines, monoamine oxidase (MAO) inhibitors, tricyclic antidepressants, high doses of corticosteroids.

Overdosage

Clinical features : Tremor, agitation, tachycardia, peripheral vasodilatation, chest pain, hypokalaemia, hyperglycaemia, lactic acidosis, acute psychosis.

Treat overdosage by emesis or gastric lavage, if appropriate. Administer atenolol 2.5 - 10 mg or propranolol 1 mg/min slowly for rate-related myocardial ischaemia.

Dosage and administration

Adults : Oral, 2 to 4 mg three or four times daily, or as directed.
Children : 2 to 6 years : Oral, 1 to 2 mg three or four times daily, or as directed.
Above 6 years : Oral, 2 mg three or four times daily, or as directed.

Note: The information given here is limited. For further information consult your doctor or pharmacist.

Storage : Store below 25°C. Protect from light and moisture.
Presentation/Packing : 1000's; Blisters of 10 x 10's.

Marketing authorization holder : HOVID Bhd., 121, Jalan Tunku Abdul Rahman, 30010 Ipoh, Malaysia.
Manufactured by : HOVID Bhd., Lot 56442, 7 1/2 Miles, Jalan Ipoh / Chemor, 31200 Chemor, Malaysia.

Information date: May 2012