

Important information. Please read carefully.

TULLYX

Elixir

COMPOSITION

Each 5ml contains Dextromethorphan Hydrobromide Monohydrate 15mg and Bromhexine Hydrochloride 4mg.

Methyl Hydroxybenzoate 0.08% w/v and Propyl Hydroxybenzoate 0.03% w/v as preservatives.

PHARMACODYNAMICS

Bromhexine is a mucolytic used in the treatment of respiratory disorders associated with productive cough, where it changes the structure of bronchial secretions leading to a reduction in the viscosity of sputum.

Dextromethorphan Hydrobromide is a cough suppressant used for the relief of non-productive cough. It has a central action on the cough centre in the medulla. It is also an antagonist of N-methyl-D-aspartate (NMDA) receptors. Dextromethorphan has no classical analgesic properties and little sedative activity.

PHARMACOKINETICS

Bromhexine is rapidly absorbed from the gastrointestinal tract and undergoes extensive first-pass metabolism in the liver. It is widely distributed to body tissues. About 85 to 90% of a dose is excreted in the urine mainly as metabolites. Bromhexine is highly bound to plasma proteins and has a terminal elimination half-life of up to about 12 hours. It crosses the blood-brain barrier and small amounts cross the placenta.

Dextromethorphan Hydrobromide is rapidly absorbed from the gastrointestinal tract. It is metabolized in the liver excreted in the urine as unchanged dextromethorphan and demethylated metabolites including dextrophan. It is reported to act within half an hour of administration by mouth and to exert an effect for up to 6 hours.

INDICATIONS

Tullyx helps for short term symptomatic relief of cough associated with reduction in sputum viscosity.

DOSAGE AND ADMINISTRATION

For oral use only.

Adult and children over 12 years:

10 ml (2 teaspoons), 3 times daily.

Children 6 to 12 years:

5 ml (1 teaspoon), 3 times daily.

Children 2 to 6 years:

A doctor's direction is recommended as young children have a greater chance of experiencing serious adverse effects.

The recommended dosage should not exceed 3ml per day.

Children below 2 years:

Not recommended.

Patients are advised to use a calibrated measuring device as 5 ml is a rough approximation of one teaspoon and different teaspoons differ in their volume.

CONTRAINDICATIONS

This medication is contraindicated with patients with sensitivity to bromhexine and dextromethorphan or other components of the formulation. The presence of other medical problems may affect the use of this medication. A doctor should be consulted before using this medicine if any other medical problems exist, especially: peptic ulceration, asthma, chronic bronchitis, productive cough, emphysema, diabetes, hepatic function impairment and respiration depression.

WARNINGS AND PRECAUTIONS

Bromhexine should be used with care in patients with a history of asthma and peptic ulceration. Clearance of bromhexine or its metabolites may be reduced in patients with severe hepatic or renal impairment.

Very rare cases of chronically associated severe skin impairments such as Stevens Johnson Syndrome, Toxic Epidermal Necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP) have been reported. In most cases, these could be explained by the severity of the underlying disease or concomitant administration of another drug. In the early stages of such severe skin reactions, initially only nonspecific flu-like symptoms appear, e.g. fever, arthralgia, runny nose, cough, and sore throat. If skin or mucous membrane damage occurs, seek medical advice immediately and discontinue treatment as a precaution.

Dextromethorphan should not be given to patients at risk of developing respiratory failure and caution is needed in patients with a history of asthma and should not be given during an acute attack.

This medicine may cause drowsiness and dizziness, less alert than normal or to feel a false sense of well-being. Patients should know how they react to this medicine before they drive, use machines or doing anything that could be dangerous if they are dizzy or are not alert and clearheaded. Children should be supervised to avoid potential harm in bike riding or other dangerous activities.

Not to be used in children less than 2 years of age. To be used with caution and doctor's advice in children 2 to 6 years of age.

DRUG INTERACTIONS

Concurrent use of this medication with combination products containing any of the following medications may interact with dextromethorphan: central nervous system (CNS) depressants, monoamine oxidase (MAO) inhibitors including furazolidone, phenelzine, procarbazine, selegiline and tranylcypromine; other enzyme inhibitors like amiodarone or fluoxetine or quinidine; and smoking tobacco.

PREGNANCY AND LACTATION

It is advisable not to use this medication during the first trimester of pregnancy. This medication also should be avoided during lactation.

SIDE EFFECTS

The side effects with this medication appear to be rare and may include: gastrointestinal disturbance, transient rise in serum aminotransferase values, headache, drowsiness, dizziness, sweating, skin rashes and occasionally produced cough or bronchospasm in susceptible subjects, confusion, constipation, nausea or vomiting and stomach pain.

Immune System Disorders

Frequency not known: Anaphylactic reactions including anaphylactic shock.

Skin and Subcutaneous Skin Disorders

Frequency not known: Severe skin reactions (including Stevens Johnson syndrome, Toxic epidermal necrolysis (TEN), Erythema Multiforme (EM) and Acute Generalized Exanthematous Pustulosis (AGEP).

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

Symptoms of overdose: Ataxia, blurred vision, coma, confusion, drowsiness or dizziness, respiratory depression, severe nausea or vomiting, severe unusual excitement nervousness, restlessness or irritability and urinary retention.

Treatment for overdose: Treatment is symptomatic and is directed toward the affected body system. Gastric lavage, assisted respiration, vital sign monitoring and intravenous naloxone are some of the possible prescribed treatment.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

Although Dextromethorphan has little sedative activity, nevertheless patients should be advised that it may affect performance of skilled tasks such as driving.

SHELF-LIFE

The expiry date is indicated on the packaging.

STORAGE CONDITION

Store below 30°C.

PRODUCT DESCRIPTION

Orange coloured oral solution with orange flavour.

PACKAGING AVAILABLE

Available in pack size of 60ml, 90ml and 120ml in Amber Polyethylene Terephthalate (PET) bottle.

KEEP OUT OF REACH OF CHILDREN/ JAUHI DARI KANAK-KANAK

For further information, please consult your pharmacist or physician.

Revision Date: 10-Oct-2020

Manufacturer and Product Registration Holder
Xepa-Soul Pattinson (Malaysia) Sdn Bhd
1-5 Cheng Industrial Estate, 75250 Melaka, Malaysia.