

MEDIDYL COUGH SYRUP

Each 5ml contains:-
Diphenhydramine HCl 14mg
Ammonium Chloride 135mg

Preservative:-
Sodium Benzoate 3.7mg

Pharmacodynamics:

Diphenhydramine is an ethanolanine-derivative antihistamine. It shares the action and uses of other antihistamines. It is also used as an antitussive for temporary relief of cough caused by minor throat and bronchial irritation such as may occur with common colds or inhaled irritants. It has been suggested that ammonium chloride may act as an expectorant by reflex stimulation of bronchial mucous glands resulting from irritation of gastric mucosa following oral administration. Sodium citrate is used as a source of bicarbonate when employed as alkalinizing agent.

Pharmacokinetics:

Diphenhydramine hydrochloride is well absorbed following oral administration but apparently undergoes first-pass metabolism in the liver and only about 40-60% of an oral dose reaches systemic circulation as unchanged diphenhydramine. Following oral administration of a single dose of diphenhydramine, the drug appears in plasma within 15 minutes and peak plasma concentrations are attained within 1-4 hours. Distribution of diphenhydramine into human body tissues and fluids has not been fully characterized. The terminal elimination half-life of diphenhydramine has not been fully elucidated but appears to range from 2.4-3.3 hours in healthy adults. The terminal elimination half-life reportedly is prolonged in adults with liver cirrhosis. Diphenhydramine is rapidly and apparently almost completely metabolized principally to diphenylmethoxyacetic acid which may further undergo conjugation. The drug also undergoes dealkylation to form the N-demethyl and N-dimethyl derivatives. Diphenhydramine and its metabolites are excreted principally in urine. Ammonium chloride, following oral administration is rapidly absorbed from the GI tract, complete absorption occurring within 3-6 hours. The drug is metabolized in the liver to form urea and hydrochloric acid.

Sodium citrate is an alkalinizing agent whose activity depends on conversion to bicarbonate. When oxidative activity is intact, oxidation of the citrate salt is essentially complete and less than 5% of the drug is excreted in urine unchanged.

Route of Administration:

Oral administration.

Dosage:

Adults: Two teaspoonsful every four hours not to exceed twelve teaspoonsful or 60ml in 24 hours. Children (6 to 11 years): One teaspoonful every four hours not to exceed six teaspoonsful or 30ml in 24 hours.

Take with food/milk to minimize gastric irritation. Drink a glass of water after each dose to help loosen mucus in lungs.

Indications:

It is indicated as an expectorant for control of coughs due to colds or allergy.

Warnings and Precautions:

Anti-histamines should be used with considerable caution in patients with narrow-angle glaucoma, stenosing peptic ulcer, pyloroduodenal obstruction, symptomatic prostatic hypertrophy or bladder-neck obstruction. In infants and children, especially, anti-histamines in overdose may cause hallucinations, convulsions or death. As in adults, anti-histamines may diminish mental alertness in children. In the young child, particularly, they may produce excitation. Experience with this drug in pregnant women is inadequate to determine whether there exists a potential for harm to the developing fetus. Patients should be warned about engaging in activities requiring mental alertness, such as driving a car, or operating appliances, machinery etc. Anti-histamines are more likely to cause dizziness, sedation and hypotension in elderly patients. Diphenhydramine hydrochloride has an atropine-like action and therefore should be used with caution in patients with a history of bronchial asthma, increased intraocular pressure, hyperthyroidism, cardiovascular diseases or hypertension. MAO inhibitors prolong and intensify the anti-cholinergic (drying) effects of anti-histamines. Consult a physician if symptoms persist for more than 7 days, or if high fever, skin rash or continuing headache is present with cough.

WARNING: When used for treatment of cough and cold:

- Not to be used in children less than 2 years of age.
- To be used with caution and doctor's/pharmacist's advice in children 2 to 6 years of age.

Use in Pregnancy and Lactation:

There is no adequate and controlled studies to date for the use of diphenhydramine maleate in pregnant and nursing mother. Hence the drug should be used in these situation only when clearly needed.

Contraindications:

This product should not be taken for persistent chronic cough such as occurs with smoking, asthma, emphysema, or when cough is accompanied by excessive secretions, or if the patient has epilepsy, glaucoma, or difficulty in urination due to enlargement of the prostate gland except under the advice and supervision of physician. Diphenhydramine hydrochloride has additive effects with alcohol and other CNS depressants (hypnotics, sedatives, tranquilizers). This drug should not be used in newborn or premature infants. Because of the higher risk of anti-histamine for infants generally and for newborns and premature infants in particular, anti-histamines therapy is contraindicated in nursing mothers. Anti-histamines should not be used to treat lower respiratory tract symptoms, including asthma. Anti-histamines are also contraindicated in the following conditions: Hypersensitivity to diphenhydramines of similar chemical structure. Monoamine oxidase inhibitor therapy.

Side Effects / Adverse Reactions:

The adverse reactions reported are:

- General: Urticaria, drug rash, anaphylactic shock, photosensitivity, excessive perspiration, chills, dryness of mouth, nose and throat.
- Cardiovascular system: Hypotension, headache, palpitations, tachycardia, extrasystoles.
- Hematologic system: Hemolytic anaemia, thrombocytopenia, agranulocytosis.
- Nervous system: Sedation, sleepiness, dizziness, distributed co-ordination, fatigue, confusion, restlessness, excitation, nervousness, tremor, irritability, insomnia, euphoria, paraesthesia, blurred vision, diplopia, vertigo, tinnitus, acute labyrinthitis, hysteria, neurosis, constipation.
- G.I. systems: Epigastric distress, anorexia, nausea, vomiting, diarrhoea, constipation.
- G. U. systems: Urinary frequency, difficult urination, urinary retention, early menses.
- Respiratory system: Thickening of bronchial secretions, tightness of chest and wheezing nasal stuffiness.

Symptoms and Treatment for Overdose and Antidotes:

Anti-histamine overdose reactions may vary from central nervous system depression to stimulation. Stimulation is particularly likely in children. Atropine-like signs and symptoms - dry mouth; fixed dilated pupils; flushing and gastro-intestinal symptoms may also occur. If vomiting has not occurred spontaneously, the patient should be induced to vomit. This is best done by having the patient drink a glass of water or milk which, patient should be made to gag. Precaution against aspiration must be taken especially in infants and children. If vomiting is unsuccessful, gastric lavage is indicated within 3 hours after ingestion and even later if large amounts of milk or cream were given before hand. Isotonic or 1/2 isotonic saline is the lavage solution of choice. Saline cathartics, as milk of magnesia, by osmosis draw water into bowel and therefore are valuable for their action in rapid dilution of bowel content. Vasopressors may be used to treat hypotension. Stimulants should not be used.

Effects on Ability to Drive and Use Machine:

Medidyl Cough Syrup may cause drowsiness. Do not drive or operate machinery.

Interactions with Other Medicaments:

Additive effects if used with hypnotics, sedatives and tranquilizers. Avoid drinking alcoholic beverages when taking Medidyl Cough Syrup.

Pack Size: Amber round plastic PET bottle with inner ring HDPE lock cap of 60ml and 120ml.

Shelf Life: 3 years

Storage Condition: Store at or below 30°C. Protect from light.

Description: Clear, dark brown syrup with raspberry flavour.

FURTHER INFORMATION CONCERNING THIS DRUG CAN BE OBTAINED FROM YOUR FAMILY PHYSICIAN / LOCAL GENERAL PRACTITIONER / PHARMACIST.

Manufacturer:

Sunward Pharmaceutical Sdn. Bhd.
No. 9, 11 & 17, Jalan Kempas 4
Taman Perindustrian Tampoi Indah,
81200 Johor Bahru, Johor, Malaysia.

Revised Date: 03/12/2019