

FUCON[®]

Presentation(s):

Fucon Capsule:
Each capsule contains:
Hyoscine-N-Butylbromide 10mg

Action(s):

- 1. Hyoscine-N-Butylbromide is an antimuscarinic agent with central and peripheral actions. It is a more powerful suppressant of salivation than atropine.
- 2. Hyoscine-N-Butylbromide is being poorly absorbed from the gastro-intestinal tract and does not readily pass the blood brain barrier.

Indication(s):

For the treatment of spasmodic pain, esophageal spasm, cholecystitis, choledocholithiasis, uriasis cystospasm, dysuria, pollakisuria, and spasmodic dysmenorrhea.

Dosage and Administration:

1 - 2 capsules three to four times daily.
To be taken orally

Precaution(s) / Warning(s):

Hyoscine should be used with caution in children and in geriatric patients with myasthenia gravis and with fever. Hyoscine may cause drowsiness, patients so affected should not drive or operate machinery. Patients should abstain from alcohol. Use with caution in patients with impaired metabolic, liver or kidney functions, as adverse CNS effects are more likely to occur in these patients. Because of a possible mydriatic effect, Hyoscine should not be administered to patients with glaucoma.

Pregnancy and lactation:

None known.

Drug Interactions:

- 1. Concurrent administration of Fucon with tricyclic antidepressant, monoamine oxidase inhibitors, antihistamines, amantadine and disopyramide can potentiate the antimuscarinic side effect of Fucon.
- 2. Fucon enhances tachycardic effects of β-adrenergic agents.
- 3. Fucon may cause reduced absorption of Ketoconazole.
- 4. The gastro-intestinal effects of dopamine antagonist such as Metoclopramide and Domperidone can be antagonised by Fucon.

Side Effect(s) / Adverse Reaction(s):

Side effects include dryness of the mouth with difficulty in swallowing and talking, thirst, dilation of the pupils (mydriasis), loss of accommodation (cycloplegia) and photophobia, flushing and dryness of the skin, transient bradycardia followed by tachycardia and urinary urgency, difficulty and retention, vomiting, giddiness, retrosternal pain, CNS depression, coma, circulatory and respiratory failure.

Contraindication(s):

Hyoscine is contraindicated in patients with closed-angle glaucoma, or in patients with a narrow-angle between the iris and the cornea and with prostatic enlargement, patients in whom it may lead to urinary retention, and in those with paralytic ileus or pyloric stenosis and patients with history of hypersensitivity.

Symptoms and Treatment for Overdosage, and Antidote(s):

Signs of overdose: Blurred vision, clumsiness or unsteadiness, confusion, convulsion, dryness of mouth, nose or throat, fever, unusual excitement nervousness, usually fast heartbeat. In treatment of overdosage, emesis or lavage of the stomach with 4% tannin acid solution and administration of activated charcoal is recommended: slow intravenous administration of 0.5mg to 2mg (0.5mg to 1mg in children, up to a total dose of 2mg) of physostigmine, at a rate not to exceed 1mg per minute, when needed, may be given in repeated doses of 1 to 4mg up to a total dose of 5mg in adults. Management of excitement may be accomplished by administering small doses of short-acting barbiturate such as thiopental sodium (100mg) or benzodiazepines or by rectal infusion of a 2% solution of chloral hydrate.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in a tight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description and Packing(s):

Fucon Capsule:
A #4 dark green cap pale green body hard gelatin capsule.
Plastic bottle of 1000's and 1500's (for export only).
Blister packing of 10's x 10 and 10's x 100.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)
Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679