



Package leaflet: Information for the User

COLLOMACK[®] Topical solution 10 % w/w

(Salicylic Acid)

Description

Clear as water R, colorless to light yellow, pungent smell of acetic acid.

Indication

For external application to the skin in order to remove the skin in the event of hardening caused by warts, corns and calluses.

Recommended dosage

Soak the affected area in warm water for 5 minutes. Dry thoroughly with a clean towel. The surface of the affected area should be rubbed with a nail file, pumice stone, emery board or coarse washcloth, with care taken not to cause bleeding. Apply the solution to the affected area using cotton bud, twice daily. Care should be taken to avoid the surrounding healthy skin. The affected area should cover with a plaster (dressing) to help the penetration of ingredients.

Clinically visible improvement should occur in 1-2 weeks, but the maximum effect may be expected after 4-8 weeks. If wart persists beyond 12 weeks of treatment, the patient should be advised to consult their pharmacist or doctor.

Not recommended for use in infants under the age of 2 years. Children under 12 years should be treated under supervision.

Route of Administration

Topical application in the form of the solution.

Contraindication

Hypersensitivity to salicylic acid or to any of the excipients.

Collomack should not be used by diabetics or patients with impaired blood circulation. Do not use if the wart or surrounding skin is inflamed or broken. Do not use on moles, birthmarks, unusual warts with hair growth, on facial warts, or in the anal or perineal region.

Special warning and Precaution for use

Collomack is for external use only. Do not permit contact with eyes or mucous membranes. If contact occurs flush with water for 15 minutes. Do not allow contact with normal skin around wart. Avoid using on areas of broken or damaged skin. Discontinue treatment if excessive irritation occurs. Excessive prolonged use of topical salicylic acid may result in symptoms of salicylism and must therefore be avoided.

Interaction with other medicaments

There are no known interactions when used as indicated. However, topical salicylic acid may increase the absorption of other topically applied medicines. Concomitant use of Collomack and other topical medicines on the treated wart should therefore be avoided.

Fertility, Pregnancy and lactation

Pregnancy

Collomack should not be used during pregnancy except on an area smaller than 5 cm².

Lactation

Collomack must not be used on the breast of breastfeeding mothers. If the doctor considers a regular treatment of large areas of skin with Collomack necessary, it is advisable to abort at an early stage as risks cannot be ruled out due to a lack of detoxification in newborns.

Effects on ability to drive and use machines

There are no special precautions required.

Side effects / Adverse Reaction

As with all pharmaceutical products Collomack can have side effects though they do not occur in every case.

Assessment of the frequency of side effects is based on the following data:

Very common :	More than 1 in 10 cases
Common :	Fewer than 1 in 10 but more than 1 in 100 cases
Occasional :	Fewer than 1 in 100 but more than 1 in 1,000 cases
Rare :	Fewer than 1 in 1,000 but more than 1 in 10,000 cases
Very rare :	Fewer than 1 in 10,000 cases including individual instances

Significant side effects or symptoms you should look out for and what to do if you are Affected.

If you are affected by one of the side effects mentioned below, discontinue using Collomack and consult your doctor as soon as possible.

Occasionally application causes local irritation which is usually a slight burning sensation but in Individual cases can produce a touch-sensitive allergic reaction.

Please inform your doctor or pharmacist if one of the specified side effects affects you significantly or if you notice side effects that are not indicated in this package leaflet.





Overdose and treatment

Symptoms of systemic salicylate poisoning have been reported after the application of salicylic acid to large areas of skin and for prolonged periods. Salicylism may also occur in the unlikely event of large quantities being ingested. Salicylism is unlikely to occur if collomack I is used as indicated.

Pharmacodynamics

Pharmacotherapeutic group: Wart and anticon preparations

ATC code: D11AF

Salicylic acid has bacteriostatic and fungicidal actions, but its keratolytic properties which are important for this medicinal product. When applied externally it produces slow and painless destruction of the epithelium. Salicylic acid is usually applied in the form of paint in a collodian base (10 to 17 %) or as a plaster (20 to 50 %) to destroy warts or corns.

Pharmacokinetic properties

The metabolism of salicylic acid is achieved by conjugation with glycine salicyluric acid, with glucuronic acid at the phenolic OH-group to etherglucuronide and at the COOH-group to form ester glucuronide or by hydroxylation to give gentisic acid or dihydroxybenzoic acid. The half-life of salicylic acid is between 2 and 3 hours in the normal dose range and may increase to 15-30 hours at high doses due to limited liver capacity to conjugate salicylic acid.

Preclinical safety data

a) Acute toxicity

Salicylic acid penetrates rapidly as a result of animal experiments and human pharmacokinetic studies, depending on the basis and penetration-influencing factors such as the skin condition. That's why the rare occurrence of intoxications in topical application is dependent on the pharmaceutical dosage form, the amount of salicylic acid applied, the duration of treatment, the frequency of treatment and the dermatological pathology. Only serum values above 30 mg/dl are associated with intoxication symptoms.

b) Chronic toxicity

Animal experiments with dermal administration over long-term effects are not available (see acute toxicity).

c) Mutagenic and tumor-producing potential

There is no relevant reference to the mutagenic effect of salicylic acid in the extensive scientific literature.
There is no tumor-producing potential of salicylic acid.

d) Reproductive toxicity

Salicylates have been found in animal experiments on several animal species at systemic application of teratogenic effects. Implantation disorders, embryo- and fetotoxic effects as well as disturbances of the learning ability in the off-spring after prenatal exposure have been described. Toxic effects are not to be expected when used as intended, since toxicologically relevant plasma levels are not reached in conventional applications.

Storage

Do not store above 30 °C.

The bottle should always be kept tightly closed. The solution is slightly volatile and flammable.

Pharmaceutical particulars

List of excipients

99 % acetic acid, docusate sodium, castor oil, pyroxyline, ethanol 96 %, acetone, isobutyl acetate 6.3

Shelf life

3 years.

Manufactured by:

P.W. Beyvers GmbH
Schaffhausener Str. 26 -34
Berlin 12099 GERMANY

Product Registration Holder:

Singapore

RX Pharma Pte Ltd
Email: rxpharma@singnet.com.sg

Malaysia

DCH AURIGA (M) SDN BHD
Lot 6, Persiaran Perusahaan,
Seksyen 23, 40300 ,
Shah Alam, Selangor,
Malaysia.

Date of Revision

January 2023

