



AmsolvanTM

Oral Solution 30mg/5ml

COMPOSITION

Each 5ml contains Ambroxol Hydrochloride 30mg.
Benzoic Acid 0.15%w/v as preservative.

PHARMACODYNAMICS

Ambroxol hydrochloride is a mucolytic that alters the structure of mucus to decrease its viscosity thereby facilitating its removal by ciliary action or expectoration.

PHARMACOKINETICS

Absorption

Absorption of ambroxol hydrochloride is rapid and complete, with dose linearity in the therapeutic range. Maximum plasma levels are reached within 1-2.5 hours following oral administration.

Distribution

In the therapeutic range, plasma protein-binding was found to be approximately 90%. Distribution of ambroxol hydrochloride from blood to tissue is rapid and pronounced, with the highest concentration of the active substance found in the lungs. The volume of distribution following oral administration was estimated to be 552 L.

Metabolism

About 30% of an orally administered dose is eliminated via first-pass metabolism. Studies in human liver microsomes have shown that CYP3A4 is responsible for the metabolism of ambroxol hydrochloride to dibromanthranilic acid. Ambroxol hydrochloride is primarily metabolized in the liver by glucuronidation and some cleavage to dibromanthranilic acid (approximately 10% of dose) aside from some minor metabolites.

Elimination

Ambroxol hydrochloride is eliminated with a terminal elimination half-life of approximately 10 hours. Total clearance is in the range of 660 ml/min, with renal clearance accounting for approximately 83% of the total clearance.

Special Populations: In patients with hepatic dysfunction, elimination of ambroxol hydrochloride is reduced, resulting in approximately 1.3 to 2-fold higher plasma levels. Due to the high therapeutic range of ambroxol hydrochloride, dose adjustments are not necessary.

Others: Age and gender were not found to affect the pharmacokinetics of ambroxol hydrochloride to a clinically relevant extent and thus, there is no necessity for adjustment of dosage regimens. Food was not found to influence the bioavailability of ambroxol hydrochloride.

INDICATIONS

Amsolvan clears the airway by thinning the thick mucus in the respiratory tract and facilitates expectoration.

DOSAGE AND ADMINISTRATION

For oral use only. Can be taken with or without food.

Amsolvan is not recommended for children under 2 years of age without medical advice.

Adults and children over 12 years old : 10ml 2 times daily

Children 6-12 years old : 5ml 2-3 times daily

Children 2-5 years old : 2.5ml 3 times daily

CONTRAINDICATIONS

Contraindicated in patients who are allergic to ambroxol hydrochloride.

Amsolvan contains sorbitol and patients who have rare hereditary fructose intolerance should not take this medicine.

WARNINGS AND PRECAUTIONS

There have been very few reports of severe skin lesions eg, Stevens-Johnson syndrome and toxic epidermal necrolysis (TEN) in temporal association with the administration of expectorants such as ambroxol hydrochloride.

Mostly these could be explained by the severity of the patient's underlying disease and/or concomitant medication.

During the early phase of a Stevens-Johnson Syndrome or TEN, a patient may first experience non-specific influenza-like prodromes. It is possible that a symptomatic treatment is started with a cough and cold medication. Therefore, if new skin or mucosal lesions occur, medical advice should be sought immediately and treatment with ambroxol discontinued as a precaution.

In the presence of impaired renal function, ambroxol hydrochloride may be used only after consulting a physician.

There is no evidence for an effect on the ability to drive and use machines. Studies on the effects on the ability to drive and use machines have not been performed.

DRUG INTERACTIONS

No clinically relevant interaction with other medications have been reported.

PREGNANCY AND LACTATION

Ambroxol hydrochloride crosses the placental barrier. The use of ambroxol hydrochloride is not recommended during first trimester of pregnancy as usual precautions in the use of drugs during pregnancy. Animal studies do not indicate direct or indirect harmful effects with respect to pregnancy, foetal development, parturition or postnatal development.

Extensive clinical experience after the 28th week of pregnancy has shown no evidence of harmful effects on the foetus.

Ambroxol hydrochloride is excreted in breast milk. It is not recommended for use in nursing mothers though unfavourable effects on breastfed infants would not be expected.

SIDE EFFECTS

- *Gastrointestinal:* Dyspepsia, nausea, vomiting, diarrhoea and abdominal pain.
- *Immune system:* Anaphylactic reactions including anaphylactic shock.
- *Skin and Subcutaneous Tissue:* Angioedema, rash, urticaria, pruritus and other hypersensitivity.
- *Respiratory, Mediastinal and Thoracic:* Oral and pharyngeal hypoaesthesia, dry mouth and dry throat.
- *Nervous System:* Dysgeusia

SYMPTOMS AND TREATMENT FOR OVERDOSAGE

No specific overdose symptoms have been reported in man to date. Based on accidental overdose and/or medication error reports, the observed symptoms are consistent with the known side effects of ambroxol hydrochloride at recommended doses and may need symptomatic treatment.

STORAGE CONDITION

Store below 30°C.

SHELF LIFE

The expiry date is indicated on the packaging.

PRODUCT DESCRIPTION, DOSAGE FORM AND PACKAGING AVAILABLE

Clear, faintly yellow solution with strawberry flavor.

Available in pack size of 100ml bottles.

KEEP OUT OF REACH OF CHILDREN/ JAUHI DARI KANAK-KANAK

For further information, please consult your physician or pharmacist.

Revision Date: 01-Sep-2016.

Manufacturer and Product Registration Holder

Xepa-Soul Pattinson (Malaysia) Sdn Bhd

1-5 Cheng Industrial Estate 75250 Melaka, Malaysia.

