

8986



Cymbalox 30mg Delayed-Release Capsules
Cymbalox 60mg Delayed-Release Capsules
(Duloxetine Hydrochloride Delayed-Release capsules)

1. NAME OF THE MEDICINAL PRODUCT

Cymbalox 30 mg Delayed-Release Capsules
Cymbalox 60 mg Delayed-Release Capsules

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each Hard Gelatin capsule contains:
Duloxetine Hydrochloride Ph. Eur Eq. to Duloxetine 30 mg
Duloxetine Hydrochloride Ph. Eur Eq. to Duloxetine 60 mg
For the full list of excipients, see section 6.1

3. PHARMACEUTICAL FORM

Cymbalox 30 mg Delayed-Release Capsules

Opaque blue/opaque white, hard gelatin capsule size "3", having imprinting "157" on body with green ink and "A" on cap with green ink and filled with white to off white coloured pellets.

Cymbalox 60 mg Delayed-Release Capsules

Opaque blue/opaque green, hard gelatin capsule size "1", having imprinting "158" on body with white ink and "A" on cap with white ink and filled with white to off white coloured pellets.

4. Clinical particulars

4.1 Therapeutic indications

Treatment of major depressive disorder.
Management of neuropathic pain associated with diabetic peripheral neuropathy in adults.
Treatment of generalised anxiety disorder.

4.2 Posology and method of administration

Posology

Major depressive disorder

The starting and recommended maintenance dose is 60 mg once daily with or without food. Doses above 60 mg once daily, up to a maximum dose of 120 mg per day have been evaluated from a safety perspective in clinical trials. However, there is no clinical evidence suggesting that patients not responding to the initial recommended dose may benefit from dose up-titrations.

Therapeutic response is usually seen after 2-4 weeks of treatment.

After consolidation of the antidepressive response, it is recommended to continue treatment for several months, in order to avoid relapse. In patients responding to duloxetine, and with a history of repeated episodes of major depression, further long-term treatment at a dose of 60 to 120 mg/day could be considered.

Generalised anxiety disorder

The recommended starting dose in patients with generalised anxiety disorder is 30 mg once daily with or without food. In patients with insufficient response the dose should be increased to 60 mg, which is the usual maintenance dose in most patients.

In patients with co-morbid major depressive disorder, the starting and maintenance dose is 60 mg once daily (please see also dosing recommendation above). Doses up to 120 mg per day have been shown to be efficacious and have been evaluated from a safety perspective in clinical trials. In patients with insufficient response to 60 mg, escalation up to 90 mg or 120 mg may therefore be considered. Dose escalation should be based upon clinical response and tolerability.

After consolidation of the response, it is recommended to continue treatment for several months, in order to avoid relapse.

Diabetic peripheral neuropathic pain

The starting and recommended maintenance dose is 60 mg daily with or without food. Doses above 60 mg once daily, up to a maximum dose of 120 mg per day administered in evenly divided doses, have been evaluated from a safety perspective in clinical trials. The plasma concentration of duloxetine displays large inter-individual variability (see section 5.2). Hence, some patients that respond insufficiently to 60 mg may benefit from a higher dose.

Response to treatment should be evaluated after 2 months. In patients with inadequate initial response, additional response after this time is unlikely.

The therapeutic benefit should be reassessed regularly (at least every three months) (see section 5.1).

Special populations

Elderly

No dose adjustment is recommended for elderly patients solely on the basis of age. However, as with any medicine, caution should be exercised when treating the elderly, especially with duloxetine 120 mg per day for major depressive disorder or generalised anxiety disorder, for which data are limited (see sections 4.4 and 5.2).

Hepatic impairment

Duloxetine must not be used in patients with liver disease resulting in hepatic impairment (see sections 4.3 and 5.2).

Renal impairment

No dose adjustment is necessary for patients with mild or moderate renal dysfunction (creatinine clearance 30 to 80 ml/min). Duloxetine must not be used in patients with severe renal impairment (creatinine clearance < 30 ml/min; see section 4.3).

Paediatric population

Duloxetine should not be used in children and adolescents under the age of 18 years for the treatment of major depressive disorder because of safety and efficacy concerns (see sections 4.4, 4.8 and 5.1). The safety and efficacy of duloxetine for the treatment of generalised anxiety disorder in paediatric patients aged 7-17 years have not been established.

The safety and efficacy of duloxetine for the treatment of diabetic peripheral neuropathic pain has not been studied. No data are available.

Discontinuation of treatment

Abrupt discontinuation should be avoided. When stopping treatment with Duloxetine the dose should be gradually reduced over a period of at least one to two weeks in order to reduce the risk of withdrawal reactions (see sections 4.4 and 4.8). If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose, but at a more gradual rate.

Method of administration

For oral use.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Concomitant use of Duloxetine with non-selective, irreversible monoamine oxidase inhibitors (MAOIs) is contraindicated (see section 4.5). Liver disease resulting in hepatic impairment (see section 5.2)

Duloxetine should not be used in combination with fluvoxamine, ciprofloxacin or enoxacin (i.e. potent CYP1A2 inhibitors) since the combination results in elevated plasma concentrations of duloxetine (see section 4.5). Severe renal impairment (creatinine clearance < 30 ml/min) (see section 4.4)

The initiation of treatment with duloxetine is contraindicated in patients with uncontrolled hypertension that could expose patients to a potential risk of hypertensive crisis (see sections 4.4 and 4.8).

4.4 Special warnings and precautions for use

Mania and seizures

Duloxetine should be used with caution in patients with a history of mania or a diagnosis of bipolar disorder, and/or seizures.

Mydriasis

Mydriasis has been reported in association with duloxetine, therefore, caution should be used when prescribing duloxetine to patients with increased intraocular pressure, or those at risk of acute narrow angle glaucoma.

Blood pressure and heart rate

Duloxetine has been associated with an increase in blood pressure and clinically significant hypertension in some patients. This may be due to the noradrenergic effect of duloxetine. Cases of hypertensive crisis have been reported with duloxetine, especially in patients with pre-existing hypertension. Therefore, in patients with known hypertension and/or other cardiac disease, blood pressure monitoring is recommended, especially during the first month of treatment. Duloxetine should be used with caution in patients whose conditions could be compromised by an increased heart rate or by an increase in blood pressure. Caution should also be exercised when duloxetine is used with medicinal products that may impair its metabolism (see section 4.5). For patients who experience a sustained increase in blood pressure while receiving duloxetine either dose reduction or gradual discontinuation should be considered (see section 4.8). In patients with uncontrolled hypertension duloxetine should not be initiated (see section 4.3).

Renal impairment

Increased plasma concentrations of duloxetine occur in patients with severe renal impairment on haemodialysis (creatinine clearance < 30 ml/min). For patients with severe renal impairment, see section 4.3. See section 4.2 for information on patients with mild or moderate renal dysfunction.

Serotonin syndrome

As with other serotonergic agents, serotonin syndrome, a potentially life-threatening condition, may occur with duloxetine treatment, particularly with concomitant use of other serotonergic agents (including SSRIs, SNRIs tricyclic antidepressants or triptans), with agents that impair metabolism of serotonin such as MAOIs, or with antipsychotics or other dopamine antagonists that may affect the serotonergic neurotransmitter systems (see sections 4.3 and 4.5).

Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g. hyperreflexia, incoordination) and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhoea).

If concomitant treatment with duloxetine and other serotonergic agents that may affect the serotonergic and/or dopaminergic neurotransmitter systems is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases.

St John's wort

Adverse reactions may be more common during concomitant use of duloxetine and herbal preparations containing St John's wort (*Hypericum perforatum*).

Suicide

Major Depressive Disorder and Generalised Anxiety Disorder

Depression is associated with an increased risk of suicidal thoughts, self-harm and suicide (suicide-related events). This risk persists until significant remission occurs. As improvement may not occur during the first few weeks or more of treatment, patients should be closely monitored until such improvement occurs. It is general clinical experience that the risk of suicide may increase in the early stages of recovery.

Other psychiatric conditions for which duloxetine is prescribed can also be associated with an increased risk of suicide-related events. In addition, these conditions may be co-morbid with major depressive disorder. The same precautions observed when treating patients with major depressive disorder should therefore be observed when treating patients with other psychiatric disorders.

Patients with a history of suicide-related events or those exhibiting a significant degree of suicidal thoughts prior to commencement of treatment are known to be at greater risk of suicidal thoughts or suicidal behaviour, and should receive careful monitoring during treatment. A meta-analysis of placebo-controlled clinical trials of antidepressant medicinal products in psychiatric disorders showed an increased risk of suicidal behaviour with antidepressants compared to placebo in patients less than 25 years old. Cases of suicidal thoughts and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation (see section 4.8).

Close supervision of patients and in particular those at high risk should accompany medicinal product therapy especially in early treatment and following dose changes. Patients (and caregivers of patients) should be alerted about the need to monitor for any clinical worsening, suicidal behaviour or thoughts and unusual changes in behaviour and to seek medical advice immediately if these symptoms present.

Diabetic Peripheral Neuropathic Pain

As with other medicinal products with similar pharmacological action (antidepressants), isolated cases of suicidal ideation and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation. Physicians should encourage patients to report any distressing thoughts or feelings at any time.

Use in children and adolescents under 18 years of age

Duloxetine should not be used in the treatment of children and adolescents under the age of 18 years. Suicide-related behaviours (suicide attempts and suicidal thoughts), and hostility (predominantly aggression, oppositional behaviour and anger), were more frequently observed among children and adolescents treated with antidepressants. If, based on clinical need, a decision to treat is nevertheless taken, the patient should be carefully monitored for the appearance of suicidal symptoms (see section 5.1). In addition, long-term safety data in children and adolescents concerning growth, maturation and cognitive and behavioural development are lacking (see section 4.8).

WARNING

Suicidality in Children and Adolescents

- Antidepressants increase the risk of suicidal thinking and behavior (suicidality) in children and adolescents with major depressive disorder (MDD) and other psychiatric disorders.
- Anyone considering the use of an antidepressant in a child or adolescent for any clinical use must balance the risk of increased suicidality with the clinical need.
- Patients who are started on therapy should be observed closely for clinical worsening, suicidality, or unusual changes in behavior.
- Families and caregivers should be advised to closely observe the patient and to communicate with the prescriber.
- The indication(s) approved in paediatric for the particular drug should be clearly stated / included.

Haemorrhage

There have been reports of bleeding abnormalities, such as ecchymoses, purpura and gastrointestinal haemorrhage with selective serotonin reuptake inhibitors (SSRIs) and serotonin/noradrenaline reuptake inhibitors (SNRIs), including duloxetine. Caution is advised in patients taking anticoagulants and/or medicinal products known to affect platelet function (e.g. NSAIDs or acetylsalicylic acid (ASA)), and in patients with known bleeding tendencies.

Hyponatraemia

Hyponatraemia has been reported when administering duloxetine, including cases with serum sodium lower than 110 mmol/l. Hyponatraemia may be due to a syndrome of inappropriate anti-diuretic hormone secretion (SIADH). The majority of cases of hyponatraemia were reported in the elderly, especially when coupled with a recent history of, or condition pre-disposing to, altered fluid balance. Caution is required in patients at increased risk for hyponatraemia, such as elderly, cirrhotic, or dehydrated patients or patients treated with diuretics.

Discontinuation of treatment

Withdrawal symptoms when treatment is discontinued are common, particularly if discontinuation is abrupt (see section 4.8). The risk of withdrawal symptoms seen with SSRIs and SNRIs may be dependent on several factors including the duration and dose of therapy and the rate of dose reduction. The most commonly reported reactions are listed in section 4.8. Generally, these symptoms are mild to moderate, however, in some patients they may be severe in intensity. They usually occur within the first few days of discontinuing treatment, but there have been very rare reports of such symptoms in patients who have inadvertently missed a dose. Generally, these symptoms are self-limiting and usually resolve within 2 weeks, though in some individuals they may be prolonged (2-3 months or more). It is therefore advised that duloxetine should be gradually tapered when discontinuing treatment over a period of no less than 2 weeks, according to the patient's needs (see section 4.2).

Elderly

Data on the use of duloxetine 120 mg in elderly patients with major depressive disorder and generalised anxiety disorder are limited. Therefore, caution should be exercised when treating the elderly with the maximum dosage (see sections 4.2 and 5.2).

Akathisia/psychomotor restlessness

The use of duloxetine has been associated with the development of akathisia, characterised by a subjectively unpleasant or distressing restlessness and need to move often accompanied by an inability to sit or stand still. This is most likely to occur within the first few weeks of treatment. In patients who develop these symptoms, increasing the dose may be detrimental.

Medicinal products containing duloxetine

Duloxetine is used under different trademarks in several indications (treatment of diabetic neuropathic pain, major depressive disorder, generalised anxiety disorder and stress urinary incontinence). The use of more than one of these products concomitantly should be avoided.

Hepatitis/increased liver enzymes

Cases of liver injury, including severe elevations of liver enzymes (> 10 times upper limit of normal), hepatitis and jaundice have been reported with duloxetine (see section 4.8). Most of them occurred during the first months of treatment. The pattern of liver damage was predominantly hepatocellular. Duloxetine should be used with caution in patients treated with other medicinal products associated with hepatic injury.

Sexual dysfunction

Selective serotonin reuptake inhibitors (SSRIs)/serotonin norepinephrine reuptake inhibitors (SNRIs) may cause symptoms of sexual dysfunction (see section 4.8). There have been reports of long-lasting sexual dysfunction where the symptoms have continued despite discontinuation of SSRIs/SNRIs.

Sucrose

This product contains sucrose. Patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrose-isomaltase insufficiency should not take this medicine.

4.5 Interaction with other medicinal products and other forms of interaction

Monoamine oxidase inhibitors (MAOIs)

Due to the risk of serotonin syndrome, duloxetine should not be used in combination with non-selective irreversible monoamine oxidase inhibitors (MAOIs), or within at least 14 days of discontinuing treatment with an MAOI. Based on the half-life of duloxetine, at least 5 days should be allowed after stopping duloxetine before starting an MAOI (see section 4.3).

The concomitant use of duloxetine with selective, reversible MAOIs, like moclobemide, is not recommended (see section 4.4). The antibiotic linezolid is a reversible non-selective MAOI and should not be given to patients treated with duloxetine (see section 4.4).

Inhibitors of CYP1A2

Because CYP1A2 is involved in duloxetine metabolism, concomitant use of duloxetine with potent inhibitors of CYP1A2 is likely to result in higher concentrations of duloxetine. Fluvoxamine (100 mg once daily), a potent inhibitor of CYP1A2, decreased the apparent plasma clearance of duloxetine by about 77% and increased AUC_{0-t} 6-fold. Therefore duloxetine should not be administered in combination with potent inhibitors of CYP1A2 like fluvoxamine (see section 4.3).

CNS medicinal products

The risk of using duloxetine in combination with other CNS-active medicinal products has not been systematically evaluated. Consequently, caution is advised when duloxetine is taken in combination with other centrally acting medicinal products or substances, including alcohol and sedative medicinal products (e.g. benzodiazepines, morphinomimetics, antipsychotics, phenobarbital, sedative antihistamines).

Serotonergic agents

In rare cases, serotonin syndrome has been reported in patients using SSRIs/SNRIs concomitantly with serotonergic agents. Caution is advisable if duloxetine is used concomitantly with serotonergic agents like SSRIs, SNRIs, tricyclic antidepressants like clomipramine or amitriptyline, MAOIs like moclobemide or linezolid, St John's wort (*Hypericum perforatum*) or triptans, tramadol, pethidine and tryptophan (see section 4.4).

Effect of duloxetine on other medicinal products

Medicinal products metabolised by CYP1A2

The pharmacokinetics of theophylline, a CYP1A2 substrate, were not significantly affected by co-administration with duloxetine (60 mg twice daily).

Medicinal products metabolised by CYP2D6

Duloxetine is a moderate inhibitor of CYP2D6. Caution is advised if duloxetine is co-administered with medicinal products that are predominantly metabolised by CYP2D6 (risperidone, tricyclic antidepressants (TCAs) such as nortriptyline, amitriptyline, and imipramine) particularly if they have a narrow therapeutic index (such as flecainide, propafenone and metoprolol).

Oral contraceptives and other steroidal agents

Results of in vitro studies demonstrate that duloxetine does not induce the catalytic activity of CYP3A. Specific in vivo drug interaction studies have not been performed.

Anticoagulants and antiplatelet agents

Caution should be exercised when duloxetine is combined with oral anticoagulants or antiplatelet agents due to a potential increased risk of bleeding attributable to a pharmacodynamic interaction. Furthermore, increases in INR values have been reported when duloxetine was co-administered to patients treated with warfarin. However, concomitant administration of duloxetine with warfarin under steady state conditions, did not result in a clinically significant change in INR from baseline or in the pharmacokinetics of R- or S-warfarin.

Effects of other medicinal products on duloxetine

Antacids and H2 antagonists

Co-administration of duloxetine with aluminium- and magnesium-containing antacids or duloxetine with famotidine had no significant effect on the rate or extent of duloxetine absorption after administration of a 40 mg oral dose.

Inducers of CYP1A2

Population pharmacokinetic analyses have shown that smokers have almost 50% lower plasma concentrations of duloxetine compared with non-smokers.

4.6 Fertility, pregnancy and lactation

Fertility

Duloxetine had no effect on male fertility, and effects in females were only evident at doses that caused maternal toxicity.

9869



8986



520 mm

150 mm
Front Side

Alembic Pharmaceuticals Limited 	
Product: Cymbalox Capsules	Colour 
Pack: Pack Insert	
Size: 150 x 520 mm (L x H)	
Date: 27-10-2025	
	DieLine Not to be Printed 

Note: Actual pharma code number is for reference & should not be printed.

Pregnancy

There are no adequate data on the use of duloxetine in pregnant women.
The potential risk for humans is unknown.

Epidemiological data have suggested that the use of SSRIs in pregnancy, particularly in late pregnancy, may increase the risk of persistent pulmonary hypertension in the newborn (PPHN). Although no studies have investigated the association of PPHN to SNRI treatment, this potential risk cannot be ruled out with duloxetine taking into account the related mechanism of action (inhibition of the re-uptake of serotonin).

As with other serotonergic medicinal products, discontinuation symptoms may occur in the neonate after maternal duloxetine use near term. Discontinuation symptoms seen with duloxetine may include hypotonia, tremor, jitteriness, feeding difficulty, respiratory distress and seizures. The majority of cases have occurred either at birth or within a few days of birth.

Duloxetine should be used in pregnancy only if the potential benefit justifies the potential risk to the foetus. Women should be advised to notify their physician if they become pregnant, or intend to become pregnant, during therapy.

Pregnancy and lactation

Observational data indicate an increased risk (less than 2-fold) of postpartum haemorrhage following SSRI/SNRI exposure within the month prior to birth.

Breast feeding

The safety of duloxetine in infants is not known, the use of duloxetine while breast-feeding is not recommended.

4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. Duloxetine may be associated with sedation and dizziness. Patients should be instructed that if they experience sedation or dizziness they should avoid potentially hazardous tasks such as driving or operating machinery.

4.8 Undesirable effects

a. Summary of the safety profile

The most commonly reported adverse reactions in patients treated with duloxetine were nausea, headache, dry mouth, somnolence, and dizziness. However, the majority of common adverse reactions were mild to moderate, they usually started early in therapy, and most tended to subside even as therapy was continued.

b. Tabulated summary of adverse reactions

Table 1: Adverse reactions

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Very common	Common	Uncommon	Rare	Very Rare
Infections and infestations				
		Laryngitis		
Immune system disorders				
			Anaphylactic reaction, Hyper-sensitivity disorder	
Endocrine disorders				
			Hypo-thyroidism	
Metabolism and nutrition disorders				
	Decreased Appetite	Hyperglycaemia (reported especially in diabetic patients)	Dehydration, Hyponatraemia, SIADH ¹	
Psychiatric disorders				
	Insomnia, Agitation, Libido decreased, Anxiety, Orgasm abnormal, Abnormal dreams	Suicidal ideation, Sleep disorder, Briuxism, Disorientation, Apathy	Suicidal behaviour, Mania, Hallucinations, Aggression and anger ¹	
Nervous system disorders				
Headache Somnolence	Dizziness, Lethargy, Tremor, Paraesthesia	Myoclonus, Akathisia, Nervousness, Disturbance in attention, Dysgeusia, Dyskinesia, Restless legs syndrome, Poor quality sleep	Serotonin syndrome, Convulsion ¹ , Psychomotor restlessness, Extra-pyramidal symptoms ⁴	
Eye disorders				
	Blurred vision	Mydriasis, Visual impairment	Glaucoma	
Ear and labyrinth disorders				
	Tinnitus ¹	Vertigo, Ear pain		
Cardiac disorders				
	Palpitations	Tachycardia, Supra-ventricular arrhythmia, mainly atrial fibrillation		
Vascular disorders				
	Blood pressure increase, Flushing	Syncope, Hypertension, Orthostatic hypotension, Peripheral coldness	Hypertensive crisis ^{3,4}	
Respiratory, thoracic and mediastinal disorders				
	Yawning	Throat tightness, Epistaxis		
Gastrointestinal disorders				
Nausea Dry mouth	Constipation, Diarrhoea, Abdominal pain, Vomiting, Dyspepsia, Flatulence	Gastrointestinal haemorrhage ⁷ , Gastroenteritis, Eructation, Gastritis, Dysphagia	Stomatitis, Haematochezia, Breath odour, Microscopic colitis ⁷	
Hepato-biliary disorders				
		Hepatitis, Elevated liver enzymes (ALT, AST, alkaline phosphatase), Acute liver injury	Hepatic failure, Jaundice ⁸	
Skin and subcutaneous tissue disorders				
	Sweating increased, Rash	Night sweats, Urticaria, Dermatitis contact, Cold sweat, Photo-sensitivity reactions, Increased tendency to bruise	Stevens-Johnson Syndrome ⁴ , Angio-neurotic oedema	Cutaneous vasculitis
Musculoskeletal and connective tissue disorders				
	Musculoskeletal pain, Muscle spasm	Muscle tightness, Muscle twitching	Trismus	
Renal and urinary disorders				
	Dysuria, Pollakiuria	Urinary retention, Urinary hesitation, Nocturia, Polyuria, Urine flow decreased	Urine odour abnormal	
Reproductive system and breast disorders				
	Erectile dysfunction, Ejaculation disorder, Ejaculation delayed	Gynaecological haemorrhage, Menstrual disorder, Sexual dysfunction, Testicular pain	Menopausal symptoms Galactorrhoea, Hyperprolactinaemia	
General disorders and administration site conditions				
	Falls, Fatigue	Chest pain ¹ , Feeling abnormal, Feeling cold, Thirst, Chills, Malaise, Feeling hot, Gait disturbance		
Investigations				
	Weight decrease	Weight increase, Blood creatine phosphokinase increased, Blood potassium increased	Blood cholesterol increased	
¹ Cases of convulsion and cases of tinnitus have also been reported after treatment discontinuation. ² Cases of orthostatic hypotension and syncope have been reported especially at the initiation of treatment. ³ See section 4.4. ⁴ Cases of aggression and anger have been reported particularly early in treatment or after treatment discontinuation. ⁵ Cases of suicidal ideation and suicidal behaviours have been reported during duloxetine therapy or early after treatment discontinuation (see section 4.4). ⁶ Estimated frequency of post-marketing surveillance reported adverse reactions; not observed in placebo-controlled clinical trials. ⁷ Not statistically significantly different from placebo. ⁸ Falls were more common in the elderly (≥ 65 years old) ⁹ Estimated frequency based on all clinical trial data				

c. Description of selected adverse reactions

Discontinuation of duloxetine (particularly when abrupt) commonly leads to withdrawal symptoms. Dizziness, sensory disturbances (including paraesthesia or electric shock-like sensations, particularly in the head), sleep disturbances (including insomnia and intense dreams), fatigue, somnolence, agitation or anxiety, nausea and/or vomiting, tremor, headache, myalgia, irritability, diarrhoea, hyperhidrosis and vertigo are the most commonly reported reactions.

Generally, for SSRIs and SNRIs, these events are mild to moderate and self-limiting, however, in some patients they may be severe and/or prolonged. It is therefore advised that when duloxetine treatment is no longer required, gradual discontinuation by dose tapering should be carried out (see sections 4.2 and 4.4).

4.9 Overdose

Cases of overdose, alone or in combination with other medicinal products, with duloxetine doses of 5400 mg were reported. Some fatalities have occurred, primarily with mixed overdoses, but also with duloxetine alone at a dose of approximately 1000 mg. Signs and symptoms of overdose (duloxetine alone or in combination with other medicinal products) included somnolence, coma, serotonin syndrome, seizures, vomiting and tachycardia.

No specific antidote is known for duloxetine but if serotonin syndrome ensues, specific treatment (such as with cyproheptadine and/or temperature control) may be considered. A free airway should be established. Monitoring of cardiac and vital signs is recommended, along with appropriate symptomatic and supportive measures. Gastric lavage may be indicated if performed soon after ingestion or in symptomatic patients. Activated charcoal may be useful in limiting absorption. Duloxetine has a large volume of distribution and forced diuresis, haemoperfusion, and exchange perfusion are unlikely to be beneficial.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Other antidepressants, ATC code: N06AX21

Mechanism of action

Duloxetine is a combined serotonin (5-HT) and noradrenaline (NA) reuptake inhibitor. It weakly inhibits dopamine reuptake with no significant affinity for histaminergic, dopaminergic, cholinergic and adrenergic receptors. Duloxetine dose-dependently increases extracellular levels of serotonin and noradrenaline in various brain areas of animals.

Pharmacodynamic effects

Duloxetine normalised pain thresholds in several preclinical models of neuropathic and inflammatory pain and attenuated pain behaviour in a model of persistent pain. The pain inhibitory action of duloxetine is believed to be a result of potentiation of descending inhibitory pain pathways within the central nervous system.

5.2 Pharmacokinetic properties

Duloxetine is administered as a single enantiomer. Duloxetine is extensively metabolised by oxidative enzymes (CYP1A2 and the polymorphic CYP2D6), followed by conjugation. The pharmacokinetics of duloxetine demonstrate large intersubject variability (generally 50-60%), partly due to gender, age, smoking status and CYP2D6 metaboliser status.

Absorption

Duloxetine is well absorbed after oral administration with a Cmax occurring 6 hours post dose. The absolute oral bioavailability of duloxetine ranged from 32% to 80% (mean of 50%). Food delays the time to reach the peak concentration from 6 to 10 hours and it marginally decreases the extent of absorption (approximately 11%). These changes do not have any clinical significance.

Distribution

Duloxetine is approximately 96% bound to human plasma proteins. Duloxetine binds to both albumin and alpha-1 acid glycoprotein. Protein binding is not affected by renal or hepatic impairment.

Biotransformation

Duloxetine is extensively metabolised and the metabolites are excreted principally in urine. Both cytochromes P450-2D6 and 1A2 catalyse the formation of the two major metabolites glucuronide conjugate of 4-hydroxy duloxetine and sulphate conjugate of 5-hydroxy 6-methoxy duloxetine. Based upon in vitro studies, the circulating metabolites of duloxetine are considered pharmacologically inactive. The pharmacokinetics of duloxetine in patients who are poor metabolisers with respect to CYP2D6 has not been specifically investigated. Limited data suggest that the plasma levels of duloxetine are higher in these patients.

Elimination

The elimination half-life of duloxetine ranges from 8 to 17 hours (mean of 12 hours). After an intravenous dose the plasma clearance of duloxetine ranges from 22 l/hr to 46 l/hr (mean of 36 l/hr). After an oral dose the apparent plasma clearance of duloxetine ranges from 33 to 261 l/hr (mean 101 l/hr).

Special populations

Gender

Pharmacokinetic differences have been identified between males and females (apparent plasma clearance is approximately 50% lower in females). Based upon the overlap in the range of clearance, gender based pharmacokinetic differences do not justify the recommendation for using a lower dose for female patients.

Age

Pharmacokinetic differences have been identified between younger and elderly females (≥ 65 years) (AUC increases by about 25% and half-life is about 25% longer in the elderly), although the magnitude of these changes is not sufficient to justify adjustments to the dose. As a general recommendation, caution should be exercised when treating the elderly (see sections 4.2 and 4.4).

Renal impairment

End stage renal disease (ESRD) patients receiving dialysis had 2-fold higher duloxetine Cmax and AUC values compared with healthy subjects. Pharmacokinetic data on duloxetine is limited in patients with mild or moderate renal impairment.

Hepatic impairment

Moderate liver disease (Child Pugh Class B) affected the pharmacokinetics of duloxetine.

Paediatric population

Pharmacokinetics of duloxetine in paediatric patients aged 7 to 17 years with major depressive disorder following oral administration of 20 to 120 mg once daily dosing regimen was characterized using population modelling analyses based on data from 3 studies. The model-predicted duloxetine steady state plasma concentrations in paediatric patients were mostly within the concentration range observed in adult patients.

Breast-feeding mothers

The disposition of duloxetine was studied in 6 lactating women who were at least 12-weeks postpartum. Duloxetine is detected in breast milk, and steady-state concentrations in breast milk are about one-fourth those in plasma. The amount of duloxetine in breast milk is approximately 7 g/day while on 40 mg twice daily dosing. Lactation did not influence duloxetine pharmacokinetics.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content:

Sugar spheres(30-35#Pharm-A-Spheres), Hypromellose (Pharmacoat 606), Taic (Luzenac Pharma Ultra Micronized grade), Purified water*, Sucrose (Compressuc PS), Hypromellose phthalate (HP-55), Triethyl citrate (ET505), Isopropyl Alcohol*, Dichloromethane*

*-Processing solvent and does not appear in the final product, except in traces.

Capsule shell:

Cymbalox Capsules 30 mg

Gelatin, Titanium dioxide, FD & C Blue # 2, Green ink

Green ink contains : Shellac NF, Dehydrated Alcohol, Isopropyl Alcohol, Butyl Alcohol, Propylene Glycol, Strong Ammonia Solution NF, Titanium Dioxide, Yellow Iron Oxide NF, FD & C#2 Aluminium lake

Cymbalox Capsules 60 mg

Gelatin, FD&C Blue 2, Titanium dioxide, Iron oxide Yellow, White ink

White ink contains: Shellac NF, Dehydrated Alcohol, Isopropyl Alcohol, Butyl Alcohol, Propylene glycol, Strong Ammonia Solution, Potassium Hydroxide, Titanium dioxide

6.2 Shelf life

24 months

6.3 Special precautions for storage

Store in a dry place below 30°C.

6.4 Pack size

Duloxetine Hydrochloride Delayed Release Capsules are packed in Alu/Alu Blister pack (OPA/ALLU/PVC).

7 capsules per blister. Such 4 blisters are further packed in a carton along with pack insert.

6.5 Special precautions for disposal

No special requirements.

Manufactured by:

Alembic Pharmaceuticals Limited

(Formulation Division),

At- Panelav, Tal- Halol, City: Panelav,

Dist: Panchmahal, Gujarat State, India.

Product Registration Holder in MY:

GENPHARMA SDN. BHD.

Lot 5016, Jalan Teratai,

5 ½ Mile Off Jalan Meru,

41050 Klang, Selangor, Malaysia.

7. DATE OF REVISION:

27 October 2025