

## HOVID CHEWETTE KIDS-MART MULTI CHEWABLE TABLET

VICHE00-0\_4

### Description

Round, 15mm in diameter, light purple, mottled uncoated chewable tablet with blackcurrant flavour, bevel-edged and flat faces.

### Composition

Each tablet contains :

|                  |             |
|------------------|-------------|
| Vitamin A        | 1500.00 i.u |
| Vitamin D3       | 100.00 i.u  |
| Thiamine         | 0.25 mg     |
| Riboflavine      | 0.25 mg     |
| Nicotinamide     | 2.50 mg     |
| Pyridoxine       | 0.21 mg     |
| Cyanocobalamin   | 2.00 mcg    |
| Folic Acid       | 0.25 mg     |
| Pantothenic Acid | 3.00 mg     |
| Ascorbic Acid    | 50.00 mg    |
| Lysine           | 40.00 mg    |

### Indications

As a health supplement.

### Contraindications

Not recommended for patients with hypervitaminosis A or D, hypercalcaemia, renal osteodystrophy with hyperphosphatemia and Leber's disease.

### Precautions

- Keep out of reach of children/Jauhi daripada kanak-kanak
- Caution in patients with arteriosclerosis, renal, hepatic or cardiac function impairment, hyperphosphatemia, hypersensitivity to effects of vitamin D, gout, haemochromatosis, sideroblastic anaemia, thalassaemia, hyperoxaluria, peptic ulcer, diabetes mellitus and glaucoma.

### Main Side/Adverse Effects

Diarrhoea, itching, headache, flushing or redness of skin, nausea or vomiting, stomach pain and yellow discolouration of urine may occur occasionally.

### Drug Interactions

- Effects of levodopa are reversed by pyridoxine.
- Co-administration of probenecid may decrease gastrointestinal absorption and urinary excretion of riboflavine.
- Prolonged administration of aspirin may reduce blood levels of ascorbic acid.
- Vitamin E may facilitate absorption, hepatic storage, utilization and reduce toxicity of vitamin A; excessive doses of Vitamin E may deplete vitamin A stores.
- Avoid concurrent use of medications containing cholestyramine, colestipol, digitalis glycosides, magnesium-containing antacids, mineral oil, phenobarbital, phenytoin, high doses of phosphorus-containing preparations unless otherwise directed by physician.

### Overdosage

Clinical features : Diarrhoea, drowsiness, headache, peeling of skin, general feeling of discomfort or illness or weakness, stomach pain, vomiting, nausea, papilledema, intracranial hypertension, portal hypertension, anorexia, pruritus, polyuria, thirst.

Treatment for overdosage :

Symptomatic and supportive measures.

### Dosage and Administration

Children 4-12 years old : Chew 1 tablet once daily with meal.

The information given here is limited. For further information, consult your doctor or pharmacist.

|                                          |                                                                                                 |
|------------------------------------------|-------------------------------------------------------------------------------------------------|
| Storage                                  | : Store below 30 °C. Protect from light and moisture.                                           |
| Presentation/Packing                     | : Bottle of 60's.                                                                               |
| Marketing Authorization Holder /         | <b>HOVID Berhad,</b>                                                                            |
| Product Registration Holder (Malaysia) : | 121, Jalan Tunku Abdul Rahman (Jalan Kuala Kangsar), 30010 Ipoh, Perak, Malaysia.               |
| Manufactured by                          | : <b>HOVID Berhad,</b> Lot 56442, 7½ Miles, Jalan Ipoh / Chemor, 31200 Chemor, Perak, Malaysia. |
| Revision date                            | : December 2020                                                                                 |