

USER INFORMATION
READ CAREFULLY

RinsA CGC 50

(Chlorhexidine Gluconate 0.05% w/v and Cetrimide 0.5% w/v Irrigation Solution)
Sterile Topical Antiseptic Solution

Product Description:

RinsA CGC 50 is a clear, orange and sterile irrigation solution.

Pharmacodynamics:

Chlorhexidine: Chlorhexidine is an antiseptic and disinfectant which is effective against a wide range of vegetative Gram-positive and Gram-negative organisms, some viruses and some fungi. It is ineffective against bacterial spores at room temperature, and acid-fast bacteria are inhibited but not killed. It is more active against Gram-positive than Gram-negative bacteria and some species of *Pseudomonas* and *Proteus* are relatively less susceptible. Chlorhexidine is most active at a neutral or slightly acid pH and its activity may be reduced by blood and other organic matter.

Cetrimide: Cetrimide is a quaternary ammonium disinfectant with properties typical of cationic surfactants. Solutions of these surfactants have emulsifying and detergent properties and bactericidal activity against both Gram-positive and Gram-negative organisms but higher concentrations are necessary to kill the latter. The combined detergent and antibacterial properties of cetrimide make it useful in cleansing dirty or infected wounds. It is however, relatively ineffective against bacterial spores, acid-fast bacteria, viruses and fungi.

Pharmacokinetics :

Chlorhexidine is poorly absorbed from the gastrointestinal tract and skin.

Indication :

RinsA CGC 50 is for cleansing and irrigating the skin and dirty wounds where an added surfactant effect is required.

Recommended Dosage :

Rinse the area to be cleaned with water, apply the minimum amount of RinsA CGC 50 necessary to cover the wound area and wash gently. Rinse again thoroughly. Apply to wound as necessary. Discard remaining solution after use.

Route of Administration :

Topical

Contraindications :

RinsA CGC 50 should not be used in patients with known hypersensitivity to either chlorhexidine or cetrimide. It should not be used to irrigate the brain, meninges, eyes or perforated eardrum.

RinsA CGC 50 also should not be used in body cavities or as enema.

Warnings and Precautions :

For external use only. RinsA CGC 50 is not intended for injection and for irrigation only. It is not isotonic and is haemolytic. Cetrimide may have a prolonging effect on the wound healing process. Use of this antiseptic should be restricted to infection control and cleansing of wounds rather than general wound management.

Paediatric Use

Use with care in neonates, particularly in premature infants. Chlorhexidine may cause irritation or chemical burns.

RinsA CGC 50 contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare. RinsA CGC 50 should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

If any signs of symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Interactions with Other Medicaments:

Chlorhexidine salts and cetrimide are incompatible with soaps and other anionic materials.

Statement On Usage During Pregnancy and Lactation:**Use in Pregnancy - Category A**

Chlorhexidine has been used by a large number of pregnant women and women of childbearing age without any proven increase in the frequency of malformations or other direct or indirect harmful effects on the fetus having been observed.

Based on negligible topical absorption and a complete lack of any unusual reported effects when used during pregnancy, chlorhexidine would not appear to present a risk to pregnant women. No special precautions or changes in methods of application or use of chlorhexidine seem needed in pregnant health care workers or patients.

Lactation

The product is safe for use in lactation.

Adverse Effects/Undesirable Effects:

Irritative skin reactions and hypersensitivity reactions to chlorhexidine have been reported. In the event that these reactions occur, discontinue use.

Chlorhexidine may cause anaphylaxis.

Cetrimide may have a prolonging effect on the wound healing process.

Immune system disorders

Frequently not known: Hypersensitivity including anaphylactic shock.

Overdose and Treatment:**In case of accidental or deliberate oral poisoning or ingestion**

In case of accidental or deliberate oral poisoning or ingestion, chlorhexidine which is taken orally is poorly absorbed. Immediate dilution with (120 - 240mL) water or milk (not to exceed 15mL/kg in children) is recommended. Supportive measures are also appropriate.

In case of accidental intravenous administration

In case of accidental intravenous administration, transfusion may be necessary to counteract the haemolysis.

Storage Condition:

Do not store above 30°C. Protect from light.

Shelf Life:

2 years from manufacturing date in the proposed storage condition.

Do not use after expiry.

Dosage Form and Packaging Available:

30 mL x 20 LDPE Ampoules per box

Manufacturer/Product Registration Holder:

AIN MEDICARE SDN. BHD.

Lot 4933, 4934 & PT2464, Jalan 6/44, Kaw. Perindustrian Pengkalan Chepa 2,
16100 Kota Bharu, Kelantan, MALAYSIA

Date of Revision: 07.12.2020
