
Unimed's ChlorSept 5% Antiseptic Solution

Name and Strength of Active Substance:

25.0% v/v of Chlorhexidine gluconate 20% solution equivalent to chlorhexidine gluconate.....5% w/v

Description:**Product Description**

A clear reddish pink solution with lavender fragrance

Pharmacodynamics

Chlorhexidine is a bisbiguanide antiseptic that is bactericidal (kills bacteria) or bacteriostatic (inhibits bacterial growth) against a wide range of Gram-positive and Gram-negative bacteria. Chlorhexidine kills bacteria by binding cationic molecule to negatively charged bacterial cell walls and microbial complexes leading to precipitation of cytoplasmic contents of bacterial cell wall which results in cell death. At lower concentration, it acts as a bacteriostatic by alteration of bacterial cell osmotic equilibrium and leakage of potassium and phosphorus.

Pharmacokinetics

Chlorhexidine is poorly absorbed from the skin.

Indications

Unimed's ChlorSept 5% Antiseptic Solution used for preoperative skin disinfection and for wounds or burns.

Recommended Dose

Pre-Operative Skin Disinfection: To be used diluted with 1: 10 in 70% alcohol for preoperative skin disinfection

General cleansing for wounds or burns: To be used with 1: 100 in 70% alcohol for wounds or burns.

Route of Administration

For topical use only.

Contraindications

Hypersensitivity including anaphylactic shock.

Warning and Precautions

For external use only

Keep this medicine out of the reach of children.

Unimed's ChlorSept 5% Antiseptic Solution contains chlorhexidine.

Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare.

Unimed's ChlorSept 5% Antiseptic Solution should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine. If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Interactions with Other Medicaments

Incompatible with soaps or other anionic agents.

Pregnancy & Lactation

No effects during pregnancy and lactation are anticipated, since systemic exposure to chlorhexidine gluconate is negligible.

Adverse Effects / Undesirable Effects

Skin and subcutaneous tissue disorder. Dermatitis, pruritus, erythema, eczema, rash, skin irritation, blisters

Immune system disorder

Frequency not known: Hypersensitivity including anaphylactic shock

Symptoms and Treatment of Overdose

Chlorhexidine gluconate taken orally is poorly absorbed into the blood. Accidental ingestion may require gastric lavage using milk, egg white, gelatine or mild soap. Employ supportive measures as appropriate. Haemolysis has been reported following ingestion or accidental intravenous administration. Haemolysis may necessitate blood transfusion.

Effects on Ability to Drive and Use Machine

No effects on ability to drive and machine use

Storage Condition

Keep container tightly closed. Protect from heat and light. Store below 30°C

Dosage forms and packaging available

120ml and 1000ml HDPE Bottle

Date of Revision:

2/11/2020

Registration Number:

MAL

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas,

Pulau Pinang, Malaysia