

Front - 80mm

Back - 80mm

Unimed's ChlorSept 0.5% with IPA 75% Antiseptic Solution with Emollient

Composition and Strength of Active Ingredients:

2.5% v/v of Chlorhexidine Gluconate 20% solution equivalent to Chlorhexidine Gluconate 0.5% v/v
Isopropyl alcohol 75% v/v

Product Description

A clear pink colour solution with rose fragrance.

Pharmacodynamics

Chlorhexidine is effective against a wide range of Gram negative and Gram positive vegetative bacteria, yeasts, dermatophyte fungi and lipophilic viruses. It is inactive against bacterial spores except at elevated temperatures.

Isopropyl alcohol is a bactericidal agent that acts by the denaturation of proteins. It is a rapidly bactericidal and a fast-acting broad-spectrum antiseptic, but is not considered persistent. At concentrations of 70% v/v, it is a much effective antibacterial preservative than ethanol (95%). The bactericidal effect of aqueous solutions increases steadily as the concentration approaches 100% v/v. However, isopropyl alcohol is ineffective against bacterial spores.

Pharmacokinetics

Chlorhexidine is cationic nature it binds strongly to skin mucosa and other tissues and is thus very poorly absorbed. There are, as a consequence, no general pharmacological studies on chlorhexidine available and its effects on internal organs are minimal. No detectable blood levels have been found in man following oral use and percutaneous absorption, if it occurs at all, is insignificant.

As for the pharmacokinetic properties, there is little absorption of isopropyl alcohol through intact skin. It is readily absorbed from the gastrointestinal tract and may be slowly absorbed through intact skin. Isopropyl alcohol is metabolized more slowly than ethanol, primarily to acetone. Metabolites and unchanged isopropyl alcohol are mainly excreted in the urine.

Indication

To be used undiluted for pre-operation surgical hand disinfection, antiseptic hand disinfection, and disinfection of patient's skin.

Recommended Dosage

Pre-op surgical hand disinfection: Spread 5ml thoroughly over both hands and forearms, rubbing vigorously. When dry apply a further 5ml and repeat procedure.

Antiseptic hand disinfection: Spread 3ml thoroughly over the hands and wrist rubbing vigorously until dry.

Disinfection of patient's skin: Prior to surgery, apply to a sterile swab and rub thoroughly over the operation site for a minimum of 2 mins.

Route of Administration

Topical

Contraindications

Known hypersensitivity to Unimed's ChlorSept 0.5% with IPA 75% Antiseptic Solution with Emollient or any of its components, especially those with a known history of Chlorhexidine-related allergic reactions.

Warning and Precautions

Unimed's ChlorSept 0.5% with IPA 75% Antiseptic Solution with Emollient contains chlorhexidine.

Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is unknown, but available literature suggests this is likely to be very rare.

Unimed's ChlorSept 0.5% with IPA 75% Antiseptic Solution with Emollient should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine. If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated.

Interactions with Other Medicaments

Chlorhexidine salts are incompatible with soaps and other anionic materials. Activity may be reduced in the presence of suspending agents such as alginates and tragacanth, insoluble powders such as kaolin, and insoluble compounds of calcium, magnesium, and zinc.

As for the isopropyl alcohol, it should not be brought into contact with some vaccines and skin test injections (patch tests) and you should consult the vaccine manufacturer's literature when in doubt. Also, isopropyl alcohol is known to be incompatible with oxidizing agents such as hydrogen peroxide and nitric acid, which may cause decomposition.

Pregnancy & Lactation

No special precautions need to be taken when used in pregnancy and lactation. Thus, Unimed's ChlorSept 0.5% with IPA 75% Antiseptic Solution with Emollient can be used during pregnancy or during breast-feeding.

Side Effects

Immune system disorders.

Frequency not known: Hypersensitivity including anaphylactic shock.

Symptoms and Treatment of Overdose

There are no reports of overdose with this product. However, Chlorhexidine Gluconate taken orally is poorly absorbed into the blood. Accidental ingestion may require appropriate gastric lavage using milk, egg white, gelatine or mild soap. Haemolysis has been reported following ingestion or accidental intravenous administration which necessitate blood transfusion.

Effects on Ability to Drive and Use Machine

Unimed's ChlorSept 0.5% with IPA 75% Antiseptic Solution with Emollient has no influence on ability to drive and use machines.

Storage Condition

Keep container tightly closed. Protect from heat and light. Store below 30°C.

Dosage Forms and Packaging Available

500 ml HDPE Bottle with pump cap.

Registration Number:

MAL21056025XZ

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,

Lintang Bayan Lepas 1, 11900 Bayan Lepas,

Pulau Pinang, Malaysia.

Date of Revision:

11 May 2021

PI0068-0

115mm