



ALLERSOOTHE ELIXIR 5MG/5ML

Sugar Free

DESCRIPTION: A clear orange syrupy liquid with banana and vanilla flavour.

Each 5ml contains Promethazine Hydrochloride 5mg with Sodium Benzoate 0.2% w/v as preservative. Sugar free.

PHARMACODYNAMICS: Promethazine hydrochloride is an antihistamine with some anticholinergic, anti-serotonergic, and marked local anaesthetic properties. It has a prolonged antihistaminic action. It also has anti-emetic and sedative actions.

PHARMACOKINETICS: Promethazine is distributed widely in the body. It enters the brain and crosses the placenta. Promethazine is slowly excreted via urine and bile. Phenothiazines pass into the milk at low concentrations.

INDICATIONS: Allergic conditions, motion sickness, nausea & vomiting and hypnosis.

CONTRAINDICATIONS: Hypersensitivity to Promethazine Hydrochloride or other antihistamine of similar chemical structure.

ADVERSE EFFECTS: The most common effects of antihistamines are sedation, inability to concentrate, lassitude, dizziness, hypotension, muscular weakness, and in-coordination. Other side effects of antihistamines include gastro-intestinal disturbances such as nausea, vomiting, diarrhoea or constipation and epigastric pain.

Antihistamines may also produce headache, blurred vision, tinnitus, elation or depression, irritability, nightmares, anorexia, difficulty in micturition, dryness of the mouth, tightness of the chest, and tingling heaviness, and weakness of the hands.

Although cardiovascular side-effects are rare, minor increase in blood pressure and occasional mild hypotension have been reported, Leucopenia and rarely agranulocytosis, jaundice, extrapyramidal reactions and photosensitivity reactions have been reported.

INTERACTIONS WITH OTHER MEDICAMENTS: Allersoothe Elixir will enhance the action of any anticholinergic agent, tricyclic anti-depressant, sedative or hypnotic. Alcohol should be avoided during treatment. Allersoothe Elixir may interfere with immunological urine pregnancy tests to produce false-positive or false-negative results. Allersoothe Elixir should be discontinued at least 72 hours before the time start of skin tests as it may inhibit the cutaneous histamine response thus producing false-negative results.

PRECAUTIONS: Since antihistamines have some anticholinergic properties, they should be used with care in conditions liable to be exacerbated or adversely affected by atropine such as glaucoma and prostatic hypertrophy. As for all phenothiazine derivatives, Promethazine Hydrochloride should be used cautiously in patients with hepatic diseases.

WARNINGS: Antihistamines may cause drowsiness and may increase the effects of alcohol. Patients should be warned about engaging in activities requiring mental alertness. Antihistamines may enhance the sedative effect of central nervous system depressants including alcohol, barbiturates, hypnotics, narcotic analgesics, sedatives and tranquilizers.

- Allersoothe Elixir 5mg/ 5ml should not be used in pediatric patients less than 2 years of age because of the potential for fatal respiratory depression.
- To be used with caution and doctor's advice in children 2 to 6 years of age (when used for treatment of cough and cold).

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE: Because the duration of action may be up to 12 hours, patients should be advised that if they feel drowsy they should not drive or operate heavy machinery.

PREGNANCY AND LACTATION: Do not use this product in pregnancy unless recommended by your health professionals. The use of this product is not recommended in the 2 weeks prior to delivery in view of the risk of irritability and excitement in the neonates. The amount excreted in milk is insignificant. However, there are risks of neonatal irritability and excitement.

ROUTE OF ADMINISTRATION: Oral

RECOMMENDED DOSAGE:

Allergic Conditions

Dosage	Single Dose at Night	2 – 3 Times A Day Dose
2 – 5 years	5 – 15 ml	5 ml
6 – 12 years	10 – 25 ml	10 ml

Hypnosis

Dosage	Single Dose at Night
2 – 5 years	5 – 15 ml
6 – 12 years	10 – 25 ml

Motion Sickness

Dosage	
2 – 5 years	5 ml to be taken the night before the journey. To be repeated after 6–8 hours as required.
6 – 12 year	10 ml to be taken the night before the journey. To be repeated after 6–8 hours as required.

Established Sickness and Vomiting Due to Other Causes

Dosage	
2 – 5 years	5 ml every 4 to 6 hours to a maximum daily dose of 15 ml.
6 – 12 year	10 ml every 4 to 6 hours to a maximum daily dose of 25 ml.

OVERDOSAGE: Antihistamines overdose may vary from central nervous system depression to stimulation. Atropine like signs and symptoms such as dry mouth, fixed dilated pupils, flushing, and gastro-intestinal symptoms may also occur.

Treatment - Small quantity- emesis unless drowsy. In severe overdose, the stomach should be emptied by aspiration and lavage. The patient should be kept quiet to minimize the excitation. Convulsion may be controlled with diazepam given intravenously.

PACKING: 100ml amber bottle (PET)

STORAGE: Keep container well closed, Store below 30°C. Protect from strong light. Keep out of reach of children.

RECOMMENDED SHELF LIFE: 18 months

PRODUCT REGISTRATION HOLDER AND MANUFACTURER:

HOE Pharmaceuticals Sdn. Bhd.

Lot 10, Jalan Sultan Mohamed 6, Bandar Sultan Suleiman, 42000 Port Klang, Selangor, MALAYSIA

Rev. Date: Mar. 2021