

IRONFORT+B12 INJECTION

Description and Composition

IRONFORT+B12 INJECTION is a dark brown solution. Each ml contains 100mg Iron dextran and 0.08mg Cyanocobalamin as its active ingredients and 5mg Phenol as its preservative.

Pharmacodynamics

The product is a macromolecular complex solution containing 100 mg of iron per mL: stimulation of hematopoiesis by progressive release of the dextran complex.

It allows, at the recommended doses, an almost immediate reconstitution of the iron reserves necessary for the formation of hemoglobin and the recharging of iron-based enzymes which play a decisive role in growth and resistance to infections.

Pharmacokinetics

The product is quickly and completely absorbed.

Indication

Treatment and prevention of iron deficiency anemia in piglets and calves.

Recommended Dose

Intramuscular route.

Piglets: 100mg to 200mg of iron dextran per piglet, i.e. 1ml to 2ml all at once during one of the first days of the piglet's life, preferably in the muscles of the neck.

Calves: 300mg to 400mg of iron dextran per calf, i.e. 3ml to 4ml all at once during one of the first days of the calf's life, preferably in the muscles of the neck.

Duration of treatment: one injection to be repeated if necessary after 8 days to 1 month.

Route of Administration

To be given by intramuscular route.

Contraindications

None known.

Warnings and Precautions

Special warnings for each target species:

None known.

Special precautions for use

Special precautions for use in animals:

Strictly observe the rules of asepsis when injecting.

Act with caution in case of particular sensitivity to iron dextran, due to nutritional predisposition of maternal origin.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

None known.

Interactions with Other Medicaments

None known.

Pregnancy and Lactation

Not applicable.

Side Effects

Staining at the injection site.

Very rare cases of piglet deaths have occurred after parenteral administration of iron preparations.

These deaths have been associated with genetic factors or with deficiencies in vitamin E and / or selenium.

Symptoms and Treatment of Overdose

None known.

Storage Condition

Store below 30°C. Protect from light.

Shelf life

3 years.

Following withdrawal of first dose, use the product within 24 hours. Any unused material should be discarded.

Withdrawal Period

Meat and offal: Zero days.

Milk: Zero days.

Packing

20ml, 50ml, 100ml, 250ml.

Revised date: 16.4.2021

	Manufacturer & Product Registration Holder
	Range Pharma Sdn Bhd No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia Tel: 603-61568708 Fax: 603-61568707 Email: info@rangepharma.com