

成品尺寸：145X210mm

色标

■ C0 M0 Y0 K100

HP-DESLORATADINE 5MG FILM-COATED TABLETS

QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 5 mg desloratadine.

PHARMACEUTICAL FORM

Film-coated, light blue, round tablets.

CLINICAL PARTICULARS

Therapeutic indications

HP- Desloratadine 5mg Film-Coated Tablets are indicated for the rapid relief of symptoms associated with allergic rhinitis, such as sneezing, nasal discharge and itching, congestion/stuffiness, as well as ocular itching, tearing and redness.

HP-Desloratadine 5mg Film-Coated Tablets are also indicated for the relief of symptoms associated with chronic idiopathic urticaria such as the relief of itching and the size and number of hives.

Posology and method of administration

Adults and Adolescents (12 years of age and older): One HP-Desloratadine 5mg Film-Coated Tablets once a day regardless of mealtime. For oral use.

Contraindications

Hypersensitivity to the active substance or to any of the excipients.

Special warnings and precautions for use

Efficacy and safety of HP-Desloratadine 5mg Film-Coated Tablets in children under 12 years of age have not been established.

Desloratadine should be administered with caution in patients with a medical or family history of seizures.

In particular, young children may be more susceptible to developing new seizures under desloratadine treatment. Healthcare providers may consider discontinuing desloratadine in patients who experience a seizure while on treatment.

Effects on ability to drive and use machines

No effects on the ability to drive and use machines have been observed (see Pharmacodynamic properties).

Interaction with other medicinal products and other forms of interaction

No clinically relevant interactions with HP-Desloratadine 5mg Film-Coated Tablets were observed. There was no effect of food or grapefruit juice on the disposition of desloratadine.

HP-Desloratadine 5mg Film-Coated Tablets taken concomitantly with alcohol did not potentiate the performance impairing effects of alcohol

Fertility, pregnancy and lactation

HP-Desloratadine 5mg Film-Coated Tablets is not to be used during pregnancy unless the potential benefits outweigh the risks.

Desloratadine is excreted into breast milk; therefore, the use of HP-Desloratadine 5mg Film-Coated Tablets is not recommended in breast-feeding women.

Undesirable effects

The most frequent adverse events reported were fatigue, dry mouth, and headache.

Very rare cases of hypersensitivity reactions, including anaphylaxis and rash, have been reported during the marketing of desloratadine. In addition, cases of tachycardia, palpitations, psychomotor hyperactivity, somnolence, seizures, elevations of liver enzymes, hepatitis, increased bilirubin and increased appetite have been reported very rarely.

Overdose

In the event of overdose, consider standard measures to remove unabsorbed active substance. Symptomatic and supportive treatment is recommended.

In adults and adolescents, in which up to 45 mg of desloratadine was administered (9 times the clinical dose), no clinically relevant effects were observed.

Desloratadine is not eliminated by hemodialysis; it is not known if it is eliminated by peritoneal dialysis.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Mechanism of action

Desloratadine is a non-sedating long-acting histamine antagonist with potent, selective peripheral H1- receptor antagonist activity.

Desloratadine has demonstrated antiallergic, antihistaminic, and anti- inflammatory activity.

Desloratadine is the primary active metabolite of loratadine. There are no qualitative or quantitative differences in the toxicity profile of desloratadine and loratadine at comparable levels of exposure to desloratadine.

After oral administration, desloratadine selectively blocks peripheral histamine H1-receptors because the drug is effectively excluded from entry to the central nervous system (CNS).

In addition to antihistaminic activity, desloratadine has demonstrated antiallergic and anti-inflammatory activity. These shown that desloratadine inhibits the broad cascade of events that initiate and propagate allergic inflammation, including

海口新明印刷有限公司 HAIKOU XINMING PRINTING CO., LTD		说明/Explain	
文件信息/File information		尊敬的客户/Dear customer: 请您仔细审阅以上内容及图稿文字字体、大小、排列、内容、色相、层次、盒型结构、尺寸, 并签字确认, 我们将以此稿作为制版依据。 Please confirm above content, and then text font, size, arrangement, content, hue, level, structure, size on the drawing, we will follow to make plate.	
☐ 纸盒/Box ☒ 说明书/Leaflet			
制作文件名/Product Name: 马来西亚-地氯雷他定片说明书		审核人 Reviewed by	
成品尺寸/Finish Product Size: 145x210MM		审核人 Reviewed by	
制作日期/Design Time: 2022.5.31		批准人 Approved by	

成品尺寸：145X210mm

色标

■ C0 M0 Y0 K100

- the release of proinflammatory cytokines including IL-4, IL-6, IL-8, IL-13,
- the release of important proinflammatory chemokines such as RANTES (Regulated upon Activation, Normal T-cell Expressed and Secreted),
- superoxide anion production by activated polymorphonuclear neutrophils,
- eosinophil adhesion and chemotaxis,
- the expression of the adhesion molecules such as P-selectin,
- IgE-dependent release of histamine, prostaglandin (PGD2), and leukotriene (LTC4),
- the acute allergic bronchoconstrictor response and allergic cough in animal models.

No statistically or clinically relevant cardiovascular effect was observed when 20 mg of desloratadine was administered daily for 14 days. No prolongation of the QTc interval was seen when desloratadine was administered at a dose of 45 mg daily (nine times the clinical dose) for ten days.

Desloratadine does not readily penetrate the central nervous system. At the recommended dose of 5 mg daily, there was no excess incidence of somnolence. Desloratadine 5mg Film-Coated Tablets even at a dose of 7.5 mg daily did not affect psychomotor performance. A single dose of desloratadine 5 mg did not affect standard measures of flight performance including exacerbation of subjective sleepiness or tasks related to flying.

No clinically relevant changes in desloratadine plasma concentrations were observed in multiple-dose ketoconazole, erythromycin, azithromycin, fluoxetine and cimetidine interaction. Co-administration of alcohol did not increase the alcohol-induced impairment in performance or increase in sleepiness.

In adult and adolescent patients with allergic rhinitis (AR), HP-Desloratadine 5mg Film-Coated Tablets were effective in relieving symptoms such as sneezing, nasal discharge and itching, congestion/stuffiness, as well as ocular itching, tearing and redness, and itching of palate. HP-Desloratadine 5mg Film-Coated Tablets effectively controlled symptoms for 24 hours.

In patients with seasonal allergic rhinitis (SAR) and concurrent asthma, desloratadine was shown to be effective in reducing the symptoms of SAR (rhinorrhea, nasal congestion, nasal itching and sneezing, itching/burning eyes, tearing/watering eyes, redness of eyes, and itching of ears or palate) and asthma (coughing, wheezing, difficulty breathing), and decreasing beta-agonist use.

In adults and adolescents with chronic idiopathic urticaria (CIU), HP-Desloratadine 5mg Film-Coated Tablets were effective in relieving pruritus and decreasing the size and number of hives as early as 1 day after initiation of treatment. The effects were sustained over the 24-hour dosing interval. Treatment with HP-Desloratadine 5mg Film-Coated Tablets also improved sleep and daytime function, as measured by reduced interference with sleep and routine daily activities.

HP-Desloratadine 5mg Film-Coated Tablets was effective in alleviating the burden of seasonal allergic rhinitis. The greatest amelioration was seen in the domains of practical problems and daily activities limited by symptoms.

Pharmacokinetic properties

Desloratadine plasma concentrations can be detected within 30 minutes of desloratadine administration. Desloratadine is well absorbed with maximum concentration achieved after approximately 3 hours; the terminal phase half-life is approximately 27 hours. The degree of accumulation of desloratadine was consistent with its half-life (approximately 27 hours) and a once daily dosing frequency. In adults and adolescents, the bioavailability of desloratadine was dose proportional over the range of 5 mg to 20 mg.

Desloratadine is moderately bound (83% - 87%) to plasma proteins. There is no evidence of clinically relevant drug accumulation following once daily dosing of desloratadine (5 mg to 20 mg) for 14 days.

The enzyme responsible for the metabolism of desloratadine has not been identified yet, and therefore some interactions with other drugs cannot be fully excluded. In vivo studies with specific inhibitors of CYP3A4 and CYP2D6 have shown that these enzymes are not important in the metabolism of desloratadine. Desloratadine does not inhibit CYP3A4 or CYP2D6 and is neither a substrate nor an inhibitor of P-glycoprotein.

7.5 mg dose of desloratadine, there was no effect of food (high-fat, high caloric breakfast) on the disposition of desloratadine. Grapefruit juice had no effect on the disposition of desloratadine.

PHARMACEUTICAL PARTICULARS

Special precautions for storage

Store below 30 °C.
Store in the original container. Protect from excessive moisture.

Nature and contents of container

Desloratadine tablets is presented in blisters consisting of PA/AL/PVC and aluminum foil. Packs of 1×10's, 3×10's, 10×10's tablets.

MANUFACTURER

Hainan Poly Pharm. Co., Ltd.,
Guilinyang Economic Development Area,
Haikou, Hainan, China 571127

DATE OF REVISION OF THE TEXT

29/11/2021

海口新明印刷有限公司 HAIKOU XINMING PRINTING CO., LTD		说明/Explain	
文件信息/File information		尊敬的客户/Dear customer: 请您仔细审阅以上内容及图稿文字字体、大小、排列、内容、色相、层次、盒型结构、尺寸，并签字确认，我们将以此稿作为制版依据。 Please confirm above content, and then text font, size, arrangement, content, hue, level, structure, size on the drawing, we will follow to make plate.	
☐ 纸盒/Box ☒ 说明书/Leaflet			
制作文件名/Product Name: 马来西亚-地氯雷他定片说明书		审核人 Reviewed by	
成品尺寸/Finish Product Size: 145x210MM		审核人 Reviewed by	
制作日期/Design Time: 2022.5.31		批准人 Approved by	