



Package Insert

1. Name of the medicinal product

Medinazine Orapaste 0.1%w/w

2. Qualitative and quantitative composition

Each gram contains Dexamethasone 1.0mg

3. Pharmaceutical form

Orabase

Description: Cream white color orabase and smells of strawberry.

4. Clinical particulars

4.1 Therapeutic indications

Intractable stomatitis or glossitis associated with erosion or ulceration.

4.2 Posology and method of administration

Usually apply an adequate amount of Medinazine to the affected area 1 to several times daily. The dose may be increased or decreased depending on the condition of the patient.

4.3 Contraindications

Medinazine should not be used on the following patients: Patients with tuberculosis, viral or other purulent infections of the oral cavity.

Patients with a past history of hypersensitivity to Medinazine.

4.4 Special warnings and precautions for use

Unsuitable for phenylketonurics.

No food or fluid should be ingested for a short time after application of Medinazine Orapaste.

Medinazine Orapaste should by no means be applied to the eye.

Visual disturbance may be reported with systemic and topical corticosteroid use. If a patient presents with symptoms such as blurred vision or other visual disturbances, the patient should be considered for referral to an ophthalmologist for evaluation of possible causes which may include cataract, glaucoma or rare diseases such as central serous chorioretinopathy (CSCR) which have been reported after use of systemic and topical corticosteroids

4.5 Interaction with other medicinal products and other forms of interaction

Medinazine may interact with the following drugs and products:

- Aspirin, Cyclosporine, Diabetes medications, Diuretic, Ketoconazole, Phenobarbital, Phenytoin
- Rifampin, Warfarin

4.6 Pregnancy, breast-feeding and children

Pregnancy:

The safety of Medinazine when used during pregnancy has not yet been established. Its prolonged use on pregnant women or women suspected of pregnancy should be avoided.

Breast-feeding:

Please discuss the risks and benefits with your doctor.

Children:

The prolonged use of Medinazine may be associated with growth and development disturbances.



4.7 Effects on ability to drive and use machines

Medinazine does not affect the ability to drive or to operate the machine.

4.8 Adverse Effects

No severe side effects of Medinazine Orapaste have been reported except for the withdrawal of the drug from one patient for a complaint of numbness at the site of application. Adverse reactions

- (1) Infection of the oral cavity: Fungal or bacterial infection of the oral cavity may occur in the course of treatment with Medinazine Orapaste. In case that any symptom or sign of such an infection has appeared, an appropriate antifungal or antibacterial treatment should concomitantly be instituted. In cases where the symptoms or signs of the infection fail to subside rapidly, withdraw Medinazine Orapaste
- (2) Hypersensitivity: On manifestation of any allergic reaction, withdraw Medinazine Orapaste
- (3) Function of the pituitary-adrenocortical system: The prolonged use of Medinazine Orapaste may be associated with hypofunction of the pituitary-adrenocortical system.
- (4) Vision, blurred

4.9 Overdose

Do not use more than prescribed dose. Taking more medication will not improve your symptoms; rather they may cause poisoning or serious side-effects. If you suspect that you or anyone else who may have overdosed of Medinazine, please go to the emergency department of the closest hospital or nursing home. Bring a medicine box, container, or label with you to help doctors with necessary information.

Do not give your medicines to other people even if you know that they have the same condition or it seems that they may have similar conditions. This may lead to overdosage.

Please consult your physician or pharmacist or product package for more information.

5. Pharmacological properties

Dexamethasone is a synthetic adrenocortical steroid and exerts anti inflammatory effects by the same mechanism of action as naturally occurring glucocorticoids, but the mineralocorticoid action of dexamethasone is weaker than those of naturally occurring glucocorticoids.

Pharmacokinetic properties:

Dexamethasone is readily absorbed from the gastro-intestinal tract. Its biological half life in plasma is about 190 minutes. Binding of dexamethasone to plasma proteins is about 77% which is less than for most other corticosteroids. Up to 65% of a dose is excreted in urine within 24 hours. Clearance in premature neonates is reported to be proportional to gestational age, with a reduced elimination rate in the most premature. Its readily crosses the placenta with minimal inactivation.

6. Pharmaceutical particulars

6.1 List of excipients

Light liquid paraffin, White petrolatum, Butylated hydroxytoluene, Aspartame, Strawberry flavor, Sodium polyacrylate and Ethanol 0.04 % w/w

6.2 Incompatibilities

Not applicable.

6.3 Special precautions for storage

Store below 30°C. Do not refrigerate.



YUAN CHOU CHEMICAL & PHARMACEUTICAL CO., LTD

No. 272 , Nangang 3rd Rd., Nantou City , Nantou County 54067, Taiwan R.O.C

6.4 Nature and contents of container

Collapsible aluminum tubes with plastic screw caps.

Pack size: 5g tubes.

6.5 Special precautions for disposal and other handling

No special requirements.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. Name and Address of Manufacturer

Yuan Chou Chemical and Pharmaceutical Co. Ltd.

No. 272, Nangang 3rd Rd, Nantou City, Nantou County 54067, Taiwan.

8. Name and Address of Product Registration Holder

World Medicare Supplies Sdn. Bhd.

Pt 5740 (1st floor) Jalan TS 2/1E,

Taman Semarak 2, 71800 Nilai,

Negeri Sembilan.

9. Date of Revision of PI

January 2022