




BELCID FORTE SUSPENSION
BELCID FORTE SACHET (ORAL SUSPENSION)
Antacid, Antiflatulent



PRODUCT DESCRIPTION

White suspension with peppermint flavor.

COMPOSITION

Each 5 ml contains:

Aluminium Hydroxide Gel equivalent to	306 mg
Aluminium Hydroxide	
Magnesium Hydroxide USP	110 mg
Simethicone USP	30 mg

INDICATION

Use as an antacid for symptomatic relief of hyperacidity associated with the diagnosis of peptic ulcer, gastritis, peptic esophagitis, gastric hyperacidity, heart-burn, or hiatal hernia. As an anti-flatulent to alleviate the symptoms of gas, including postoperative gas pain.

RECOMMENDED DOSE

Adults and Children ≥ 12 years :

BELCID FORTE : 1 - 3 teaspoonfuls 4 times daily between meals, at bed time.

BELCID FORTE SACHET : 1 sachet 4 times daily between meals, at bed time.

PHARMACODYNAMICS

Pharmacological actions :

BELCID FORTE has an action of antacid, antiflatulence.

BELCID FORTE acts by neutralizing the gastric acid and also has antifoaming action.

Mechanism of actions :

The clinical use of antacids is based on their ability to increase the pH of gastric secretions: Aluminium hydroxide or oxide reacts with hydrochloric acid to form aluminium chloride and water. Magnesium hydroxide rapidly reacts with hydrochloric acid to form magnesium chloride and water.

Simethicone acts by coagulating the mucus-surrounded gas bubbles in the GI tract and expelling them.

PHARMACOKINETICS

Aluminium hydroxide or oxide is slowly solubilized in the stomach after neutralizing hydrochloric acid to form aluminium chloride, about 17 - 30% of the aluminium chloride is absorbed and is rapidly excreted by the kidney in patients with normal renal function. Magnesium hydroxide neutralizes hydrochloric acid in the stomach form magnesium chloride and about 15 - 30% of the magnesium chloride is absorbed and is rapidly excreted by the kidney in patients with normal renal function.

Simethicone is not absorbed from GI tract and is excreted unchanged in feces.

SIDE EFFECTS

The most common adverse effect with prolonged administration of antacids are constipation from Aluminium and diarrhea from Magnesium.

WARNING AND PRECAUTIONS

BELCID FORTE may cause dose-related rebound hyperacidity because they may increase gastric secretion or serum gastrin levels.

Use this drug with care in patients who have recently suffered massive upper GI hemorrhage.

CONTRAINDICATION

Should not take this medicine longer than 2 weeks.

Do not take the other medicine within 2 hours of administration of this drug.

INTERACTION OF OTHER MEDICAMENTS

BELCID FORTE may interfere with drug by:

Increasing the gastric pH altering disintegration, dissolution, solubility, ionization and gastric emptying time.

Absorption of weakly acidic drugs is decreased, possibly resulting in decreased drug effect (e.g. digoxin, phenytoin, chlorpromazine, isoniazid, chlorthalazepoxide, indomethacin, iron salts). Doses of these drugs should be spaced as far apart as possible from doses of BELCID FORTE.

Weakly basic drug absorption is increased possibly resulting in toxicity or adverse reactions (e.g. pseudoephedrine, levodopa).

Absorbing or binding drugs to their surface resulting in decreased bioavailability (e.g. tetracycline); therefore, doses of tetracyclines should be spaced 1 - 2 hours from doses of BELCID FORTE.

Increasing urinary pH affecting the rate of drug elimination. The effect is inhibition of the excretion of basic drugs (e.g. quinidine, amphetamines) and enhanced excretion of acidic drugs (e.g. salicylates).

PATIENTS INFORMATION

Further information can be obtained from the doctor or the pharmacist.

PREGNANCY AND LACTATION

BELCID FORTE should only be given in pregnancy when benefit outweighs risk.

(Please discuss and take as directed by your doctor)

Pregnancy

Antacids are generally considered safe as long as chronic high doses are avoided. Adequate and well-controlled studies in humans have not been done; however, there have been reports of antacids causing such adverse effects as hypercalcemia, hypomagnesemia, hypermagnesemia, and increased tendon reflexes in fetuses and/or neonates whose mothers were chronic users of aluminium, calcium and/or magnesium-containing antacids, especially in high doses. Studies have not been done in animals.

Lactation

Problems in humans have not been documented; although some aluminium, calcium and magnesium may be distributed into breast milk, the concentration is not great enough to produce an effect in the neonate.

SYMPTOMS AND TREATMENT OF OVERDOSE

Acute Overdosage

Acute Overdosage with antacid may cause osmotic catharsis, leading to fluid and electrolyte losses. Hypermagnesemia and central nervous system depression may follow overdoses of magnesium-containing antacid.

Treatment

Specific treatment is rarely needed in either suicidal or non-suicidal patients. Gastrointestinal decontamination is not indicated unless toxic coingestants are suspected.

Hypermagnesemia may be treated with fluids and loop diuretics. Hemodialysis reduced plasma levels of magnesium. Aluminium may be removed with deferoxamine followed by hemodialysis.

STORAGE

Do not store above 30°C. Avoid freezing.

SUPPLY

BELCID FORTE : PET Bottle of 120, 240 mL.

BELCID FORTE SACHET : Aluminium sachet 15 mL. 16 and 30 sachets in paper box.

Date of revision : March 2022



Manufactured by
BIOLAB CO., LTD.
625 Soi 7A Bangpoo Industrial Estate, Sukhumvit Road,
Moo 4, Prakasa, Muang, Samutprakarn, 10280, Thailand

EES026/A

Code : EES026/A		Draft Revision : 3		Final Artwork Revision : B	
Name : EES026-A_Rev.B_Leaflet_Belcid Forte Suspension & Sachet.ai					
Remark :					
Design by		Review by QCP		Approved by QCP	
Signature		Signature		Signature	
Date		Date		Date	
Review by RA			Approved by RA		
Signature			Signature		
Date			Date		
Specification	Size : 105 x 230 MM. (+/- 3 MM.)				
	Color : พิมพ์ 1 สี คือ สีดำ				
	Material : ใช้กระดาษปอนด์ น้ำหนัก 55 แกรม				
	Other :				
Note : - Draft Rev.0 = ส่วน SUSPENSION กับ SACHET (28/10/20) - Draft Rev.1 = ส่วน BELCID SUSP. หัวข้อ COMPOSITION (28/10/20) - Draft Rev.2 = เปลี่ยน Belcid Forte Sachet ตอน (05/11/20) - Draft Rev.3 = เปลี่ยนจาก " ORAL SUSPENSION " (07/03/22) ☐ ลูกคำพิจารณาอนุมัติฉบับร่าง ☐ ทำเสนอราคา ☒ RA ชื่นชม:เขียน ☐ ส่งพิมพ์ Color Guide BLACK BIOLAB					