
Unimed's ChlorSept 0.5% with Ethanol 77% Antiseptic Solution with Emollient

Composition and Strength of Active Ingredients:

Chlorhexidine gluconate.....	0.5% w/v
Ethanol	77% v/v

Product Description

A clear pink colour solution with rose fragrance.

Pharmacodynamics

Chlorhexidine is effective against a wide range of Gram negative and Gram positive vegetative bacteria, yeasts, dermatophyte fungi and lipophilic viruses. It is inactive against bacterial spores except at elevated temperatures.

Alcohol is bacteriostatic at low concentrations but has bactericidal activity at higher concentrations; it does not, however, destroy bacterial spores. The mechanism of action appears to be denaturation of proteins. Its bactericidal activity drops sharply when diluted below a 50 % concentration and the optimal bactericidal concentration is 60 to 90% by volume. Alcohol also has some fungicidal and virucidal activity.

Pharmacokinetics

Chlorhexidine is cationic nature it binds strongly to skin mucosa and other tissues and is thus very poorly absorbed. There are, as a consequence, no general pharmacological studies on chlorhexidine available and its effects on internal organs are minimal. No detectable blood levels have been found in man following oral use and percutaneous absorption, if it occurs at all, is insignificant.

Alcohol is rapidly distributed throughout the body. It readily crosses the placenta. Alcohol is mainly metabolised in the liver and is converted by alcohol dehydrogenase to acetaldehyde and is then further oxidised to acetate. A hepatic microsomal oxidising system is also involved. About 90% to 98% of alcohol is oxidised and the remainder is excreted unchanged by the kidneys and the lungs and also in breast milk, saliva, sweat and other secretions.

Indication

To be used undiluted for pre-operation surgical hand disinfection, antiseptic hand disinfection, and disinfection of patient's skin.

Recommended Dosage

Pre-op surgical hand disinfection: Spread 5ml thoroughly over both hands and forearms, rubbing vigorously. When dry apply a further 5ml and repeat procedure.

Antiseptic hand disinfection: Spread 3ml thoroughly over the hands and wrist rubbing vigorously until dry.

Disinfection of patient's skin: Prior to surgery, apply to a sterile swab and rub thoroughly over the operation site for a minimum of 2 mins.

Route of Administration

Topical

Contraindications

Do not use in patients with a known hypersensitivity to the product or any of its components, especially in those with a history of possible chlorhexidine related allergic reactions.

Do not use in contact with eyes, brain, meninges, middle ear or external ear with a perforated tympanic membrane.

Do not inject.

When use is to be followed by diathermy do not allow pooling of the fluid to occur and ensure that the skin and surrounding drapes are dry. Do not use in body cavities.

Warning and Precautions

Unimed's ChlorSept 0.5% with Ethanol 77% Antiseptic Solution with Emollient contains chlorhexidine. Chlorhexidine is known to induce hypersensitivity, including generalised allergic reactions and anaphylactic shock. The prevalence of chlorhexidine hypersensitivity is

unknown, but available literature suggests this is likely to be very rare. Unimed's ChlorSept 0.5% with Ethanol 77% Antiseptic Solution with Emollient should not be administered to anyone with a possible history of an allergic reaction to chlorhexidine.

If any signs or symptoms of a suspected hypersensitivity reaction such as itching, skin rash, redness, swelling, breathing difficulties, light headedness, and rapid heart rate develop, immediately stop using the product. Appropriate therapeutic countermeasures must be instituted as clinically indicated

Remove any soaked materials, drapes or gowns before proceeding with the intervention. Do not use excessive quantities and do not allow the solution to pool in skin folds or under the patient or drip of sheets or other material in direct contact with the patient. Where occlusive dressings are to be applied to areas previously exposed to this product, care must be taken to ensure no excess product is present prior to application of the dressing.

Interactions with Other Medicaments

None stated.

Pregnancy & Lactation

None stated.

Side Effects

Immune system disorders

- Frequency not known: Hypersensitivity including anaphylactic shock

Skin disorders

Frequency not known:

- Allergic skin reactions such as dermatitis, pruritus, erythema, eczema, rash, urticarial, skin irritation, and blisters.
- Chemical burns in neonates and infants

Symptoms and Treatment of Overdose

None stated.

Effects on Ability to Drive and Use Machine

None stated.

Storage Condition

Store below 30°C. Keep container tightly closed.

Dosage Forms and Packaging Available

500ml bottle.

Registration Number:

Manufactured by & Product Registration Holder

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia.

Date of Revision:

7 May 2022