

Aw. No.: 2000XXXXX

Front

2000XXXXX

SYNERRV FEXOFENADINE 120
SYNERRV FEXOFENADINE 180
FEXOFENADINE HYDROCHLORIDE 120 MG TABLETS USP
FEXOFENADINE HYDROCHLORIDE 180 MG TABLETS USP

COMPOSITION:

Synerrv Fexofenadine 120 (Fexofenadine Hydrochloride 120 mg Tablets USP)
Each film coated tablet contains:
Fexofenadine hydrochloride USP ... 120 mg
Excipients: Lactose spray dried, Pregelatinized starch, Croscarmellose sodium, Colloidal anhydrous silica, Magnesium stearate, Povidone and Opadry pink 03B54819
Colours: Iron Oxide Red, Iron Oxide Yellow and Titanium Dioxide
Synerrv Fexofenadine 180 (Fexofenadine Hydrochloride 180 mg Tablets USP)
Each film coated tablet contains:
Fexofenadine hydrochloride USP ... 180 mg
Excipients: Lactose spray dried, Pregelatinized starch, Croscarmellose sodium, Colloidal anhydrous silica, Magnesium stearate, Povidone and Opadry pink 03B54819
Colours: Iron Oxide Red, Iron Oxide Yellow and Titanium Dioxide

DESCRIPTION:

For 120mg: Peach coloured, oblong shaped, biconvex film coated tablets with both sides plain. The tablets are approximately 15.5 mm in length and 7 mm in breadth.
For 180mg: Peach coloured, oblong shaped, biconvex film coated tablets with both sides plain. The tablets are approximately 18 mm in length and 8 mm in breadth.


DOSAGE FORM: Film Coated Tablets

PRESENTATION: 10's, 30's and 50's Tablets (1 x 10's, 3 x 10's & 5 x 10's Alu/PVC/PVdC Blister Strips)

PROPERTIES:

Fexofenadine, a pharmacologically active metabolite of Terfenadine, is a non-sedating antihistamine with selective peripheral H1 receptor antagonist with the chemical name (±)-4-[1 hydroxy-4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-butyl]- , -dimethyl benzeneacetic acid hydrochloride. The molecular formula is C₂₆H₃₉NO₄HCL and its molecular weight is 538.13.

STRUCTURAL FORMULA:



CLINICAL PHARMACOLOGY:

PHARMACODYNAMICS:

Pharmacotherapeutic group: Antihistamines for systemic use, ATC Code: R06AX26
Mechanism of Action
Fexofenadine hydrochloride is a non-sedating H₁ antihistamine. Fexofenadine is a pharmacologically active metabolite of Terfenadine.
Efficacy and Safety
Single and twice daily doses of Fexofenadine hydrochloride provides antihistaminic effects beginning within one hour, achieving maximum at 6 hours and lasting for 24 hours. No tolerance to the effects after 28 days of dosing. Doses of at least 130 mg is required to achieve a consistent effect that is maintained over a 24-hour period.

PHARMACOKINETICS:

Absorption
Fexofenadine hydrochloride is rapidly absorbed into the body following oral administration, with T_{max} occurring at approximately 1-3 hours post dose. The mean C_{max} value was approximately 494 ng/ml following the administration of a180 mg dose once daily.
Distribution
Fexofenadine is 60-70% plasma protein bound.
Biotransformation and elimination
Fexofenadine undergoes negligible metabolism (hepatic or non-hepatic), as it was the only major compound identified in urine and faeces of animals and man. The plasma concentration profiles of fexofenadine follow a bi-exponential decline with a terminal elimination half-life ranging from 11 to 15 hours after multiple dosing. The single and multiple dose pharmacokinetics of fexofenadine are linear for oral doses up to 120 mg BID. A dose of 240 mg BID produced slightly greater than proportional increase (8.8%) in steady state area under the curve, indicating that fexofenadine pharmacokinetics are practically linear at these doses between 40 mg and 240 mg taken daily. The major route of elimination is believed to be via biliary excretion while up to 10% of ingested dose is excreted unchanged through the urine.

INDICATIONS:

For 120mg: Fexofenadine is indicated for the relief of symptoms associated with allergic rhinitis in adults and children 12 years of age and older. Symptoms treated effectively include sneezing, rhinorrhoea, itchy nose/palate/throat itchy/watery red eyes.
For 180mg: Fexofenadine is indicated for the relief of symptoms associated with chronic idiopathic urticaria and allergic rhinitis in adults and children 12 years of age and older. Symptoms treated effectively include sneezing, rhinorrhoea, itchy nose/ palate/ throat, itchy/ watery red eyes.

DOSAGE AND ADMINISTRATION:

For 120 mg:
Adults and Children 12 years and older
The recommended dose is one tablet once daily with water.
Children under 12 years of age
The efficacy and safety of Fexofenadine hydrochloride has not been studied in children under 12.
Special risk group
It is reported that for special risk groups (elderly, renally or hepatically impaired patients), it is not necessary to adjust the dose of Fexofenadine hydrochloride in these patients.
For 180 mg:
Seasonal Allergic Rhinitis and Chronic Idiopathic Urticaria
Adults and Children: 12 years and older – The recommended dose of Fexofenadine tablets is 180 mg once daily with water. A dose of 60 mg once daily is recommended as the starting dose in patients with decreased renal function.

Fexo 120/180mg PIL

Back

ROUTE OF ADMINISTRATION: Oral

CONTRAINDICATIONS:

Hypersensitivity to the active substance or to any of the excipients listed in this leaflet.

WARNINGS AND PRECAUTIONS:

As with most new medicinal products there is only limited data in the elderly and renally or hepatically impaired patients. Fexofenadine hydrochloride should be administered with care in these special groups. Patients with a history of or ongoing cardiovascular disease should be warned that, antihistamines as a medicine class have been associated with the adverse reactions, tachycardia, and palpitations.

INTERACTIONS WITH OTHER MEDICATIONS:

Fexofenadine does not undergo hepatic biotransformation and therefore will not interact with other medicinal products through hepatic mechanisms. Co-administration of Fexofenadine hydrochloride with Erythromycin or Ketoconazole results in a 2-3 times increase in the level of Fexofenadine in plasma. The changes were not accompanied by any effects on the QT interval and were not associated with any increase in adverse reactions compared to the medicinal products given singly. No interaction between Fexofenadine and Omeprazole was observed. However, the administration of an antacid containing Aluminium and Magnesium hydroxide gels 15 minutes prior to Fexofenadine hydrochloride reduces bioavailability, most likely due to binding in the gastrointestinal tract. It is advisable to leave 2 hours between administration of Fexofenadine hydrochloride and Aluminium and Magnesium hydroxide containing antacids. Fruit juices such as grapefruit, orange and apple may reduce the bioavailability and exposure to Fexofenadine.

PREGNANCY AND LACTATION:

Pregnancy

There are no adequate data from the use of Fexofenadine hydrochloride in pregnant women. Fexofenadine hydrochloride should not be used during pregnancy unless clearly necessary.

Lactation

There are no data on the content of human milk after administering Fexofenadine hydrochloride. However, when Terfenadine was administered to nursing mothers Fexofenadine was found to cross into human breast milk. Therefore, Fexofenadine hydrochloride is not recommended for mothers breastfeeding their babies.

Fertility

No human data on the effect of Fexofenadine hydrochloride on fertility are available. In mice, there was no effect on fertility with Fexofenadine hydrochloride treatments.

SIDE EFFECTS:

In adults, the following undesirable effects have been reported in clinical trials, with an incidence similar to that observed with placebo:
Nervous system disorders
Common: headache, drowsiness, dizziness
Gastrointestinal disorders
Common: nausea
General disorders and administration site conditions
Uncommon: fatigue
In adults the following undesirable effects have been reported in post-marketing surveillance. The frequency with which they occur is not known (cannot be estimated from available data):
Immune system disorders: Hypersensitivity reactions with manifestations such as angioedema, chest tightness, dyspnoea, flushing and systemic anaphylaxis.
Psychiatric disorders: insomnia, nervousness, sleep disorders or nightmares/ excessive dreaming (paroniria)
Cardiac disorders: tachycardia and, palpitations
Gastrointestinal disorders: diarrhoea
Skin and subcutaneous tissue disorders: rash, urticaria, pruritus

SYMPTOMS AND TREATMENT OF OVERDOSE:

Dizziness, drowsiness, fatigue and dry mouth have been reported with overdose of Fexofenadine hydrochloride. Single doses up to 800 mg and doses up to 690 mg twice daily for 1 month or 240 mg once daily for 1 year have been administered without the development of clinically significant adverse reactions. The maximum tolerated dose of Fexofenadine hydrochloride has not been established. Standard measures should be considered to remove any unabsorbed medicinal product. Symptomatic and supportive treatment is recommended. Haemodialysis does not effectively remove Fexofenadine hydrochloride from blood.

EFFECT ON ABILITY TO DRIVE AND USE MACHINES:

On the basis of the pharmacodynamic profile and reported adverse reactions it is unlikely that Fexofenadine hydrochloride tablets will produce an effect on the ability to drive or use machines. Fexofenadine hydrochloride has been shown to have no significant effects on central nervous system function. This means that patients may drive or perform tasks that require concentration. However, in order to identify sensitive people who have an unusual reaction to medicinal products, it is advisable to check the individual response before driving or performing complicated tasks.

STORAGE:

Store below 30°C. Store in original package.

SHELF LIFE:

48 months

MANUFACTURER:

Ind-Swift Laboratories Limited
Off- NH-21, Village Jawaharpur,
Tehsil Derabassi, Distt. S.A.S Nagar
(Mohali) Punjab (India) Pin -140507

PRODUCT REGISTRATION HOLDER:

SYNERRV SDN. BHD.,
SO-29-2, Menara 1, KL Eco City,
Jalan Bangsar, Kg. Haji Abdullah Hukum,
59200 Bangsar, Wilayah Persekutuan Kuala Lumpur

LEAFLET REVISION DATE: September 2025

Black

Dimension: 130 x 250mm
Paper should be Maplitho (60 GSM)

Item Code No.	2000XXXXX		Revision No.:	02	
			Prev. Aw. No.:	200012940	
Packaging Development	Marketing	Regulatory Affairs	Production	Quality Control	Quality Assurance

Pharmacode	No.
Front 	4533
Back 	4534