
OTTOZOLE EAR DROPS (CLOTRIMAZOLE 1% W/V)

Name and Strength of Active Substance(s)

Clotrimazole..... 1% w/v

Preservatives:

Sodium Methyl Paraben 0.05% w/v

Propyl Paraben 0.01% w/v

Product Description

Clear, colourless viscous liquid.

Pharmacodynamics

Clotrimazole is an imidazole antifungal agent and is effective against a wide variety of fungi including. Epidermophyton, Microsporum and Trichophyton spp, Coccidioides immitis, Histoplasma capsulatum, Aspergillus and Candida spp. Other fungi responding to Clotrimazole include Cryptococcus neoformans, Malassezia furfur [Pityrosporum orbiculare], Paracoccidioides brasiliensis, Sporothrix schenckii, Psedallescheria, Medurella and and Phialophora spp.

Clotrimazole also has antibacterial action and some strains of Staphylococcus aureus and Streptococcus Pyogenes have been reported to be sensitive.

Clotrimazole is believed to exert its activity on the cell membrane of the fungus. It appears that the drug inhibits protein synthesis and so influences the utilisations and the incorporation of the acylated amino acids essential for fungi.

Pharmacokinetics

Clotrimazole appears to be well absorbed in humans following oral administration and is eliminated mainly as inactive metabolites. Following topical and vaginal administration, however, clotrimazole appears to be minimally absorbed. Six hours after the application of radioactive clotrimazole 1% cream and 1 % solution onto to intact acutely inflamed skin, the concentration of clotrimazole varied from 100 mcg / cm³ in the stratum corneum to 0.5 to 1 mcg / cm³ in the stratum reticulare, and 0.1 mcg / cm³ in the subcutis. No measurable amount of radioactivity [<0.0001 mcg/ml] was found in the serum within 48 hours after application under occlusive dressing of 0.5ml of the solution or 0.8 g of the cream. Only 0.5% or less of the applied radioactivity was excreted in the urine.

Indications

- . Otomycosis
- . Concomitant therapy with Antibiotic and Corticosteroid Ear Drops.
- . Prophylaxis for swimmers.

Recommended Dose

4 -5 drops are instilled in the ear 3-4 times daily or as directed by the physician.

Route of Administration

Auricular (Otic)

Contraindications

Hypersensitivity to clotrimazole.

Warning and Precautions

Perforated eardrum.

Interactions with Other Medicaments

No known drug interaction.

Pregnancy & Lactation**Pregnancy:**

There is a limited amount of data from the use of clotrimazole in pregnant women. Animal studies with clotrimazole have shown reproductive toxicity at high oral doses. At the low systemic exposures of clotrimazole following topical treatment, harmful effects with respect to reproductive toxicity are not predicted. Clotrimazole can be used during pregnancy, but only under the supervision of a physician or midwife.

Lactation:

Available pharmacodynamic/toxicological data in animals have shown excretion of clotrimazole/metabolites in milk after intravenous administration. A risk to the suckling child cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from clotrimazole therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

Adverse Effects / Undesirable Effects

Local reactions such as irritation and burning sensations rarely occurs subject to application.

Overdose and Treatment

No risk of acute intoxication is seen as it is unlikely to occur following a single dermal application of an overdose (application over a large area under conditions favourable to absorption) or inadvertent oral ingestion. There is no specific antidote.

However, in the event of accidental oral ingestion, routine measures such as gastric lavage should be performed only if clinical symptoms of overdose become apparent (e.g. dizziness, nausea or vomiting). Gastric lavage should be carried out only if the airway can be protected adequately.

Effects on Ability to Drive and Use Machine

Clotrimazole has no or negligible influence on the ability to drive or use machines.

Storage Condition

Keep container tightly closed. Store below 30°C. After opening, store below 30°C and discard any remaining solution after 7 days.

Dosage Forms and Packaging Available

10ml bottle. The filled bottle is packed in a preprinted carton with package insert.

Registration Number:

MAL

Manufactured by & Product Registration Holder:

KCK Pharmaceutical Industries Sdn. Bhd.

Plot 61, Bayan Lepas Industrial Park Phase 4,
Lintang Bayan Lepas 1, 11900 Bayan Lepas,
Pulau Pinang, Malaysia

Date of Revision:

10 February 2022